

ERRATUM

## Erratum to: Anesthetic Selection and Disease-Free Survival Following Optimal Primary Cytoreductive Surgery for Stage III Epithelial Ovarian Cancer

Kevin M. Elias, MD<sup>1,2,3,4,5</sup>, Stephanie Kang, BS<sup>2</sup>, Xiaoxia Liu, MS<sup>2</sup>, Neil S. Horowitz, MD<sup>1,4,5</sup>, Ross S. Berkowitz, MD<sup>1,4,5</sup>, and Gyorgy Frendl, MD, PhD, FCCM<sup>2,3,5</sup>

<sup>1</sup>Division of Gynecologic Oncology, Department of Obstetrics and Gynecology and Reproductive Biology, Brigham and Women's Hospital, Boston, MA; <sup>2</sup>Surgical ICU Translational Research Center, Brigham and Women's Hospital, Boston, MA; <sup>3</sup>Department of Anesthesiology, Perioperative and Pain Medicine, Brigham and Women's Hospital, Boston, MA; <sup>4</sup>Dana-Farber Cancer Institute, Boston, MA; <sup>5</sup>Harvard Medical School, Boston, MA

**ERRATUM TO: ANN SURG ONCOL**  
**DOI: 10.1245/S10434-014-4112-9**

Table 4 in the original article contains some errors. Following is the corrected table:

**TABLE 4** Cox's proportional hazards model of selected parameters and relationship to disease free survival

Parameter	Parameter estimate	Hazard ratio	95% hazard ratio confidence limits		<i>p</i> value
Age	−0.00718	0.993	0.974	1.012	0.45
BMI	0.02556	1.026	0.991	1.062	0.15
Race	0.42777	1.534	0.977	2.407	0.06
Anesthesia					
General alone		1.00 (ref)			
General and epidural	−0.30690	0.736	0.491	1.103	0.14
Carcinoma	1.46396	4.323	1.950	9.585	<0.001
Cancer stage					
IIIa	−1.05845	0.347	0.123	0.976	0.04
IIIb	−0.28471	0.752	0.388	1.457	0.40
IIIc		1.00 (ref)			
Estimated blood loss	−0.0001034	1.000	0.999	1.000	0.62
RBC transfusion	0.52880	1.697	0.995	2.894	0.05
Length of case	−0.12754	0.880	0.690	1.123	0.30

The online version of the original article can be found under doi:[10.1245/s10434-014-4112-9](https://doi.org/10.1245/s10434-014-4112-9).

© Society of Surgical Oncology 2015

Published Online: 29 January 2015

K. M. Elias, MD  
e-mail: kelias1@partners.org

**TABLE 4** continued

Parameter	Parameter estimate	Hazard ratio	95% hazard ratio confidence limits		<i>p</i> value
Volatile anesthetic					
Sevoflurane		1.00 (ref)			
Desflurane	-0.57364	0.563	0.330	0.962	0.04
Level of surgical debulking					
No residual		1.00 (ref)			
<5 mm residual	0.57596	1.779	1.166	2.713	0.01
5mm-1 cm residual	1.12904	3.093	1.727	5.538	<0.001