



ASO VISUAL ABSTRACT

ASO Visual Abstract: Association of Tumor Molecular Subtype and Stage with Breast and Axillary Pathologic Complete Response After Neoadjuvant Chemotherapy for Breast Cancer

Sara P. Myers, MD, PhD¹, Gillian M. Ahrendt, BS¹, Joanna S. Lee, MD², Jennifer Steiman, MD², Atilla Soran, MD, MPH², Ronald R. Johnson, MD², Priscilla F. McAuliffe, MD, PhD², and Emilia J. Diego, MD²

¹Department of Surgery, University of Pittsburgh School of Medicine, Pittsburgh, PA; ²Division of Surgical Oncology, Department of Surgery, University of Pittsburgh College of Medicine, Magee Women's Hospital of UPMC, Pittsburgh, PA

Our retrospective study (<https://doi.org/10.1245/s10434-021-10195-8>) postulated that breast cancer molecular phenotype and stage were related to axillary pathologic complete response. Low clinical T stage and hormone-receptor negativity were associated with pathologically negative axilla, indicating that de-escalation of axillary staging may be safe for some.

Publisher's Note Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

DISCLOSURE SPM, GMA, JSL, JS, AS, RRJ, PFM, and EJD have no relevant disclosures or conflicts of interest to disclose.

