



ASO VISUAL ABSTRACT

ASO Visual Abstract: How to Safely Perform Repeat Sentinel Node Biopsy After a Previous Mastectomy—Technical Features and Oncological Outcomes

Elisa Vicini, MD¹, Maria Cristina Leonardi, MD², Sabrina Kahler Ribeiro Fontana, MD, PhD¹, Eleonora Pagan, PhD³, Vincenzo Bagnardi, PhD³, Laura Gilardi, MD⁴, Anna Cardillo, MD⁵, Paola Rafaniello Raviele, MD⁶, Manuela Sargenti, MD¹, Consuelo Morigi, MD¹, Mattia Intra, MD, PhD¹, Paolo Veronesi, MD, PhD^{1,7}, and Viviana Galimberti, MD¹

¹Division of Breast Surgery, European Institute of Oncology IRCCS, Milan, Italy; ²Division of Radiation Oncology, European Institute of Oncology, IRCCS, Milan, Italy; ³Department of Statistics and Quantitative Methods, University of Milan-Bicocca, Milan, Italy; ⁴Division of Nuclear Medicine, European Institute of Oncology, IRCCS, Milan, Italy; ⁵Division of Medical Senology, European Institute of Oncology, IRCCS, Milan, Italy; ⁶Division of Pathology, IEO, European Institute of Oncology IRCCS, Milan, Italy; ⁷Department of Oncology and Hematology-Oncology, University of Milan, Milan, Italy

Repeat sentinel node biopsy after mastectomy is a nonstandard procedure. This practice can represent a safe option to detect and remove occult axillary disease in order

to achieve better local control and could influence the choice of adjuvant treatment (<https://doi.org/10.1245/s10434-021-10986-z>).

