POLYMORPHISM OF MICROSATELLITE SEQUENCE WITHIN ABC TRANSPORTER GENES IN PHYTOPATHOGENIC FUNGUS, *MAGNAPORTHE GRISEA*

Lin Liu¹, Chengyun Li^{1,*}, Jing Yang¹, Jinbin Li², Yuan Su¹, Yunyue Wang¹, Yong Xie¹, Youyong Zhu¹

¹ Key Laboratory for Agricultural Biodiversity and Pest Management of China Education Ministry, Plant Protection College, Yunnan Agricultural University, Kunming, 650201, China

² Plant Protection Research Institute of Yunnan Academy of Agricultural Sciences, Kunming, 650205, China

* Corresponding author, Address: Chengyun Li, Key Laboratory for Agricultural Biodiversity and Pest Management of China Education Ministry, Yunnan Agricultural University, Hei Longtan, Kunming, 650201, P. R. China, Tel: +86-871-5227552, Fax: 86-871-5227552, Email: li.chengyun@gmail.com

Abstract: Thirteen polyporphic microsatellite markers suitable for population genetic structure analysis and ABC transporter and signal transduction coding genes variation measurement were developed for rice blast fungus, *Magnaporthe grisea*. Polymorphism was evaluated by using forty-six isolates collected from diverse geographical locations and rice varieties. Preliminary results indicated that each locus resolved multiple alleles ranging from two to ten. There results showed that these SSR-containing genes are also polymorphic in the natural population.

Keywords: *Magnaporthe grisea*, ABC transporter, microsatellite

1. INTRODUCTION

Rice blast disease, caused by *Magnaporthe grisea*, is the devastated disease of cultivated rice in most rice-growing regions worldwide. The

fungus exhibits a high level of pathotype variation. Potential mechanisms contributing to this variation include mutation, migration, parasexual recombination or an as yet unobserved sexual stage in the field (Ou, 1985). Disease management strategies would greatly benefit from an increased understanding of the amount and distribution of genetic diversity in this pathogen. The completion of the fungus genome sequence project has made it possible to determine not only the total number of genes, but also the exact number of genes of a particular type and analysis their structure and function in details (Ou, 1985; Zeigler et al., 1997). As a consequence, we now know exactly how many regulatory gene are encoded by the blast fungus genome, and how many genes contain simple sequence repeats (SSRs) within protein coding regions. Trinucleotide repeats are clustered in regulatory genes in *Saccaromyces cerevisiae* (Young et al., 2000) and rice blast fungus (Li et al., 2005), but all these SSRs are structurally and functionally polymorphisms, are still unknown.

Microsatellites are founded in both eukaryotes and prokaryotes. It nonrandomly distributes either in expressed sequence tags (ESTs) and genes, including protein-coding, 3'-UTRs and 5'-UTRs, or in introns. The consequences of SSRs repeat-number changes are different in those regions of both prokaryotes and eukaryotes. For example, 14% of protein-coding regions of all known proteins in eukaryotes was proved to containing repeated sequences, and it is three times higher abundance of repeats than in prokaryotes (Marcotte et al., 1998). Characterized with relatively rapid and inexpensive, microsatellites are favored for genetic research, it was not only applied to polymorphic resolve within species but also commonly used to identify specific chromosomal regions consistently across populations.

Genes involved in ABC transporters play a key role in development and pathogenicity of fungal pathogens. The ATP-binding cassette (ABC) superfamily of active transporters is composed of about 50 functionally diverse prokaryotic and eukaryotic transmembrane proteins (Higgins, 1992; Michaelis and Berkower, 1995). The ABC transporters not only carry a variety of substrates into or out of the cell, but also are involved in intracellular compartmental transport. These proteins utilize energy derived from the hydrolysis of ATP to transport the substrate across the membrane against a concentration gradient.

The previous work showed that microsatellite sequences, especially trinucleotide repeats are richness in protein kinase and ABC transporter coding genes of fungus (Keleher et al., 1992). The objective of this study was to determine the polymorphism of these microsatellite loci by PCR assay of loci among natural population in *M. grisea*.

2. MATERIALS AND METHOD

The DNA sequence, a database of known and predicted open reading frames (ORF) of eukaryotic ABC transporters were obtained from the *Maganaporthe grisea* genome database World Wide Web site: <http://www.genome.wi.mit.edu/annotation/fungi/magnaporthe/> on July 14, 2005, and was made sure by *Maganaporthe grisea* genome database World Wide Web site: http://www.broad.mit.edu/annotation/genome/magnaporthe_grisea/ on May 12, 2006. We used the program software tandem repeats finder (TRF) written by Benson (Benson, 1999) with the following options: minimum size = 15 bp, 80% matches (namely number of matched bases between two repetitive elements is 80%) and abundance was removed.

Polymorphic loci were detected by screening a subset of 46 isolates of *M. grisea* collected from different regions (including *japonica*, *indica* rice grown regions) and various rice varieties of Yunnan Province, China. The genomic DNA were extracted from mycelia using a simple extraction protocol (Sweigard et al., 1990). Primers were designed for DNA sequence with microsatellite motifs using PRIMER3 (Rozen and skaletsky, 2000) software and synthesized by Invitrogen Biotechnology Co. Ltd. Shanghai, China.

PCR amplifications were performed in 20 μ L volumes containing 1 $\times$ PCR buffer (10 mM Tris-HCl pH 8.5, 50 mM KCl, 1.5 mM MgCl₂, and 0.001% gelatin), 125 μ M each dNTP, 5 pmol of each primer, and 0.5 U of *Taq* DNA polymerase (Sino-American Biotechnology Co., Beijing). Approximately 50 ng of genomic DNA was used for each reaction. Amplification were performed in a Eppendoff PCR thermal Mastercycler with the cycling parameters; 2 min and 30 sec at 94°C, 35 cycles of 30 sec at 94°C, 1 min at 55°C and 1 min at 72°C followed by a final extension for 10 min at 72°C. In initial experiments, amplified fragments were visualized by electrophoreses in 1.5% agarose gels stained with ethidium bromide. Those loci appeared polymorphic were further examined by 8% polyacrylamide gel to determine the product size of the PCR product and number of alleles per locus. Fragment size of PCR products were estimated on Bio-Imaging System E5000.

3. RESULTS AND DISCUSSION

Thirteen of the fifteen polymorphic loci produced amplicons from a majority of 46 isolates, and displayed two to ten alleles (Table 1). Observed

Table 1. Polymorphisms of SSRs in ABC transporter genes in *M. grisea*. I, Shannon's information index, Ho, expected homozygosity*, Hg, expected heterozygosity*

	Primer(F'-3')	Primer (R'-3')	Gene name	Estimate product size	Super ranging	I	Motif	Repeat No.	Product size range	H _o	H _g
SMS1	ACAAAGCCAGTCGCAGTCAC	CTAACCCGTACGCTTCTTC	MGG_00937.5	250	5.194	0.6902	CAGCAA	3	206-216	0.6092	0.3908
SMS2	CATTGCCCTCGATCGTTTC	TGTTGAGCCACTCGATATGC	MGG_02572.5	250	5.193	1.8668	AAG	6	235-263	0.1887	0.8113
SMS3	GGCCGTACGAGGACTATGAC	TGCGTTTCGGGTTTGATTC	MGG_13490.5	282	5.134	2.0419	GTTGGG	3	245-304	0.1409	0.8591
SMS4	TGCATCCAGGGTAACAGTGA	GTTCGAGCAAGAAGCCTGTC	MGG_05009.5	290	5.175	1.0438	GGTAGC	4	234-324	0.4639	0.5361
SMS5	CCCTGATAGTCGCCCTCAIA	GATCCGGACCAAGCTTGAGTA	MGG_06707.5	254	5.186	1.7759	CGA	6	243-275	0.1906	0.8094
SMS6	CCGACATTGTTCTCGACCTC	ATCCGAACTGGGCTGAACAC	MGG_06939.5	275	5.186	1.5362	CGCCAT	3	293-331	0.2145	0.7855
SMS7	GAGCTGCTGACGTTTGAGG	TCATGCCCTAACCTTTTTC	MGG_07375.5	282	5.191	1.6115	GCAGCT	3	281-308	0.2317	0.7683
SMS8	AGCCTGCACACTACACCAAA	CGGGTAAGCTTTTCCATCAA	MGG_07848.5	256	5.183	1.4919	CTC	5	265-339	0.2480	0.7520
SMS9	ATCAACCCGCAAGACCAAC	ATGATCTGTGAGCCCTGAC	MGG_08309.5	296	5.195	2.0016	GGC	5	328-355	0.1345	0.8657
SMS10	CGTTCATACGAGCGTTTCA	TACGGGAACCAAGAGAGCAC	MGG_12035.5	260	5.187	1.6648	CAAGGC	3	268-301	0.2059	0.7941
SMS11	ATCGTGGGTTGATCGAGAG	GGACCTCCACCAATTTGATGT	MGG_09931.5	276	5.186	1.9762	AAG	5	248-296	0.1677	0.8323
SMS12	AAGGTGGGCACCTCTTC	CTCTCGGGTTGTAAATGA	MGG_10277.5	273	5.179	1.6606	GCT	5	264-302	0.2099	0.7901
SMS13	GAATCACCAGCGAATGTT	GACTCTGAAGCGTTGGAGGT	MGG_11035.5	266	5.187	1.9533	CTCGT	3	237-293	0.1524	0.8476

* Expected homozygosity and heterozygosity were computed using Levene (1949)

heterozygosity and expected heterozygosity values by software GENEPOP (V1.34), were shown in table 1. The results suggested that genes harbored these SSR sequence are also diversity in isolates used.

4. CONCLUSIONS AND FUTURE WORKS

The high degree of polymorphism in this set of microsatellite markers can be used to analysis of population structure and strain distribution in association with particular commodities and locations, as well as complemented for understanding function of regulatory genes in the fungus. With integration of such information into strategies of the functional genomics, it would facilitate SSR functions Study.

ACKNOWLEDGEMENTS

This work is supported by National Basic Research Program of China (2006CB100202), Education Ministry Foundation (307025) and Doctorial Foundation of Education Ministry of China (20050676001).

REFERENCES

- Benson G. 1999, Tandem repeats finder: a program to analyze DNA sequences. *Nucleic Acids Res* 27, 573-580.
- Cecconi F., Alvarez-Bolado G., Meyer B.I., Roth K.A., Gruss P. 1998, Apaf1 (CED-4 homolog) regulates programmed cell death in mammalian development, *Cell* 94, 727-737.
- Dean Ralph A., Talbot Nicholas J., Ebbole Daniel J. Ebbole, Mark L. Farman, Thomas K. Mitchell, Marc J. Orbach, Michael Thon, Resham Kulkarni, Jin-Rong Xu, Huaqin Pan, Nick D. Read, Yong-Hwan Lee, Ignazio Carbone, Doug Brown, Yeon Yee Oh, Nicole Donofrio, Jun Seop Jeong, Darren M. Soanes, Slavica Djionovic, Elena Kolomlets, Cathryn Rehmeier, Welxi Li, Michael Hardling, Soonok Klm, Marc-Henri Lebrun, Heidi Bohnert, Sean Coughlan, Jonathan Butler, Sarah Calvo, Li-Jun Ma, Robert Nicol, Seth Purcell, Chad Nusbaum, James E. Galagan & Bruce W. Birren 2005, The genome sequence of the rice blast fungus *Magnaporthe grisea*. *Nature* 434, 980-986.
- Edward M. Marcotte, Matteo pellegrini, Todd O. Yeates and David Eisenberg 1998, A census of protein repeats, *J. Mol. Biol.* 293, 151-160.
- Higgins C.F. 1992, ABC transporters: from microorganisms to man. *Annu. Rev. Cell. Biol.* 8, 67-113.
- Keleher C.A., Redd M.J., Schultz J., Carlson M., Johnson A.D. 1992, Ssn6-Tup1 is a general repressor of transcription in yeast, *Cell* 68, 709-719.
- Li Cheng-yun, Li Jinbin, Zhou Xiao-gang, Zhang Shao-song, Dong Ai-rong, Xu Ming-hui 2005, Frequency and Distribution of Microsatellite sequence in Open Reading Frames of Rice Blast Fungus, *Magnaporthe grisea*. *Chinese J. Rice Science* 19(2), 167-173, 2005. (in Chinese with English abstract).

- Michaelis S., Berkower C. 1995, Sequence comparison of yeast ATP-binding cassette proteins. Cold Spring Harbor Symp Quant Biol 60, 291-307.
- Ou S.H. Rice diseases 1985, 2nd Edition. Commonwealth Mycological Institute, Kew, UK, 1985. 380.
- Rozen S., Skaletsky H. 2000, Primer3 on the WWW for general users and for biologist programmers. Methods Mol. Biol. 132, 365-386.
- Saxena K., Gaitatzes, C., Walsh, M.T., Eck, M., Neer, E.J., and Smith, T.F. 1996, Analysis of the physical properties and molecular modeling of Sec13: a WD repeat protein involved in vesicular traffic, Biochemistry 35, 15215-15221.
- Stifani S., Blaumueller C.M., Redhead N.J., Hill R.E., Artavanis Tsakonas S. 1992, Human homologs of a Drosophila Enhancer of split gene product define a novel family of nuclear proteins [published erratum appears in Nat. Genet. Dec2(4), 119-127.
- Swigard, J.A., Orbach, M.J., Valent, B., Chumley, F.G. 1990, A miniprep procedure for isolating genomic DNA from *Magnaporthe grisea*, Fungal. Genet. Newslett 37, 4
- Vaisman N., Tsouladze A., Robzyk K., Ben-Yehuda S., Kupiec M., Kassir Y. 1995 The role of *Saccharomyces cerevisiae* Cdc40p in DNA replication and mitotic spindle formation and/or maintenance, Mol. Gen. Genet 247, 123-136.
- Young E.T., Sloan, J.S. and Riper K.V. 2000, Trinucleotide repeats are clustered in regulatory genes in *Saccharomyces cerevisiae*. Genetics 154, 1053-1068.
- Zeigler, R.S., Scott, R.P., Leung, H., Bordeos, A.A., Kumar, J., and Nelson, R.J. 1997, Evidence of parasexual exchange of DNA in the rice blast fungus challenges its exclusive clonality. Phytopathology 87, 284-294.