ON EXTENDED RATE OF CONVERGENCE RESULTS FOR THE INVARIANCE PRINCIPLE

BY C. C. HEYDE

Australian National University

Let $\{X_i, i = 1, 2, 3, \dots\}$ be a sequence of independent and identically distributed random variables for which $EX_i = 0$, $\text{var } X_i = \sigma^2$, $E|X_i|^{2+a} = b < \infty$, $a > 0$. $C[0, 1]$ is the space of continuous real-valued functions on the interval $[0, 1]$ with the sup norm topology. We define a “random broken line” as follows:

$$\begin{aligned}\xi_n(t) &= \sum_{i=1}^k X_i / (\sigma n^{\frac{1}{2}}) && \text{for } t = n^{-1}k \\ \xi_n(t) &= \xi_n(n^{-1}k) + n[\xi_n(n^{-1}(k+1)) - \xi_n(n^{-1}k)](t - n^{-1}k) \\ &&& \text{for } n^{-1}k \leq t \leq n^{-1}(k+1), 1 \leq k \leq n.\end{aligned}$$

THEOREM. *Let g be any uniformly continuous functional, $g: C[0, 1] \rightarrow \text{real line}$, such that there exists a constant $L > 0$ with the property that*

$$|\Pr\{g(\xi(t)) \leq x + h\} - \Pr\{g(\xi(t)) \leq x\}| \leq L|h|$$

where $\xi(t)$ is the standard Wiener process. Then, there exists a constant $A > 0$ such that for all $n > 1$,

$$(1) \quad |\Pr\{g(\xi_n(t)) \leq x\} - \Pr\{g(\xi(t)) \leq x\}| \leq A(\log n)^\lambda n^{-\mu},$$

where $\lambda = (1 + \frac{1}{2}a)/(a+3) (< \frac{1}{2})$, $\mu = \min(a, 1 + \frac{1}{2}a)/2(a+3)$. This result extends that of Theorem 5 of Rosenkrantz [3] where the bound (1) is, in essence, given in the form

$$|\Pr\{g(\xi_n(t)) \leq x\} - \Pr\{g(\xi(t)) \leq x\}| \leq A(\log n)^{\frac{1}{2}} n^{-\mu}$$

subject to the restriction that $a \leq 2$. Only minor modifications of the work of Rosenkrantz and necessary to furnish this extension; we shall just indicate these and refer the reader to [3] for full details of the proof.

The condition $a \leq 2$ was imposed by Rosenkrantz as a consequence of employing a moment inequality of von Bahr and Esseen [1] which is valid for exponent r , $1 \leq r \leq 2$ (note that $2r = 2 + a$). This is used to obtain the vital inequality (25) of [3]. For the case $a > 2$ ($r > 2$), we make use of a result of Dharmadhikari, Fabian and Jogdeo [2] which gives in place of (25),

$$(2) \quad E|z_{nn}|^r \leq C_r n^{\frac{1}{2}} \nu_n,$$

where $C_r = [8(r-1) \max(1, 2^{r-3})]^r$.

Thus, combining (2) with (25) of [3], we can replace the estimate (27) of [3]

Received 14 February 1969.

by the general estimate

$$(3) \quad \Pr (\max_{1 \leq k \leq n} |z_{nk}| > \delta) \leq \max (2, C_r) n^{-a(r)} \delta^{-r}$$

where $a(r) = \min(r - 1, \frac{1}{2}r)$, $r > 1$.

The details of proof of [3] are then followed exactly up to the stage of choosing ϵ_n and δ_n tending to zero at appropriate rates. Instead of the versions chosen in (37), we employ

$$\delta_n = n^{-2a(r)(2r+1)^{-1}} (\beta \log n)^{-(2r+1)^{-1}}, \quad \epsilon_n^2 = 2\beta\delta_n \log n,$$

where $\beta > 1 + a(r)/(2r + 1)$. The result of the theorem then follows as with [3].

REFERENCES

- [1] BAHR, B. VON and ESSEEN, C. G. (1965). Inequalities for the r th absolute moment of a sum of random variables, $1 \leq r \leq 2$. *Ann. Math. Statist.* **36** 299-303.
- [2] DHARMADHIKARI, S. W., FABIAN, V. and JOGDEO, K. (1968). Bounds on the moments of martingales. *Ann. Math. Statist.* **39** 1719-1723.
- [3] ROSENKRANTZ, W. A. (1967). On rates of convergence for the invariance principle. *Trans. Amer. Math. Soc.* **129** 542-552.