Chapter 1

Introduction

Atlantic bluefin tuna, *Thunnus thynnus* (L.), from the Atlantic and Mediterranean, hereinafter referred to as ABFT, has undergone various crises in recent times owing to overfishing, which is why the *International Commission for the Conservation of Atlantic Tunas* (ICCAT) recently adopted highly restrictive conservation measures for the protection of the resource. The ICCAT Standing Committee on Research and Statistics (SCRS) carries out periodic assessments of resources to examine the impact of such conservation measures on the population.

Since the middle of the 20th century fishing pressure on ABFT resources has been growing constantly, causing increasing damage to stocks. New fisheries have been created and fishing effort has increased throughout the Atlantic and Mediterranean to the extent that on two occasions, one in 1993 (Safina 1993) and the other in 2009 (Fromentin et al. 2014) the species' inclusion under Appendix 1 of the *Convention of International Trade in Endangered Species of Wild Fauna and Flora* (CITES) was proposed. Were it to be included it would mean the prohibition of international trade in ABFT. This did not happen on those two occasions, but it was a close call. ICCAT's intervention was crucial on both occasions in redirecting the situation and returning the resource to a situation of apparent sustainability.

Among the most relevant events in recent decades the following can be highlighted:

- Overfishing in the eastern Atlantic in the 1960s (when ICCAT had not yet been founded), which led to a permanent crisis in the traps fishery of the Strait of Gibraltar and the collapse of the northern European fisheries (Cort and Abaunza 2015). From the 1970s purse seine fishing developed rapidly in the Mediterranean, which became the main fishing gear for the catch of ABFT in this sea (ICCAT 2010). Meanwhile, the spawner fisheries of the eastern Atlantic remained at very low levels from the beginning of the 1960s and in 1974 led ICCAT to adopt the first conservation measure (ICCAT 1974), which established a minimum catchable length of age 1 fishes ($W = 6.4$ kg). This measure was not implemented in any eastern stock fisheries until the beginning of the 2000s, which for decades resulted in the illegal catch of millions of fishes below the minimum length.

- At the beginning of the 1980s a second ABFT crisis arose as a result of overfishing, this time in the fisheries of the western stock (a definition of a stock is provided later), a crisis that led to ICCAT's adoption of the first TACs (Total Allowable Catches) for this species, which have been set between 2000 and 2500 t/year since 1982 (Fromentin and Powers 2005). Later, following the recent disappearance of traditional fisheries such as the Sicilian traps, the continued proliferation of purse seine fleets in the Mediterranean and illegal fishing by vessels with flags of convenience had taken ABFT catches to an official figure of 50,000 t in 1995 (ICCAT 2014; Fromentin et al. 2014), in 1998 ICCAT adopted the first TACs for the eastern stock fisheries (32,000 t/year), though the quantity recommended by the scientists at that time was between 15,000 and 25,000 t/year. Nevertheless, following the adoption of this conservation measure there was a complete lack of monitoring among the fleets and the measure was only implemented in the traps.
- Lastly, the Mediterranean overfishing in the 2000s coincided with the start of purse seine fishing for fattening on farms, an activity that brought about the plundering of the species in the Mediterranean (WWF 2008). According to estimations made by the SCRS, the real catch between 1998 and 2006 was between 50,000–61,000 t/year, though the TAC remained at 32,000 t (Fromentin et al. 2014). The scientific reports of 2008 and 2010 pointed to a fall in the spawning biomass of up to 80% below that of the maximum sustainable yield (ICCAT 2012; ICCAT 2014). In view of all this, in 2007 ICCAT adopted a Pluri-Annual Recovery Plan (PARP) throughout all the fisheries of the eastern Atlantic and Mediterranean (ICCAT 2006) involving a very low TAC (for example, 12,900 t in 2012) among other measures such as limiting the minimum weight at catch to 30 kg (ages 1–4), all under the management of the international commission. There were, however, some exceptions to the minimum weight, such as in the Bay of Biscay, Adriatic Sea and the artisanal coastal Mediterranean fishery, where it is just 8 kg (ICCAT 2007).

The PARP has important consequences:

(i) It has led to a considerable reduction in the TAC, which has in turn led to a large fall in the number of fishing vessels in the Mediterranean; (ii) the increase in the minimum size of ABFT (from 10 to 30 kg) has brought about the disappearance of most of the juvenile fisheries; and (iii) it has brought with it strict monitoring of the landings of this species. All of this has meant that in the ten years that the PARP has been in force the spawning stock biomass of ABFT has increased very significantly, something to which the high recruitments in some years may have contributed as a result of favorable environmental conditions (Piccinetti et al. 2013). In the most recent assessments (ICCAT 2012, 2014, 2017) and from different fishing indicators available to the assessment group [the traps in Sardinia (Addis et al. 2012); the Moroccan traps of the Atlantic (Abid et al. 2017); the larvae of the Balearic Sea (Álvarez-Berastegui et al. 2018); the Spanish purse seiners in the Balearic Sea (Gordoa 2014, 2017); the Portuguese traps in the Strait of Gibraltar (Lino et al. 2018); the Japanese long liners in the Atlantic (Kimoto and Itoh 2017); the aerial surveys in the Gulf of Lion (Rouyer et al. 2018); and the Tunisian purse seiners in

the central Mediterranean (Zarrad and Missaoui 2017)] a very considerable recovery of resources has been recorded, which has given rise to a continual increase in quotas (TAC = 36,000 t by 2020) which may bring the PARP to an end.

Since the very first fishing season following the implementation of the PARP (2008), a minimum of 840,000 juvenile specimens (<30 kg) were no longer being caught each year in the central and western Mediterranean (Cort and Martinez 2010). This figure is mainly based on the catches of the 83 vessels of Italy, France and Spain (>40 m) registered at ICCAT, which caught around 100–150 t/year of juveniles according to estimates made by the SCRS (ICCAT 2008). Most of these juveniles have now joined the spawning stock and this is possibly one of the reasons behind the continuing rise of the abundance indices of spawners since then. A simulation made by Belda and Cort (2011) in a scenario in which there were no juvenile catches (<30 kg) by the EU fleet reveals the same trend and quantity of biomass as that reached in the assessment of the ABFT group in 2017 (ICCAT 2017).

Since the adoption and implementation of the PARP the situation has been completely turned around, such that now, under the monitoring of fisheries by member states, the resources of this species are very much recovered (ICCAT 2014, 2017). Moreover, scientific activities on ABFT have since multiplied (Di Natale et al. 2018).

The present study focuses on one of the events described above: overfishing in the 1950s, which was what brought about the disappearance of most of the traps of the Strait of Gibraltar and led to the collapse of the northern European fisheries from the 1960s; the answers provided by science concerning these facts; how the catches of the juvenile fisheries affected this crisis; and how the crisis in catches has now been overcome.

To deal with this matter, and considering the traps as a central theme of the study, the following chapters are described:

- A brief review of the ABFT, which includes some characteristics such as the description of the species, its habitat, stocks, migrations, growth, reproduction, fishing, etc.
- The catch of the species, taking into account the traps of the Strait of Gibraltar, making a tour from their ancient history to the present. This section includes accounts based on archaeological research in the Strait of Gibraltar.
- The trap fisheries from a historical point of view, though given the immense bibliography on this subject and the numerous scientists involved in it, scientific activities related to this fishing gear by Spanish scientists from the twentieth century are cited.
- ABFT fisheries in the eastern Atlantic, the interaction among them and associated research. Special attention is paid to the Bay of Biscay fishery, both from the historical point of view and from fisheries research, as this is now one of the most closely studied ABFT fisheries.
- The latest assessments of ABFT resources carried out by the SCRS group of experts, the most important scientific reference regarding the situation of the stocks of this species.

References

- Abid N, Malouli M, Ben Mhamed A (2017) Standardized CPUE of bluefin tuna, *Thunnus thynnus*, caught by Moroccan Atlantic traps for the period 1986–2016. ICCAT SCRS/2017/038, 7 p
- Addis P, Secci M, Locci I, Cau A, Sabatini A (2012) Analysis of Atlantic bluefin tuna catches from the last *Tonnara* in the Mediterranean Sea: 1993–2010. Fish Res 127(128):133–141. <https://doi.org/10.1016/j.fishres.2012.05.010>
- Álvarez-Berastegui D, Ingram GW Jr, Reglero P, Ferrà C, Alemany F (2018) Changes of bluefin tuna (*Thunnus thynnus*) larvae fishing methods over time in the western Mediterranean, calibration and larval indices updating. Col Vol Sci Pap ICCAT 74(6):2772–2783
- Belda E, Cort JL (2011) Simulation of biomass trends of eastern bluefin tuna (*Thunnus thynnus*) stock under current management regulations. Col Vol Sci Pap ICCAT 66(2):989–994
- Cort JL, Abaunza P (2015) The fall of tuna traps and collapse of the Atlantic bluefin tuna, *Thunnus thynnus* (L.), fisheries of Northern Europe in the 1960s. Rev Fish Sci Aquac 23:4, 346–373. <http://dx.doi.org/10.1080/23308249.2015.1079166>
- Cort JL, Martínez D (2010) Posibles efectos del Plan de Recuperación de atún rojo (*Thunnus thynnus*) en algunas pesquerías españolas. Col Vol Sci Pap ICCAT 65:868–874
- Di Natale A, Lino P, López González JA, Neves dos Santos M, Pagá García A, Piccinetti C, Tensek S (2018) Unusual presence of small bluefin tuna YOY in the Atlantic Ocean and in other areas. Col Vol Sci Pap ICCAT 73(6):3510–3514
- Fromentin JM, Powers J (2005) Atlantic bluefin tuna: population dynamics, ecology, fisheries and management. Fish 6:281–306
- Fromentin JM, Bonhommeau S, Arrizabalaga H, Kell LT (2014) The spectre of uncertainty in management of exploited fish stocks: the illustrative case of Atlantic bluefin tuna. Mar Policy 47:8–14
- Gordoa A (2014) Catch rates and catch size structure of the Balfegó purse seine fleet in Balearic waters from 2000 to 2014; two years of size frequency distribution based on video techniques. Col Vol Sci Pap ICCAT 71(4):1803–1812
- Gordoa A (2017) Updated bluefin tuna CPUE and catch structure from Balfegó purse seine fleet in Balearic waters from 2000 to 2016. Col Vol Sci Pap ICCAT 73(6):2020–2025
- ICCAT (1974) Recomendación 74-1. Talla límite y mortalidad por pesca del atún rojo. <https://www.iccat.int/Documents/Recs/compendiopdf-s/1974-01-s.pdf>
- ICCAT (2006) Informe del período bienal 2006–2007, 1ª parte (2006), vol 1. http://www.iccat.int/Documents/BienRep/REP_ES_06-07_I_1.pdf
- ICCAT (2007) Report for biennial period, 2006–07. Part I (2006), vol 2, 240 p. https://www.iccat.int/Documents/BienRep/REP_EN_06-07_I_2.pdf
- ICCAT (2008) Report for biennial period, 2006–07. Part II (2007), vol. 1, 276 p
- ICCAT (2010) ICCAT manual. Description of species. Chapter 2; 2.1.5 Atlantic bluefin tuna, vol 99. Madrid, ICCAT, pp 93–111. http://iccat.int/Documents/SCRS/Manual/CH2/2_1_5_BFT_ENG.pdf
- ICCAT (2012) Report of the 2012 Atlantic bluefin tuna stock assessment session. Madrid, Spain, 124 p, 4–11 September 2012. http://www.iccat.int/Documents/Meetings/Docs/2012_BFT_ASSESS.pdf
- ICCAT (2014) Report of 2014 Atlantic bluefin tuna stock assessment session. Madrid, Spain, 178 p, 20–28 July 2014. http://iccat.int/Documents/Meetings/Docs/2014_BFT_ASSESS-ENG.pdf
- ICCAT (2017) Report of the 2017 ICCAT bluefin stock assessment meeting. Madrid, Spain, 106 p, 22–27 July 2017. http://iccat.int/Documents/Meetings/Docs/2017_BFT_ASS_REP_ENG.pdf
- Kimoto A, Itoh T (2017) Simple update of the standardized bluefin tuna CPUE of Japanese longline fishery in the Atlantic up to 2016 fishing year. Col Vol Sci Pap ICCAT 73(6):1957–1976
- Lino P, Rosa D, Coelho R (2018) Update on the bluefin tuna catches from the tuna trap fishery off southern Portugal (NE Atlantic) between 1998 and 2016, with a preliminary CPUE standardization. Col Vol Sci Pap ICCAT 74(6):2719–2733

- Piccinetti C, Di Natale A, Arena P (2013) Eastern bluefin tuna (*Thunnus thynnus*, L.) reproduction and reproductive areas and season. Col Vol Sci Pap ICCAT 69(2):891–912
- Rouyer T, Brisset B, Bonhommeau S, Fromentin JM (2018) Update of the abundance index for juvenile fish derived from aerial surveys of bluefin tuna in the western Mediterranean Sea. Col Vol Sci Pap ICCAT 74(6):2887–2902
- Safina C (1993) Bluefin tuna in the West Atlantic: negligent management and the making of an endangered species. Originally published in Conservation Biology, vol 7, pp 229–234. <http://www.seaweb.org/resources/articles/writings/safina2.php>
- WWF (2008) Race for the last bluefin. WWF Mediterranean Project, Zurich, 126 p. https://www.wwf.or.jp/activities/lib/pdf/0811med_tuna_overcapacity.pdf
- Zarrad R, Missaoui H (2017) Update of CPUE bluefin tuna, *Thunnus thynnus* (L. 1758) caught by Tunisian purse seiners in the Central Mediterranean. Col Vol Sci Pap ICCAT 73(6):2183–2187

Open Access This chapter is licensed under the terms of the Creative Commons Attribution 4.0 International License (<http://creativecommons.org/licenses/by/4.0/>), which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons license and indicate if changes were made.

The images or other third party material in this chapter are included in the chapter's Creative Commons license, unless indicated otherwise in a credit line to the material. If material is not included in the chapter's Creative Commons license and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder.