Chapter 21

Comments

1. The MAF method for solution of the boundary value problem (7.1), (7.2) in angles $\Phi < \pi$ was introduced in [52] for general strongly elliptic operators (7.6). We give its complete presentation in Chaps. 7–19 for the most important case of the Helmholtz operator (7.4) with complex $\omega \notin \mathbb{R}$.

This presentation also contains many important details and results, which we publish for the first time. In particular, Lemma 7.1 is new, as well as all results of Chap. 8 which concern distributional solutions.

Theorem 19.2 is new. Theorem 19.7 was formulated in [52, Theorem 6.1], however, its proof was rather sketchy. We give its complete proof in Sect. 19.2 of the present book for the first time.

2. The reduction of boundary problems in angles to the Riemann-Hilbert problem was given for the first time by Malyuzhinets in the case of the impedance (Robin) boundary condition [76, 78].
3. The extension of the MAF method to $\Phi > \pi$ was done in [58, 61].
4. The Helmholtz operator (4.18) with real ω is not strongly elliptic. In this case the geometry of the regions V_l^+ and V^* is quite different from the case $\text{Im } \omega \neq 0$, and the relation of type (16.6) holds only for $t \in (0, 1)$. Therefore, this Riemann-Hilbert problem is ill-posed, as well as the problem (7.1), (7.2). This is the well-known difficulty for the Helmholtz equation in unbounded domains, where some selection rules must be added to make the problem well-posed: either the Limiting absorption principle, or the Limiting Amplitude Principle, or the Sommerfeld radiation condition.

This problem was solved by Merzon [88], who showed that the Limiting Absorption Principle makes this problem well-posed in the case of Dirichlet and Neumann boundary conditions. Namely, he proved that this principle implies the relation (16.6) with $\check{G}(t) = 0$ on the interval $(-1, 0)$.