Ribes hirtum Willd. ex Roem. & Schult.

Grossulariaceae

Syn.: *Ribes frigidum* Kunth

Description: Shrubs, 30–100 cm tall, stems erect, branches glandular hirsute, leaves ovate to cordate, 1–3 cm long, glabrous above, pubescent below, 3-lobed, central lobe usually prominent, margins biserrate, flowers orange-yellow, 2–3 mm across, in nodding axillary racemes, 3–12 cm long, berries red, stipitate glandular.

Distribution: South America (Colombia, Ecuador).

Habitat: High-altitude forests, shrublands, among rocks.

Photo is *R. andicola* (similar species), Petr Sklenář

Stem with a diameter of 10 mm. Xylem to bark ratio 2:1. Rings poorly visible. Approximately 4 rings. Mean ring width 0.5 mm. Analyzed: 1 individual.

Xylem

Transverse section

Vessel diameter 15–20 µm. Vessel outline angular. Vessels lignified, thin- to thick-walled, solitary. Fibers thin- to thick-walled. Parenchyma unrecognizable.

Tangential section

Rays uniseriate and 5–10-seriate, with sheath cells.

Radial section

Perforation plates scalariform. Intervessel pits round. Vessel length 80–120 µm.

Bark

Sieve tubes in horizontal rows, arranged in tangentially layers. With crystal druses.

Pith

Round.

Features according to IAWA and Crivellaro and Schweingruber 2015

Xylem 2 - 9 - 12 - 14 - 22 - 40.1 - 50.1 - 52.2 - 62 - 68 - 75 - 98 - 103 - 105