Erythranthe tilingii (Regel) G.L.Nesom

Phrymaceae

Syn.: *Mimulus langsdorffii* var. *tilingii* (Regel) Greene, *M. lucens* Greene, *M. tilingii* Regel

Description: Perennial herbs, 2–35 cm tall, rhizomatous, forming loose mats, glabrous, or slightly hairy, stems erect, leaves elliptic to ovate, margins irregularly remotely dentate, flowers yellow with throat spotted with red, 2–4 cm long, by 1–3 in upper leaf axils.

Distribution: W North America (from Alaska to California and New Mexico, east to Montana).

Habitat: Streambanks, mossy spring areas, moist alpine meadows.

CC BY 4.0, Miguel Vieira

Rhizome with a diameter of 1.5 mm. Xylem to bark ratio 1:3. Rings indistinct. Analyzed: 1 individual.

Xylem

Transverse section

Only one ring. Vessel diameter 20–25 µm. Vessel outline angular. Vessels solitary. Fibers thin- to thick-walled. Parenchyma pervasive near the center, absent at the periphery.

Tangential section

Rays absent.

Radial section

Perforation plates simple. Intervessel pits round, in alternating position. Vessel length 90–110 µm.

Bark

Phloem small, with small groups of sieve elements. Cortex cells very large (100 µm in diameter). Phellem absent. Crystals absent.

Pith

Round.

Features according to IAWA and Crivellaro and Schweingruber 2015

Xylem 2 - 9 - 12 - 13 - 22 - 40.2 - 50.2 - 52.2 - 62 - 70 - 75 - 79.1 - 117