Thesium tenuissimum Hook. f.

Santalaceae

Description: Subshrubs, 10–25 cm tall, glabrous, stems numerous, angular, ascending, leaves linear-lanceolate, flowers greenish, 3.5–5 mm long, in many-flowered lax racemes.

Distribution: Africa (Guinea, Sierra Leone, Ivory Coast, Nigeria, Cameroon, Bioko Isl.).

Habitat: Montane grasslands, rocky slopes.

Martin Dančák

Root collar with a diameter of 10 mm. Xylem to bark ratio 1:1. Rings recognizable. Maximal rings 3. Mean ring width 0.3 mm. Analyzed: 1 individual.

Xylem

Transverse section

Diffuse-porous. Ring boundary marked by 1–2 rows of flat fibers. Vessel diameter 15–20 µm. Vessel outline angular. Vessels solitary. Fibers thin- to thick-walled. Parenchyma difficult to recognize.

Tangential section

Rays 1–6-seriate.

Radial section

Perforation plates simple. Vessel length 60–80 µm. Ray cells upright.

Bark

Sieve elements in small in groups, radially arranged. Rays dilate. Sclerenchyma in small, irregularly formed groups. Crystals absent.

Pith

Round.

Features according to IAWA and Crivellaro and Schweingruber 2015

Xylem 1 - 5 - 9 - 12 - 13 - 22 - 40.1 - 50.2 - 52.2 - 69 - 75 - 98 - 105