P

Pica

Paul Newman
Department of Medical Psychology and
Neuropsychology, Drake Center, Cincinnati
OH, USA

Definition

Ingesting or craving of substances with no nutritional value.

Current Knowledge

Examples of substances may include dirt, paint chips, clay, leaves, pebbles, etc. Pica is considered a disorder when the behavior is not developmentally appropriate (e.g., >18–24 mo) and persists for at least 1 month. The course of pica is most often in the range of months, rather than years. It is most common in children but has also been studied in pregnant women. The etiology of pica has not empirically been determined, but hypotheses include nutritional deficiencies, cultural factors (ingestion of starch, soil, or clay is regarded as

acceptable by various groups), low socioeconomic status, learned behavior (especially in children with developmental disabilities), or underlying biochemical disorder. Prevalence is unclear, because it is often unrecognized and underreported. Pica is a potentially serious disorder depending on the toxicity of the substance being ingested (such as the ingestion of lead from paint chips).

Cross-References

► [Lead Exposure](#)

References and Readings

- Lacey, E. P. (1990). Broadening the perspective of pica: Literature review. *Public Health Reports*, 105, 29–35.
- McAdam, D. B., Sherman, J. A., Sheldon, J. B., & Napolitano, D. A. (2004). Behavioral interventions to reduce the pica of persons with developmental disabilities. *Behavior Modification*, 28, 45–72.
- Rose, E. A., Porcerelli, J. H., & Neale, A. V. (2000). Pica: Common but commonly missed. *The Journal of the American Board of Family Practice*, 13, 353–358.