H

Hemiplegia

Theslee Joy DePiero

Boston University School of Medicine, Braintree
Rehabilitation Hospital, Boston, MA, USA

Definition

Hemiplegia refers to the paralysis of one side of the body, arm and leg, with or without weakness of the face. Hemiplegia implies a severe or total paralysis, as opposed to hemiparesis, which implies some retained movement.

Causes

Hemiplegia is most commonly caused by lesions of the corticospinal tract in the brain, brain stem, or cervical spinal cord. Depending on the location of the lesion, there may be accompanying abnormalities, such as cognitive or language disorders, visual impairments, or sensory disorders. In lacunar strokes, the hemiplegia involves the face, as well as arm and leg. Dysarthria may be present,

but there are no sensory, cognitive, or language deficits. In the spinal cord, there are almost always other abnormalities associated with hemiplegia, typically sensory deficits.

Types

Hemiplegia may be spastic or flaccid. Often there is initial flaccidity followed by spasticity.

Etiology

Stroke, tumors, trauma, hemorrhages, and congenital.

Cross-References

- ▶ [Corticospinal Tract](#)
- ▶ [Hemiparesis](#)
- ▶ [Lacunar Stroke](#)

References and Readings

- Ropper, A. H., Samuels, M. A., & Klein, J. P. (2014). *Adams and Victor's principals of neurology* (10th ed.). New York: McGraw-Hill.