Spatial and Temporal Variability of Annual Precipitation during 1958-2007 in Loess Plateau, China

Rui Guo, Fengmin Li, Wenying He, Sen Yang, and Guojun Sun*

Ministry of Education Key Laboratory of Arid and Grassland Ecology,
Lanzhou University, China 730000
sungj@lzu.edu.cn

Abstract. The non-parametric Mann-Kendall method and Sliding t -test were applied in this study to analyze the spatial and temporal patterns of trends of the precipitation in the Loess Plateau from 1958 to 2007. The main trends of the precipitation in this region is downward, only four upward time series have not reach the significant level. Most of the abrupt changes happened in northeast of Loess Plateau during 1980s and 1990s. Furthermore, the abrupt changes happened in the northeast portion of the Loess Plateau were relatively later than those in the west portion of the Loess Plateau.

Keywords: Trend analysis, Abrupt change, Mann-Kendall method, Sliding t -test method.

1 Introduction

Loess Plateau is located in arid and semi-arid area of China. It faces too many problems such as low vegetation cover, water scarcity and severe soil erosion and so on. Lots of the problems were affected by the precipitation. The precipitation is also one of the major factors limited agricultural and forestry productivity. Fierce human activities increase the possibility of abrupt changes in climate[1]. It is a meaningful work to find out the regulation of the precipitation in Loess Plateau.

Mann-Kendall trend test is a non-parametric, rand-based method for assessing the significance of trends in time-series data such as water quality, streamflow, temperature, and precipitation[2]. Not only can examine the trends in time-series decline or increase, but also reflect the degree of changes, Mann-Kendall trend test could be a very good description of the characteristics of the time-series trend. Now, the Mann-Kendall trend test has been widely used and tested as an effective method to evaluate a presence of a statistically significant trend in hydrological and climatological time series[2-4]. In the study, we use this method to test trends on precipitation in Loess Plateau from 1958 to 2007. The Sliding t -test[5] was used to detect the abrupt changes in the study.

* Corresponding author.

The objectives of this study are (1) to reveal the spatial and temporal patterns of trends of the precipitation in the Loess Plateau; (2) to analyze the abrupt changes of the precipitation in the Loess Plateau from 1958 to 2007.

2 Study Area and Data Processing

2.1 Loess Plateau

The Loess Plateau is a subsystem of mutually interlinked and feedback system of northwest Gobi desert, middle Loess plateau and east alluvial plain[6]. It is located in the middle reaches of the Yellow River and extends 7 latitudes (35° - 41° N) and 13 longitudes (102° - 114° E), with total area of more than $630,000\text{km}^2$ and a population of 86 million. The plateau area is 1000-1600 m above sea level and surface is covered by an average 100-meter thickness of loess-paleosol layers. The Loess Plateau belongs to the continental monsoon region with annual average temperature of 4.3°C in northwest and 14.3°C in southeast. Annual precipitation ranges from 300mm in northwest to 700mm in southwest[7]. There is much sunshine in the Loess Plateau. The soil erosion rate on the Loess Plateau of China is among the highest in the world. Loess Plateau contains arid and semi-arid regions; the eco-environment is fragile.

2.2 Precipitation Data

The precipitation data were downloaded from China Meteorological Data Sharing Service System. The time series of annual precipitation records at 42 meteorological stations were used in this study from 1958 to 2007. The stations in the Loess Plateau were shown in Fig.1.

Fig. 1. The location of the Loess Plateau in China. The locations of the Meteorological stations are shown by the black triangle.

2.3 Methods

2.3.1 Mann-Kendall Method

The non-parametric Mann-Kendall method was used to assess the significance of a trend, and originated from Mann (1945)[8] and rephrased by Kendall (1948)[9]. The test statistic is calculated as follows:

$$S = \sum_{i=1}^{n-1} \sum_{j=i+1}^n \text{sgn}(x_i - x_j) . \quad (1)$$

Where n is the length of the data set, and x is the data point at time i and j , and

$$\text{sgn}(\theta) = \begin{cases} 1 & \text{for } \theta > 0 \\ 0 & \text{for } \theta = 0 \\ -1 & \text{for } \theta < 0 \end{cases} . \quad (2)$$

For independent, identically distributed random variables with on tied data values:

$$E(S) = 0 . \quad (3)$$

$$\text{Var}(S) = \frac{n(n-1)(2n+5)}{18} = \sigma^2 . \quad (4)$$

When some data values are tied, the correction to $\text{Var}(S)$ is:

$$\text{Var}(S) = \frac{n(n-1)(2n+5) - \sum_{i=1}^n t_i(i)(i-1)(2i+5)}{18} . \quad (5)$$

Where t_i is the number of ties of extent i . The standardized test statistic Z is computed by:

$$Z = \begin{cases} \frac{S-1}{\sigma} & \text{for } S > 0 \\ 0 & \text{for } S = 0 \\ \frac{S+1}{\sigma} & \text{for } S < 0 \end{cases} . \quad (6)$$

The standardized MK statistic Z follows a standard normal distribution with mean of zero and variance of one under the null hypothesis of no trend. A positive Z value indicates an upward trend, whereas a negative one indicates a downward trend. The p

value (probability value p) of the MK statistic S of sample data can be estimated using the normal cumulative distribution function:

$$p = 0.5 - \phi(|Z|) \quad (7)$$

where

$$\phi(|Z|) = \frac{1}{2\pi} \int_0^{|Z|} e^{-\frac{t^2}{2}} dt \quad (8)$$

If the p value is small enough, the trend is quite unlikely to be caused by random sampling. At the significance level of 0.05, if $p \leq 0.05$, then the existing trend is assessed to be statistically significant [10, 11].

2.3.2 Sliding t -Test Method

One moment is set to be a reference point in time series “ x ” with a sample size “ n ”. The two subsequences before and after the reference point were defined to be x_1 and x_2 .

$$t = \frac{\bar{X}_1 - \bar{X}_2}{s \cdot \sqrt{\frac{1}{n_1} + \frac{1}{n_2}}} \quad (9)$$

Where

$$s = \sqrt{\frac{n_1 s_1^2 + n_2 s_2^2}{n_1 + n_2 - 2}} \quad (10)$$

n_1, n_2 ——the sample size of x_1 and x_2 , separately.

$\bar{X}_1, \bar{X}_2$ ——the average of x_1 and x_2 , separately.

s_1^2, s_2^2 ——the variance of x_1 and x_2 , separately.

If $|t_i| < t_\alpha$, there is no significant difference between the two subsequence. Otherwise, there is an abrupt change at the reference point[5].

3 Results and Discussion

3.1 The Spatial Structure of the Precipitation

The linear regression was used to detect the relationship between the average rainfall for this 50 years and longitude, latitude, elevation (Fig.2). The results presented that the

average precipitation was linear positively related to the longitude, but it is not significant. The average precipitation was not the linear relation with the latitude or the elevation. It was suggested that the spatial differentiation of the precipitation was not significant on the Loess Plateau from 1958 to 2007.

Fig. 2. Relationship between the average rainfall and longitude, latitude, elevation in Loess Plateau, the linear models and R^2 also gave in the figure

3.2 Trend Analysis and Abrupt Changes

The non-parametric Mann-Kendall method was used to assess the significance of the trend of precipitation. By this method the significant trends were detected for 42 time series of annual precipitation for different stations in the Loess Plateau (Fig. 3).

There were only 7 among 42 stations showed a significant trend ($p=0.05$), the remaining 35 stations showed no significant trend. Moreover, all these 7 stations with significant trend ($p=0.05$) displayed a downward trend. The four stations with an upward trend showed in Fig.3 were not assessed to be statistically significant. In other words, the standardized MK statistic Z of these four stations is positive, but did not reach significant levels. The standardized MK statistic Z of the remaining 31 stations with no significant trend is negative. It can be easily found from the distribution of the trends for the annual precipitation that no trend and upward trend were mainly in the western portion of Loess Plateau. The downward trend concentrated in the eastern portion of central Loess Plateau. The results also indicated that the precipitation possesses the longitude zonality. This result was similar as Q. Liu et al. [12] that studied the precipitation during 1961-2006 in Yellow River Basin. The precipitation changed

Fig. 3. The trends in annual precipitation during 1958-2007 detected by the Mann-Kendall method in Loess Plateau

Fig. 4. The abrupt change tested by the Sliding t -test for annual precipitation series in Youyu station from 1962 to 2002 ($n_1=n_2=5$)

obviously from west to northeast. This situation may be due to the monsoon, as the precipitation was sensitive to the climatic change. And the influenced by human activities might another reason.

The Sliding t -test method was used to test the abrupt changes. The Youyu station ($112^{\circ}27'E$, $40^{\circ}00'N$) was selected as an example to demonstrate the abrupt change

Table 1. The time of the abrupt change detected by the Sliding t-test. (* $p<0.05$. ** $p<0.01$. The year that the abrupt change happened has been simplified in two figures, e.g., 67 for 1967).

Stations		Sliding t-test			Stations			Sliding t-test			
	n1=n2=5	n1=n2=7	n1=n2=10	n1=n2=11	n1=n2=12		n1=n2=5	n1=n2=7	n1=n2=10	n1=n2=11	n1=n2=12
Yan'an	87*, 92*, 01*					Jingyuan					
Datong			96*			Huanxian		68*	68*, 70*	68*, 69*, 71*	69*, 70*
Hecui	87*					Jingtai	66*				91*
Wutaishan	80*	92*	88*, 89*, 92*, 96*	88*, 89*, 90*, 92*, 95*, 96*	87*, 88*, 89*, 90*, 91*, 92*, 93*, 94*	Weixian		92*	92*	92*, 95*	
Wuzhai						Sannexia	80*, 85*				85*, 89*, 90*
Xingxian	78*, 79*		96*			Changwu					
Yuanning			96*			Luchuan					85*
Lishi	72**	64*, 67*	67*, 90*	90*	90*	Yanan			90*	68*, 69*, 88*	69*, 88*, 89*, 90*, 94*
Taiyuan			96*		69*	Yushe		74*			
Yangquan						Suide	70**		95*, 96*	94*, 95*, 96*	88*, 94*, 95*
Yushe						Hengshan			90*		88*
Xixian	96*					Yulin					
Jiexiu	93*, 94**	93**	89*, 90*, 91*, 92*	88*, 89*, 90*, 91*, 92*, 93*	85*, 88*, 89*, 90*, 91*, 92*, 93*, 94*	Guide	02*				
Linfen				76*	77*	Minhe	79*	79*	79*, 91*	79*, 91*	78*, 79*, 91**
Yangcheng			67*	68*		Xining					91*, 92*, 93*
Liniao		79*	92*			Menyuan					
Huajialing			85*, 86*	85*, 86*, 87*	84*, 85*, 86*, 90*	Yanchi					69*, 70*, 71*, 83*
Kongtong	68*					Guyuan	68*		67*		
Linxia						Haiyuan					
Xifeng		00*				Tongxin					
Yuzhong						Xiji	68*	67*, 68*, 95*	67*, 68*, 92*	68*	69*, 70*, 94*, 95*

Fig. 6. The times of abrupt change detected by the Sliding t -test ($n_1=n_2=10$) in different decades

Moreover, the times of abrupt changes happened in different decades showed in Fig.6. It can be found from the chart that the times of abrupt change increase with the time except from 1971 to 1980. Only one station which located in the west of Loess Plateau happened abrupt change from 1971 to 1980. The abrupt change mainly happened in west of Loess Plateau from 1961 to 1970. However, during 1981-1990 and 1991-2000, the abrupt change was mainly happened in the northwest of Loess Plateau. That is the abrupt changes happened in the northeast portion of the Loess Plateau were relatively later than those in the west portion of the Loess Plateau. Moreover, there were 6 stations among 12 which locate in the northeast happened abrupt change in 1996. It indicated that the northeast of Loess Plateau became more and more sensitive to the climate change since 1980s. The reasons that affected the precipitation in the Loess Plateau need further study on the interaction between the climate change and human activities. The results in this study are significance for the further studies and forecast of climate in Loess Plateau, and also offer the basis for decision-makers.

4 Conclusions

In this study, the Mann-Kendall method and Sliding t -test were used to analyze the precipitation from 1958 to 2007 in Loess Plateau. The results presented as follows:

- (1) The relationship between precipitation and longitude/latitude/elevation is not linear. The precipitation didn't have clear trend with the longitude/latitude/elevation increasing.
- (2) The results of Mann-Kendall method indicated that only 7 stations showed a significant trend, the other 35 stations have no significant trend. Only 4 stations have positive Z values but didn't reach the 95% significant level.
- (3) The results of Sliding t -test showed that there were 31 abrupt changes happening in 18 meteorological stations. Most of the abrupt changes happened in the northeast of Loess Plateau. Moreover, the abrupt changes happened in the northeast

concentrated during 1980s and 1990s. And the abrupt changes happened in the northeast portion of the Loess Plateau were relatively later than those in the west portion of the Loess Plateau.

- (4) The northeast of Loess Plateau was sensitive to climate change recent years, as most of the abrupt changes occurred at here during 1980s and 1990s. More attention should be paid to this region, and the changes in land cover/use could be considered in further studies.

Acknowledgments

This work is financial supported by National Key Basic Research Program (973) project (2007CB106804) and the Construction of an Information Platform/module in Eco-agricultural Assessment and Management (EAM) (2010DFA31450). The authors are grateful to the anonymous referees for their careful corrections and constructive comments on the manuscript. The authors thank the China Meteorological Data Sharing Service System for offering the precipitation data.

References

1. Alley, R.B., Marotzke, J., Nordhaus, W.D., et al.: Abrupt climate change. *Science* 229, 2005–2010 (2003)
2. Yue, S., Pilon, P., Cavadias, G.: Power of the Mann-Kendall and Spearman's rho tests for detecting monotonic trends in hydrological series. *Journal of Hydrology* 259, 254–271 (2002)
3. Douglas, E.M., Vogel, R.M., Kroll, C.N.: Trends in floods and low flows in the United States: impact of spatial correlation. *Journal of Hydrology* 240, 90–105 (2000)
4. Birsan, M.V., Molnar, P., Burlando, P., Pfaundler, M.: Streamflow trends in Switzerland. *Journal of Hydrology* 314, 312–329 (2005)
5. Wei, F.: The statistics and prediction of climate. China Meteorological Press (1999) (in Chinese)
6. He Xiubin, L.Z., Hao, M., Tang, K., Zheng, F.: Down-scale analysis for water scarcity in response to soil–water conservation on Loess Plateau of China. *Agriculture, Ecosystems and Environment* 94, 355–361 (2003)
7. Jie We, J.Z., Tian, J., He, X., Tang, K.: Decoupling soil erosion and human activities on the Chinese Loess Plateau in the 20th century. *Catena* 68, 10–15 (2006)
8. Mann, H.B.: Non-parametric test against trend. *Econometrika* 13, 245–259 (1945)
9. Kendall, M.G.: Rank Correlation Methods. Hafner, New York (1948)
10. Douglas, E.M., Vogel, R.M., Kroll, C.N.: Trends in floods and low flows in the United States: impact of spatial correlation. *Journal of Hydrology* 240, 90–105 (2000)
11. Bouza-Deaño, R., Ternero-Rodríguez, M., Fernández-Espinoza, A.J.: Trend study and assessment of surface water quality in the Ebro River (Spain). *Journal of Hydrology* 361, 227–239 (2008)
12. Liu, Q., Yang, Z.F., Cui, B.S.: Spatial and temporal variability of annual precipitation during 1961–2006 in Yellow River Basin, China. *Journal of Hydrology* 361, 330–338 (2008)