Density Theorems for Intersection Graphs of t -Monotone Curves

Andrew Suk*

Department of Mathematics, Massachusetts Institute of Technology, Cambridge, MA
asuk@math.mit.edu

Abstract. A curve γ in the plane is t -monotone if its interior has at most $t - 1$ vertical tangent points. A family of t -monotone curves F is *simple* if any two members intersect at most once. It is shown that if F is a simple family of n t -monotone curves with at least ϵn^2 intersecting pairs (disjoint pairs), then there exists two subfamilies $F_1, F_2 \subset F$ of size δn each, such that every curve in F_1 intersects (is disjoint to) every curve in F_2 , where δ depends only on ϵ . We apply these results to find pairwise disjoint edges in simple topological graphs.

1 Introduction

Given a collection of objects C in the plane, the *intersection graph* $G(C)$ has vertex set C and two objects are adjacent if and only if they have a nonempty intersection. Intersection graphs have recently received a lot of attention due to its applications in VLSI design [14], map labeling [1,3], graph drawing [9,11,24,25], and graph theory [15]. Over the past several decades, many researchers have shown that intersection graphs of certain geometric objects, such as of segments, chords of a circle, axis parallel rectangles, etc., have very strong properties (see [13,4,23,9]). The aim of this paper, is to establish density-type theorems for intersection graphs of t -monotone curves in the plane.

A *curve* in the plane is the image of a continuous injective function $f : I \rightarrow \mathbb{R}^2$, whose domain is a closed interval $I \subset \mathbb{R}$. For integer $t \geq 1$, a curve γ in the plane is *t -monotone*,¹ if its interior has at most $t - 1$ vertical tangent points. In 2001, Pach and Solymosi proved the following two theorems.

Theorem 1 ([18]). *Let F be a family of n segments in the plane with at least ϵn^2 intersecting pairs. Then there exists a δ that depends only on ϵ , and two subfamilies $F_1, F_2 \subset F$, such that $|F_1|, |F_2| \geq \delta n$, and every segment in F_1 intersects every segment in F_2 .*

Theorem 2 ([18]). *Let F be a family of n segments in the plane with at least ϵn^2 disjoint pairs. Then there exists a δ that depends only on ϵ , and two subfamilies $F_1, F_2 \subset F$, such that $|F_1|, |F_2| \geq \delta n$, and every segment in F_1 is disjoint to every segment in F_2 .*

* Research funded by an NSF Postdoctoral Fellowship.

¹ A 1-monotone curve is often referred to as *x -monotone*. Every t -monotone curve can be decomposed into t 1-monotone curves.

These theorems were later generalized by Alon et al. to semi-algebraic sets in $\mathbb{R}^d$ [2], and by Basu [5] to definable sets belonging to some fixed definable family of sets in an o-minimal structure. In all three papers [18], [2], [5], the authors only considered geometric objects with bounded or fixed “description complexity”. Previously, there were no known generalizations of Theorem 2 to geometric objects with large complexity.

Our main result generalizes Theorems 1 and 2 to t -monotone curves, which can have arbitrarily large complexity. We say that a family of curves F is *simple* if any two members intersect at most once.

Theorem 3. *Let F be a simple family of n t -monotone curves in the plane with at least ϵn^2 intersecting pairs. Then there exists a constant c_t that depends only on t , and two subfamilies $F_1, F_2 \subset F$, such that $|F_1|, |F_2| \geq \epsilon^{c_t} n$, and every curve in F_1 intersects every curve in F_2 .*

Theorem 4. *Let F be a simple family of n t -monotone curves in the plane with at least ϵn^2 disjoint pairs. Then there exists a constant c_t that depends only on t , and two subfamilies $F_1, F_2 \subset F$, such that $|F_1|, |F_2| \geq \epsilon^{c_t} n$, and every curve in F_1 is disjoint to every curve in F_2 .*

Interestingly, Theorem 4 does not hold if one drops the simple condition. That is, there exists a family F of n 1-monotone curves in the plane, with at least $n^2/4$ disjoint pairs, such that for any two subsets $F_1, F_2 \subset F$ of size $\Omega(\log n)$ each, there exists a curve in F_1 that crosses a curve in F_2 .

While Theorem 4 is a new result, Theorem 3 is a special case of a theorem due to Fox, Pach, and Tóth. In [9], Fox, Pach, and Tóth generalized Theorem 1 to families of curves in the plane with the property that any two curves intersect at most a constant number of times. Let us remark that our proof is conceptually simpler.

1.1 Applications to Topological Graphs

A *topological graph* is a graph drawn in the plane such that its vertices are represented by points and its edges are represented by non-self-intersecting arcs connecting the corresponding points. The arcs are allowed to intersect, but they may not intersect vertices except for their endpoints. Furthermore, no two edges are tangent, i.e., if two edges share an interior point, then they must properly cross at that point in common. A topological graph is *simple* if every pair of its edges intersect at most once. Two edges of a topological graph *cross* if their interiors share a point, and are *disjoint* if they neither share a common vertex nor cross.

Over 40 years ago, Conway asked what is the maximum number of edges in a *thrackle*, that is, a simple topological graph with no two disjoint edges. He conjectured that every n -vertex thrackle has at most n edges. Lovász, Pach, and Szegedy [16] were the first to establish a linear bound, proving that all such graphs have at most $2n$ edges. Despite recent improvements by Cairns and Nikolayevsky [6] and Fulek and Pach [11], this conjecture is still open. In the special case that the edges are drawn as 1-monotone curves, Pach and Sterling settled Conway’s conjecture in the affirmative [19].

Determining the maximum number of edges in a simple topological graph with no k pairwise disjoint edges, seems to be a difficult task. Pach and Tóth [22], showed that every simple topological graph with no k pairwise disjoint edges has at most $O(n \log^{4k-8} n)$ edges. They conjectured that for every fixed k , the number of edges in such graphs is at most $O(n)$. A linear bound was obtained by Pach and Töröcsik [20], in the special case that the edges are drawn as 1-monotone curves (see also [25]). As an application of Theorem 4, we improve (for large k) the Pach and Tóth bound, in the special case that the edges are drawn as t -monotone curves (where t is independent of n).

Theorem 5. *Let $G = (V, E)$ be an n -vertex simple topological graph with edges drawn as t -monotone curves. If G does not contain k pairwise disjoint edges, then $|E(G)| \leq n(\log n)^{c'_t \log k}$, where c'_t depends only on t .*

In 2009, Fox and Sudakov [10] showed that all dense simple topological graphs have at least $\Omega(\log^{1+\delta} n)$ pairwise disjoint edges, where $\delta \approx 1/40$. As an immediate Corollary to Theorem 5, we improve this lower bound (to nearly polynomial) in the special case that the edges are drawn as t -monotone curves

Corollary 1. *Let $G = (V, E)$ be an n -vertex simple topological graph with edges drawn as t -monotone curves. If $|E(G)| \geq \epsilon n^2$, then G has at least $n^{\delta / \log \log n}$ pairwise disjoint edges, where δ depends only on ϵ and t .*

We note that Suk recently showed that every complete n -vertex simple topological graph has at least $\Omega(n^{1/3})$ pairwise disjoint edges [24].

2 A Two-Color Theorem

In this section, we will prove the following two-color theorem.

Theorem 6. *Given a family B of n blue t -monotone curves and a family R of n red t -monotone curves in the plane such that $B \cup R$ is simple, there exist a $c'_t > 0$ that depends only on t , and subfamilies $B' \subset B$, $R' \subset R$, such that $|B'|, |R'| \geq n/c'_t$, and either each curve in B' intersects every curve in R' , or each curve in B' is disjoint to every curve in R' .*

In what follows, we will prove a sequence of lemmas that will lead to the proof of Theorem 6. First we need some definitions. Let F be a simple family of curves in the plane. For $\gamma \in F$, the endpoint with the left (right) most x -coordinate we refer to as the *left (right) endpoint* of γ . By a slight perturbation, we can assume that all endpoints have unique x -coordinates, no endpoint of one curve lies on another curve, and no three curves in F have a nonempty intersection. For any simply connected region $\Delta \subset \mathbb{R}^2$, we denote $bd(\Delta)$ as the boundary of Δ . For the rest of the paper, we will assume that all regions are simply connected.

A point q is called a *critical point* of $\gamma \in F$, if q is an endpoint of γ or if γ has a vertical tangent at q . Now given a subset $S \subset F$, the *vertical decomposition* of the

arrangement of S is constructed by subdividing the cells of the arrangement $\mathcal{A}(S) = \bigcup_{\gamma \in S} \gamma$ into trapezoid-like regions $\Delta_1, \Delta_2, \dots, \Delta_s$, by drawing a vertical line in both directions through every intersection point of a pair of curves and through every critical point of a curve in S until it hits a piece of some element in S . We let $\mathcal{T}(S)$ be the vertical decomposition of S , and for simplicity we will call the elements in $\mathcal{T}(S)$ *trapezoids*. See Figure 1. Let $Reg = \bigcup_{S \subset F} \mathcal{T}(S)$ be the set of all trapezoids that can ever appear in the vertical decomposition for some $S \subset F$. For each trapezoid $\Delta \in Reg$, let $D(\Delta)$ be the set of curves in F that intersects the boundary of Δ but does not intersect the interior of Δ . One can easily check that $|D(\Delta)| \leq 4$ (see [17]). Finally, we let $I(\Delta)$ denote the set of curves of F intersecting the interior of Δ .

Fig. 1. Trapezoids Δ_1 and Δ_2 . Notice that $|D(\Delta_1)| = 2$ and $|D(\Delta_2)| = 3$.

Lemma 1. *Given a simple family F of n t -monotone curves and a set P of n points in the plane with no point in P lying on any curve from F , there exists a constant C_1 , subsets $F' \subset F$ and $P' \subset P$ of size $n/(C_1 t \log^2 t)$ each, and a region Δ , such that $P' \subset \Delta$ and no curve in F' intersects the interior of Δ .*

Proof: Let S be a random subset of F by selecting each curve in F independently with probability $p = r/n$. Notice that for any subset $S \subset F$, $|\mathcal{T}(S)| \leq C_2(|S|^2 + |S|t)$ for some constant C_2 , since F is simple. An easy calculation shows that $\mathbb{E}[|S|t] = rt$ and $\mathbb{E}[|S|^2] \leq r^2 + r$, and thus

$$\mathbb{E}[|\mathcal{T}(S)|] \leq C_2(r^2 + r + rt) \leq 3C_2r^2t.$$

For $\Delta \in Reg$, we let $p(\Delta)$ denote the probability that Δ appears in the vertical decomposition of S . Since Δ appears if and only if all curves of $D(\Delta)$ are selected into S and none of $I(\Delta)$ is selected, we have

$$p(\Delta) = p^{|D(\Delta)|}(1-p)^{|I(\Delta)|}.$$

We call a trapezoid $\Delta \in Reg$ *bad* if $|I(\Delta)| \geq n/2$, otherwise it is *good*. Let $Bad = \{\Delta \in Reg : |I(\Delta)| \geq n/2\}$, and let X denote the number of bad trapezoids in the vertical decomposition of S . Since $|D(\Delta)| \leq 4$, and any four curves defines at most $20t$ trapezoids, we have

$$\mathbb{E}[X] = \sum_{\Delta \in Bad} p^{|D(\Delta)|}(1-p)^{|I(\Delta)|} \leq \sum_{1 \leq i \leq 4} 20t \binom{n}{i} \left(\frac{r}{n}\right)^i \left(1 - \frac{r}{n}\right)^{n/2}.$$

For $r = C_3 \log t$, where C_3 is a sufficiently large constant and $t \geq 2$, we have

$$\mathbb{E}[X] \leq 80tr^4 e^{-r/2} = 80t(C_3 \log t)^4 t^{-C_3/2} \leq \frac{1}{3}.$$

Hence

$$\mathbb{E} \left[\frac{1}{9C_2 r^2 t} |\mathcal{T}(S)| + X \right] \leq \frac{2}{3}.$$

By setting $C_1 = 9C_2(C_3)^2$, there exists a sample S such that

$$|\mathcal{T}(S)| \leq 9C_2 r^2 t = 9C_2(C_3 \log t)^2 t = C_1 t \log^2 t$$

and $X = 0$. By the pigeonhole principle, there exists a good trapezoid $\Delta \in \mathcal{T}(S)$ that contains at least $n/(C_1 t \log^2 t)$ points from P , and at least $n/2$ curves from F do not intersect the interior of Δ . This completes the proof. $\square$

Lemma 2. *Given a family R of n red t -monotone curves and a family B of n blue t -monotone curves in the plane such that $R \cup B$ is simple, there exists a constant C_4 , subsets $R' \subset R, B' \subset B$ of size $n/(C_4 t \log^2 t)^4$ each, and regions $\Delta_{bl}, \Delta_{br}, \Delta_{rl}, \Delta_{rr} \subset \mathbb{R}^2$, such that*

1. *the left endpoint of each blue curve in B' lies inside Δ_{bl} ,*
2. *the right endpoint of each blue curve in B' lies inside Δ_{br} ,*
3. *the left endpoint of each red curve in R' lies inside Δ_{rl} ,*
4. *the right endpoint of each red curve in R' lies inside Δ_{rr} ,*
5. *$\Delta_{bl} \cup \Delta_{br}$ is disjoint to $\Delta_{rl} \cup \Delta_{rr}$, and*
6. *no curve in R' intersects $\Delta_{bl} \cup \Delta_{br}$, and no curve in B' intersects $\Delta_{rl} \cup \Delta_{rr}$.*

Proof: Let P_{bl} be the set of left endpoints among the blue curves in B , and apply Lemma 1 to the red t -monotone curves R and the point set P_{bl} . Then we obtain subsets $P'_{bl} \subset P_{bl}, R_1 \subset R$, and a trapezoid Δ_{bl} , such that $|P'_{bl}|, |R_1| \geq n/(C_1 t \log^2 t)$, $P'_{bl} \subset \Delta_{bl}$, and no curve in R_1 intersects the interior of Δ_{bl} . Let $B_1 \subset B$ be the blue curves whose left endpoint belongs to P'_{bl} , and discard all curves not in $B_1 \cup R_1$. See Figure 2(a).

Let P_{br} be the right endpoint of the curves in B_1 , and apply Lemma 1 to P_{br} and R_1 . Then again, we obtain subsets $P'_{br} \subset P_{br}, R_2 \subset R_1$, and a trapezoid Δ_{br} , such that $|P'_{br}|, |R_2| \geq n/(C_1 t \log^2 t)^2$, and no curve in R_2 intersects the interior of Δ_{br} . Let $B_2 \subset B_1$ be the blues curves whose right endpoint belongs to P'_{br} , and discard all curves not in $B_2 \cup R_2$. See Figure 2(b).

We repeat this entire process to the curves in B_2 with the endpoints of R_2 , and obtain subsets $B_3 \subset B_4, R_3 \subset R_4$, trapezoids $\Delta_{bl}, \Delta_{br}, \Delta_3, \Delta_4$, such that

1. $|B_3|, |R_3| \geq n/(C_1 t \log^2 t)^4$,
2. the left endpoint of each blue curve in B_3 lies inside Δ_{bl} ,
3. the right endpoint of each blue curve in B_3 lies inside Δ_{br} ,
4. the left endpoint of each red curve in R_3 lies inside Δ_3 , and
5. the right endpoint of each red curve in R_3 lies inside Δ_4 .

Fig. 2. Finding regions Δ_{bl} , Δ_{br} , Δ_3 , Δ_4

See Figure 2(c). Now we will find regions $\Delta_{rl} \subset \Delta_3$ and $\Delta_{rr} \subset \Delta_4$ that satisfies the disjointness property (property 5). Let P_{rl} be the set of left endpoints of R_3 . Recall that regions Δ_{bl} , Δ_{br} , Δ_3 , and Δ_4 are trapezoids. Since $R \cup B$ is simple, this implies that the arrangement

$$bd(\Delta_3) \cup bd(\Delta_{bl} \cup \Delta_{br})$$

partitions the plane into at most (say) 80 cells. Since no point from P_{rl} lies inside $\Delta_{bl} \cup \Delta_{br}$, by the pigeonhole principle there exists a simply connected region Δ_{rl} in the arrangement that contains at least $n/(3C_1 t \log^2 t)^4$ points of P_{rl} , and Δ_{rl} is disjoint to $\Delta_{bl} \cup \Delta_{br}$. Let $P'_{rl} \subset P_{rl}$ be the points that lie inside Δ_{rl} , and let R_4 be the red curves whose left endpoints are in P'_{rl} .

We repeat this process to find region $\Delta_{rr} \subset \Delta_4$ and $R_5 \subset R_4$, such that $|R_5| \geq n/(9C_1 t \log^2 t)^4$, Δ_{rr} contains the right endpoints of R_5 , and Δ_{rr} is disjoint to $\Delta_{bl} \cup \Delta_{br}$. By letting $C_4 = 9C_1$, the statement of the lemma follows. $\square$

We are now ready to prove Theorem 6.

Proof of Theorem 6. We start by applying Lemma 2, to obtain subsets $R_1 \subset R$ and $B_1 \subset B$, and regions Δ_{bl} , Δ_{br} , Δ_{rl} , Δ_{rr} , with the properties described in Lemma 2. Suppose every curve in B_1 lies completely inside $\Delta_{bl} \cup \Delta_{br}$. Then all of these blue curves must be disjoint to every red curve in R_1 , and we are done. Therefore, fix a curve $\alpha \in B_1$ that does not lie completely inside $\Delta_{bl} \cup \Delta_{br}$. Then let α' be a subcurve of α that lies in $\mathbb{R}^2 \setminus (\Delta_1 \cup \Delta_2)$, and has endpoints on $bd(\Delta_1)$ and $bd(\Delta_2)$. See Figure 3(a). Now the proof falls into several cases.

Case I. Suppose that for at least $|B_1|/4$ curves $\gamma \in B_1$, regions Δ_{rl} and Δ_{rr} both lie in the same cell in the arrangement

$$\gamma \cup \alpha' \cup bd(\Delta_{bl} \cup \Delta_{br}).$$

See Figure 3(b). Then each red curve $\beta \in R_1$ intersects γ if and only if β intersects α' . Indeed, if β intersects α' , then β must intersect γ in order to come back inside the cell

(since $R_1 \cup B_1$ is simple and β does not intersect $\Delta_{bl} \cup \Delta_{br}$). Likewise, if β is disjoint to α' , then β must lie completely inside of the cell. Since at least half of the red curves in R_1 either intersect or is disjoint to α' , statement of the theorem follows.

Fig. 3. Drawing of α' , and Cases 1 and 2

Case 2. Suppose that for at least $|B_1|/4$ curves $\gamma \in B_1$, regions Δ_{rl} and Δ_{rr} lie in distinct cells in the arrangement

$$\gamma \cup \alpha' \cup bd(\Delta_{bl} \cup \Delta_{br}),$$

and one of these cells is not incident to α' (i.e. the cell is surrounded by γ and $\Delta_{bl} \cup \Delta_{br}$). See Figure 3(c). Then clearly every red edge in R_1 must intersect γ , and the statement follows.

Therefore, we can assume that we are not in case 1 or 2. Hence there exists a subset $B_2 \subset B_1$ such that $|B_2| \geq |B_1|/2$, and for each curve $\gamma \in B_2$, regions Δ_{rl} and Δ_{rr} lie in distinct cells in the arrangement

$$\gamma \cup \alpha' \cup bd(\Delta_{bl} \cup \Delta_{br}),$$

and both of these cells are incident to α' . Notice that if $\gamma \in B_2$ is disjoint to α' , then there are exactly two (adjacent) such cells, and if γ intersects α' , then there are three or four such cells incident to $\gamma \cap \alpha'$. Also notice that α' appears exactly once on the boundary of these cells, since $R \cup B$ is simple. Now we have two final cases.

Case 3. Suppose that for at least $|B_2|/2$ curves $\gamma \in B_2$, in the arrangement

$$\gamma \cup \alpha' \cup bd(\Delta_{bl} \cup \Delta_{br}),$$

Δ_{rl} and Δ_{rr} lie in distinct cells that share $\alpha'' \subset \alpha'$ as a common side. See Figures 4(a) and 4(b). Then each red curve $\beta \in R_1$ intersects γ if and only if β is disjoint to α' . Indeed, starting at Δ_{rl} , suppose β first intersects α' . Since α' appears exactly once on the boundary of both cells, β must have crossed α'' and is now in the cell that Δ_{rr} lies in. Since $R \cup B$ is simple, β must be disjoint to γ .

Now, if β crossed γ first, then β must be in the same cell as Δ_{rr} since otherwise β must return to the first cell to cross α'' , which contradicts the fact that $R \cup B$ is simple. Hence β is disjoint to α' .

Case 4. Suppose that for at least $|B_2|/2$ curves $\gamma \in B_2$, in the arrangement

$$\gamma \cup \alpha' \cup bd(\Delta_{bl} \cup \Delta_{br}),$$

Δ_{rl} and Δ_{rr} lie in distinct non-adjacent cells as in Figure 4(c). This is the final case. Then clearly, each red curve in R_1 intersects γ and the statement of the theorem follows. $\square$

Fig. 4. Cases 3 and 4

3 Proof of Theorems 3 and 4

The proof of Theorems 3 and 4 are now a standard application of Szemerédi's regularity lemma (see [17], [18]). Let us recall the weak bipartite regularity lemma.

Theorem 7 ([17]). Let $G = (X_1, X_2, E)$ be a bipartite graph with parts X_1 and X_2 such that $|X_1| = |X_2| = n$. Let $c \geq 2$. If $|E(X_1, X_2)| \geq \epsilon n^2$, then there exists subsets $Y_1 \subset X_1, Y_2 \subset X_2$ such that

1. $|Y_1| = |Y_2| = \epsilon^2 n$, and
2. $|E(Y_1, Y_2)| \geq \epsilon |Y_1| |Y_2|$, and
3. $|E(Z_1, Z_2)| > 0$ for any $Z_i \subset Y_i$, with $|Z_i| \geq |Y_i|/c$ for $i \in \{1, 2\}$.

Proof of Theorem 3. Let F be a simple family of n t -monotone curves with ϵn^2 intersecting pairs. Then by a random partition, we can split F into two subfamilies F_1, F_2 , such that $|F_1|, |F_2| \geq n/3$ and the number of pairs of curves, one from F_1 and one from F_2 , that intersect is at least $\epsilon n^2/2$.

Set $c = c_t''$, where c_t'' is defined in Theorem 6. Then by Theorem 7, there exist subsets $Y_1 \subset F_1, Y_2 \subset F_2$, of size $(\epsilon/2)^2 n/3$ each, such that for any subsets $Z_1 \subset Y_1, Z_2 \subset Y_2$ with $|Z_i| \geq |Y_i|/c$ for $i \in \{1, 2\}$, there must be a curve in Z_1 that intersects a curve in Z_2 . By Theorem 6, there exists subsets $Z_1 \subset Y_1, Z_2 \subset Y_2$ such that every curve in Z_1 intersects every curve in Z_2 , and

$$|Z_i| \geq \frac{|Y_i|}{c} \geq \frac{(\epsilon/2)^2}{3c} n \geq \epsilon^{c_t} n,$$

where c_t depends only on t . $\square$

Theorem 4 follows by replacing the word “intersect” with “disjoint” in the proof above.

4 Simple Topological Graphs with No k Pairwise Disjoint Edges

As defined in [21], the *odd-crossing number* $\text{odd-cr}(G)$ of a graph G is the minimum possible number of unordered pairs of edges that cross an odd number of times over all drawings of G . The *bisection width* of a graph G , denoted by $b(G)$, is the smallest nonnegative integer such that there is a partition of the vertex set $V = V_1 \dot{\cup} V_2$ with $\frac{1}{3} \cdot |V| \leq |V_i| \leq \frac{2}{3} \cdot |V|$ for $i = 1, 2$, and $|E(V_1, V_2)| = b(G)$. The following lemma, due to Pach and Tóth, relates the odd-crossing number of a graph to its bisection width.

Lemma 3 ([22]). *There is an absolute constant c_1 such that if G is a graph with n vertices of degrees $d_1, \dots, d_n$, then*

$$b(G) \leq c_1 \log n \sqrt{\text{odd-cr}(G) + \sum_{i=1}^n d_i^2}.$$

Since all graphs contain a bipartite subgraph with at least half of its edges, Theorem 5 immediately follows from the following theorem.

Theorem 8. *Let $G = (V, E)$ be an n -vertex simple topological bipartite graph with edges drawn as t -monotone curves. If G does not contain k pairwise disjoint edges, then $|E(G)| \leq n(\log n)^{c'_t \log k}$, where c'_t is a constant that depends only on t .*

Proof: We define $f(n, k)$ to be the maximum number of edges in an n -vertex simple topological bipartite graph with edges drawn as t -monotone curves, that does not contain k pairwise disjoint edges. We will prove by induction on n and k that

$$f(n, k) \leq n(\log n)^{c'_t \log k}.$$

Note that $f(n, k) \leq \binom{n}{2}$, and by [11] $f(n, 2) \leq 1.43n$. Now assume the statement is true for $n' < n$ and $k' < k$, and let G be an n -vertex simple topological bipartite graph with edges drawn as t -monotone curves, and does not contain k pairwise disjoint edges. The proof falls into two cases.

Case 1. Suppose there are at least $|E(G)|^2 / ((2c_1)^2 \log^6 n)$ disjoint pairs of edges in G . By Theorem 4, there exist subsets $E_1, E_2 \subset E(G)$ such that $|E_1|, |E_2| \geq |E(G)| / (c \log n)^{6c_t}$, and every edge in E_1 is disjoint to every edge in E_2 , and c is an absolute constant. Since G does not contain k pairwise disjoint edges, this implies that there exists an $i \in \{1, 2\}$ such that $|E_i|$ does not contain $k/2$ pairwise disjoint edges. Hence

$$\frac{|E(G)|}{(c \log n)^{6c_t}} \leq |E_i| \leq f(n, k/2).$$

By the induction hypothesis, we have

$$f(n, k/2) \leq n(\log n)^{c'_t \log(k/2)} \leq n(\log n)^{c'_t \log(k) - c'_t}.$$

For sufficiently large c'_t , we have $|E(G)| \leq n(\log n)^{c'_t \log(k)}$.

Case 2. Suppose there are at most $|E(G)|^2 / ((2c_1)^2 \log^6 n)$ disjoint pairs of edges in G . In what follows, we will apply a redrawing technique that was used by Pach and Tóth [22]. Since G is bipartite, let V_a and V_b be its vertex class. By applying a suitable homeomorphism to the plane, we can redraw G such that

1. the vertices in V_a are above the line $y = 1$, the vertices in V_b are below the line $y = 0$,
2. edges in the strip $0 \leq y \leq 1$ are vertical segments,
3. we have not created nor removed any crossings.

Now we reflect the part of G that lies above the $y = 1$ line about the y -axis. Then erase the edges in the strip $0 \leq y \leq 1$ and replace them by straight line segments that reconnects the corresponding pairs on the line $y = 0$ and $y = 1$. Note that our topological graph is no longer simple.

Notice that if any two edges crossed in the original drawing, then they must cross an even number of times in the new drawing. Indeed, suppose the edges e_1 and e_2 crossed in the original drawing. Since G is simple, they share exactly 1 point in common. Let k_i denote the number of times edge e_i crosses the strip for $i \in \{1, 2\}$, and note that k_i must be odd. After we have redrawn our graph, these $k_1 + k_2$ segments inside the strip will now pairwise cross, creating $\binom{k_1 + k_2}{2}$ crossing points. Since edge e_i will now cross itself $\binom{k_i}{2}$ times, this implies that there are now

$$\binom{k_1 + k_2}{2} - \binom{k_1}{2} - \binom{k_2}{2} \quad (1)$$

crossing points between edges e_1 and e_2 inside the strip. One can easily check that (1) is odd when k_1 and k_2 are odd. Since e_1 and e_2 had 1 point in common outside of the strip, this implies that e_1 and e_2 cross each other an even number of times. Note that one can easily get rid of self-intersections by making local modifications around these crossings.

Hence, the odd-crossing number in our new drawing is at most the number of disjoint pair of edges in the original drawing of G , plus the number of pair of edges that share a common vertex. Since there are at most

$$\sum_{v \in V(G)} d^2(v) \leq 2|E(G)|n$$

pairs of edges that share a vertex in G , this implies

$$\text{odd-cr}(G) \leq \frac{|E(G)|^2}{(2c_1)^2 \log^6 n} + 2|E(G)|n.$$

By Lemma 3, there is a partition of the vertex set $V = V_1 \dot{\cup} V_2$ with $\frac{1}{3} \cdot |V| \leq |V_i| \leq \frac{2}{3} \cdot |V|$ for $i = 1, 2$ and

$$b(G) \leq c_1 \log n \sqrt{\frac{|E(G)|^2}{(2c_1)^2 \log^6 n} + 4n|E(G)|}.$$

If $|E(G)|^2 / ((2c_1)^2 \log^6 n) \leq 4n|E(G)|$, then we have $|E(G)| \leq n(\log n)^{c'_t \log k}$ and we are done. Therefore we can assume

$$b(G) \leq c_1 \log n \sqrt{\frac{2|E(G)|^2}{(2c_1)^2 \log^6 n}} \leq \frac{|E(G)|}{\log^2 n}.$$

Let $|V_1| = n_1$ and $|V_2| = n_2$. By the induction hypothesis we have

$$\begin{aligned} |E(G)| &\leq b(G) + n_1(\log n_1)^{c'_t \log k} + n_2(\log n_2)^{c'_t \log k} \\ &\leq \frac{|E(G)|}{\log^2 n} + n(\log(2n/3))^{c'_t \log k} \\ &\leq \frac{|E(G)|}{\log^2 n} + n(\log n - \log(3/2))^{c'_t \log k}, \end{aligned}$$

which implies

$$|E(G)| \leq n(\log n)^{c'_t \log k} \frac{(1 - \log(3/2)/\log n)^{c'_t \log k}}{1 - 1/\log^2 n} \leq n(\log n)^{c'_t \log k}.$$

□

5 Concluding Remarks

It would be interesting to see if one can remove the t -monotone condition in Theorem 4 and in Lemma 1.

Problem 1. Given a simple family F of n curves in the plane, and an n -element point set P such that no point of P lies on a curve from F , does there exist a constant $\epsilon > 0$, a region $\Delta \subset \mathbb{R}^2$, subsets $F' \subset F, P' \subset P$ of size ϵn each, such that $P' \subset \Delta$ and every curve in F' does not intersect the interior of Δ ?

We list two more unsolved problems related to this paper.

Problem 2. Let G be an n -vertex simple topological graph with edges drawn as 2-monotone curves. If G has no two disjoint edges, then does G have at most n edges? What if the edges are drawn as 3-monotone curves?

Problem 3. Given a simple family F of n 2-monotone curves in the plane, with no 3 pairwise disjoint members, can one color the members in F with at most c colors, such that each color class consists of pairwise crossing members?

References

1. Agarwal, P.K., van Kreveld, M., Suri, S.: Label placement by maximum independent set in rectangles. *Comput. Geom. Theory Appl.* 11, 209–218 (1998)
2. Alon, N., Pach, J., Pinchasi, R., Radoicic, R., Sharir, M.: Crossing patterns of semi-algebraic sets. *J. Comb. Theory Ser. A* 111, 310–326 (2005)

3. Asano, T., Imai, H.: Finding the connected components and a maximum clique of an intersection graph of rectangles in the plane. *J. Algorithms* 4, 310–323 (1983)
4. Asplund, E., Grünbaum, B.: On a coloring problem. *Math. Scand.* 8, 181–188 (1960)
5. Basu, S.: Combinatorial complexity in o-minimal geometry. *Proc. London Math. Soc.* 100, 405–428 (2010)
6. Cairns, G., Nikolayevsky, Y.: Bounds for generalized thrackles. *Discrete Comput. Geom.* 23, 191–206 (2000)
7. Ehrlich, G., Even, S., Tarjan, R.E.: Intersection graphs of curves in the plane. *J. Combinatorial Theory, Ser. B* 21, 8–20 (1979)
8. Erdős, P., Hajnal, A.: Ramsey-type theorems. *Discrete Appl. Math.* 25, 37–52 (1989)
9. Fox, J., Pach, J., Tóth: Intersection patterns of curves. *Journal of the London Mathematical Society* 83, 389–406 (2011)
10. Fox, J., Sudakov, B.: Density theorems for bipartite graphs and related Ramsey-type results. *Combinatorica* 29, 153–196 (2009)
11. Fulek, R., Pach, J.: A computational approach to Conway’s thrackle conjecture. *Comput. Geom.* 44, 345–355 (2011)
12. Garey, M.R., Johnson, D.S.: *Computers and Intractability: A Guide to the Theory of NP-Completeness*. Freeman, New York (1979)
13. Gyárfás, A.: On the chromatic number of multiple intervals graphs and overlap graphs. *Discrete Math.* 55, 161–166 (1985)
14. Hochbaum, D.S., Maass, W.: Approximation schemes for covering and packing problems in image processing and VLSI. *J. ACM* 32, 130–136 (1985)
15. Kozik, J., Krawczyk, T., Lasoń, M., Micek, P., Pawlik, A., Trotter, W., Walczak, B.: Triangle-free intersection graphs of segments in the plane with arbitrarily large chromatic number (submitted)
16. Lovász, L., Pach, J., Szegedy, M.: On Conway’s thrackle conjecture. *Discrete Comput. Geom.* 18, 369–376 (1997)
17. Matoušek, J.: *Lectures on Discrete Geometry*. Springer, New York (2002)
18. Pach, J., Solymosi, J.: Crossing patterns of segments. *J. Comb. Theory Ser. A* 96, 316–325 (2001)
19. Pach, J., Sterling, E.: Conway’s conjecture for monotone thrackles. *Amer. Math. Monthly* 118, 544–548 (2011)
20. Pach, J., Töröcsik, J.: Some geometric applications of Dilworth’s theorem. *Discrete Comput. Geom.* 12, 1–7 (1994)
21. Pach, J., Tóth, G.: Which crossing number is it anyway? *J. Comb. Theory Ser. B* 80, 225–246 (2000)
22. Pach, J., Tóth, G.: Disjoint Edges in Topological Graphs. In: Akiyama, J., Baskoro, E.T., Kano, M. (eds.) *IJCCGGT 2003*. LNCS, vol. 3330, pp. 133–140. Springer, Heidelberg (2005)
23. Suk, A.: Coloring intersection graphs of x -monotone curves in the plane (submitted)
24. Suk, A.: Disjoint edges in complete topological graphs (to appear)
25. Tóth, G.: Note on geometric graphs. *J. Comb. Theory Ser. A* 89, 126–132 (2000)