Namophila HYACINTHACEAE

U. Eggli

Namophila U. & D. Müller-Doblies (Feddes Repert. 108(1–2): 77–79, ill., 1997). **Type:** *Namophila urotepala* U. & D. Müller-Doblies. — **Hyacinthoideae** — **Massonieae** — **Lit:** Martínez-Azorín & al. (2015: classification). **Distr:** S Namibia. **Etym:** Gr. ‘nama’, spring, stream; and Gr. ‘philos’, friend; for the occurrence in a valley with a seasonal stream.

Bulbous geophytes to 5 cm tall, bulbs usually solitary, globose, 1.5–2.3 cm Ø; **L** 2, ovate, 11–16 × 8–11.7 cm, (sub-) succulent, prostrately spreading; **Inf** 1, corymbose-capitate, between the leaves, 2- to 3-flowered (to 19-flowered in cultivation), raceme 1–4 cm; **Bra** linear, inconspicuous, lowest 17–36 × (1.3–) 2.5–3.5 mm; **Ped** 7–18 mm; **Fl** stellate, green, to 33 mm Ø; **Tep** 19–23 mm, basally united and forming a 4–6 mm long tube, lobes (12–) 15–17 × 3.5–4.5 mm, narrowly triangular, with a distinctly caudate tip 1–4.7 mm long; **St** basally fused to the tepal tube; free part of **Fil** 6–8 mm, white; **Anth** 2–2.5 mm, green with yellow pollen; **Ov** obclavate, green, 3.5–5 × 3–3.7 mm; **Sty** 8–10 mm, green; **Sti** punctiform; **Fr** enveloped by the fleshy perianth, 10 mm; **Se** globose, 2 × 1.2–1.8 mm, black, glossy.

The monotypic genus *Namophila* is closely related to *Massonia* according to the molecular

study by Manning & al. (2004). See *Massonia* for further comments on the classification of the group.

N. urotepala U. & D. Müller-Doblies (Feddes Repert. 108(1–2): 77–79, ill., 1997). **Type:** Namibia, Witputz (*Müller-Doblies* 88064g [WIND, B, BOL, BR, BTU, E, G, K, LI, M, MO, NBG, PRE, S, Z]). — **Lit:** Martínez-Azorín & al. (2015: with ill.). **Distr:** S Namibia (Witputz: S Namibian Escarpment); winter-rainfall area, ±700 m, spring-flowering.

Description as for the genus.

References

- Manning, J. C. [& al. 2004], Goldblatt, P. & Fay, M. F. (2004) A revised generic synopsis of *Hyacinthaceae* in sub-Saharan Africa, based on molecular evidence, including new combinations and the new tribe *Pseudoprosperaeae*. *Edinburgh J. Bot.* 60(3): 533–568, keys. <https://doi.org/10.1017/S0960428603000404>
- Martínez-Azorín, M. [& al. 2015], Pinter, M. & Wetschnig, W. (2015) *Desertia*, a new genus in *Massonieae* (*Asparagaceae*, *Scilloideae*), including the description of *Desertia luteovirens* and the taxonomic revisions of *Whiteheadia* and *Namophila*. *Phytotaxa* 221(3): 201–225, ill., maps, key. <https://doi.org/10.11646/phytotaxa.221.3.1>

U. Eggli (✉)
Sukkulanten-Sammlung Zürich, Grün Stadt Zürich,
Zürich, Switzerland
e-mail: Urs.Eggli@zuerich.ch