Experimental Hydrogen Production in Hydrothermal and Fault Systems: Significance for Habitability of Subseafloor H₂ Chemoautotroph Microbial Ecosystems

Katsuhiko Suzuki, Takazo Shibuya, Motoko Yoshizaki, and Takehiro Hirose

Abstract

Hydrogen generated in hydrothermal and fault systems has recently received considerable attention as a potential energy source for hydrogen-based microbial activity such as methanogenesis. Laboratory experiments that have reproduced conditions for the serpentinization of ultramafic rocks such as peridotite and komatiite have clarified the chemical and petrological processes of H₂ production. In a frictional experimental study, we recently showed that abundant H₂ can also be generated in a simulated fault system. This result suggests that microbial ecosystems might exist in subseafloor fault systems. Here we review the experimental constraints on hydrogen production in hydrothermal and fault systems.

Keywords

Friction experiment • Hydrogen generation • Hydrothermal experiment • Simulated fault system • Ultramafic rocks

8.1 Introduction

The discovery of chemolithoautotrophic microbial ecosystems in the modern ocean has suggested that hydrothermal fluids enriched in H₂ fuel H₂-driven primary producers such as hyperthermophilic hydrogenotrophic

methanogens (Cannat et al. 1997; Takai et al. 2004; Kelley et al. 2005). Moreover, multidisciplinary studies have indicated that an H₂-driven chemolithoautotrophic ecosystem may have supported the earliest life on Earth (Russell and Hall 1997; Sleep et al. 2004; Kelley et al. 2005; Canfield et al. 2006; Takai et al. 2006). H₂-rich hydrothermal fluids are generated by the serpentinization of Mg- and Fe-rich ultramafic rocks, which are igneous and meta-igneous

K. Suzuki (✉)

Institute for Research on Earth Evolution (IFREE), Japan Agency for Marine-Earth Science and Technology (JAMSTEC), 2-15 Natsushima, Yokosuka 237-0061, Japan

Precambrian Ecosystem Laboratory, Japan Agency for Marine-Earth Science and Technology (JAMSTEC), 2-15 Natsushima, Yokosuka 237-0061, Japan

Seafloor Resources Research Project (SRRP), Japan Agency for Marine-Earth Science and Technology (JAMSTEC), 2-15 Natsushima, Yokosuka 237-0061, Japan
e-mail: katz@jamstec.go.jp

T. Shibuya

Precambrian Ecosystem Laboratory, Japan Agency for Marine-Earth Science and Technology (JAMSTEC), 2-15 Natsushima, Yokosuka 237-0061, Japan

Seafloor Resources Research Project (SRRP), Japan Agency for Marine-Earth Science and Technology (JAMSTEC), 2-15 Natsushima, Yokosuka 237-0061, Japan

M. Yoshizaki

Precambrian Ecosystem Laboratory, Japan Agency for Marine-Earth Science and Technology (JAMSTEC), 2-15 Natsushima, Yokosuka 237-0061, Japan

Department of Earth and Planetary Sciences, Tokyo Institute of Technology, 2-12-1 Ookayama, Meguro-ku, Tokyo 152-8551, Japan

T. Hirose

Kochi Institute for Core Sample Research, Japan Agency for Marine-Earth Science and Technology (JAMSTEC), 200 Monobe-otsu, Nankoku, Kochi 783-8502, Japan

rocks consisting mainly (>90 %) of mafic minerals with high magnesium and iron contents. These rocks have a very low silica content (<45 %), MgO generally more than 18 %, high FeO, and low potassium. The Earth's mantle is composed of ultramafic rocks. The production of hydrogen by the serpentinization of ultramafic rocks is a central tenet of the hypothesis that life first emerged on Earth in an H₂-rich hydrothermal environment (Russell et al. 2010).

In the modern ocean, the basement rocks of the oceanic crust are generally composed of mid-ocean ridge basalt (MORB). Along slow-spreading ridges such as the North Atlantic Ridge and the Southwest Indian Ridge, volcanic activity is low and severely serpentinized peridotites (peridotite: ultramafic rocks composed of olivine and pyroxene) are exposed along transform faults (e.g., Miyashiro et al. 1969; Aumento and Loubat 1971; Cannat 1993). Hydrogen-enriched hydrothermal fluids are common in such peridotite-dominated slow-spreading ridge settings (Kelley et al. 2001; Früh-Green et al. 2004). Since the first discovery of natural peridotite-hosted hydrothermal vents with abundant H₂ at Logachev field on the Mid-Atlantic Ridge in 1995 (Krasnov et al. 1995), other peridotite-hosted and -associated hydrothermal vents have been found, namely, the Rainbow, Nibelungen, Lost City, and Achaze fields on the Mid-Atlantic Ridge (Charlou et al. 1998, 2002, 2008; Douville et al. 2002; Kelley et al. 2005; Melchert et al. 2008) and the Kairei field on the Central Indian Ridge (Van Dover et al. 2001). A pronounced feature of peridotite-hosted systems is the presence of hydrothermal solutions enriched not only in H₂ but also in CH₄ and other hydrocarbons (Charlou et al. 2002). Measured H₂ concentrations in these fluids range from 2.5 to 16 mmol/kg, with the highest concentrations reported at steady-state, unsedimented mid-ocean ridge hydrothermal vents. The generated H₂ supports ecosystems, including hyperthermophilic subsurface lithoautotrophic ecosystems (HyperSLiME) (Takai et al. 2004; Nealson et al. 2005), in which methanogens utilize H₂ and CO₂ and produce methane as a metabolic product.

Several petrographical (Cressey 1979) and theoretical (Wetzel and Shock 2000; Sleep et al. 2004) studies have examined the H₂-generating geochemical and mineralogical reactions associated with the serpentinization of ocean-floor ultramafic rocks, and experimental investigations (Berndt et al. 1996; Allen and Seyfried 2003; Seyfried et al. 2007) have significantly increased our understanding of the mechanisms of H₂ production during serpentinization. In the present paper, we focus on these experimental results, because it is generally difficult to directly observe chemical processes in seafloor environments. Thermodynamic modeling of seawater–peridotite reactions is also an important tool, both for understanding the experimental results and for inferring the natural serpentinization processes (Klein et al. 2009; McCollom and Bach 2009). McCollom and

Fig. 8.1 Predicted alteration mineralogy and fluid composition during hydrothermal alteration of harzburgite over a range of temperatures at a 1:1 water:rock ratio, modified from McCollom and Bach (2009). (a) Equilibrium mineral composition: amount of each mineral per kilogram of harzburgite reacted. (b) H₂ concentration in the fluid

Bach (2009) examined the effect of temperature on the mineral assemblage and fluid composition produced by serpentinization of harzburgite (a peridotite consisting mainly of olivine and orthopyroxene) (Fig. 8.1). They reported that at temperatures below 315 °C, the serpentinized rock was composed of typical serpentinite minerals, such as serpentine, brucite, magnetite, and minor secondary clinopyroxene. With increasing temperature, the amount of magnetite increased, and consequently, the concentration of H₂ generated by serpentinization also increased with temperature up to ~360 mmol/kg. Above 315 °C, however, olivine became stable and coexisted in equilibrium with other secondary minerals and fluid, and above 390 °C it remained almost completely unaltered. Therefore, the amount of Fe converted to magnetite decreased as the temperature increased above 315 °C, with the result that the H₂ concentration in the fluid was lower at these higher temperatures. Contrary to expectation, theoretical modeling of these water–rock interactions does not always produce results that are quantitatively consistent with the experimental results. Therefore, further investigation is needed to

clarify the reasons for the discrepancy between theoretical and experimental results.

Another extreme environment where H_2 may be abundant is found in natural fault systems. Wakita et al. (1980) first reported high concentrations of H_2 (up to 3 % v/v) in soil gas from sites in the Yamasaki fault zone, southwestern Japan, and Wiersberg and Erzinger (2008) reported high concentrations of H_2 in drilling cores obtained near micro-earthquake hypocenters along the San Andreas fault in California. These observations led to the hypothesis that methanogenic ecosystems might also be found in deep fault systems below the seafloor. To examine whether fault-driven H_2 generation can produce enough H_2 to maintain a chemolithoautotrophic microbial ecosystem, we conducted high-velocity sliding experiments using velocities and displacements typical of natural earthquakes (Hirose et al. 2011, 2012).

Thus, both hydrothermal and frictional experimental systems can be effectively used to elucidate physicochemical processes in natural systems. In this chapter, we review experiments, performed both as part of Project TAIGA (Trans-crustal Advection and In-situ bio-geochemical processes of Global sub-seafloor Aquifer) and by other groups, examining the generation of hydrogen by water/rock interactions at high temperature and pressure, including both reactions between water and komatiite, an Archean volcanic ultramafic rock, and those in frictional fault systems.

8.2 Constraints on H_2 Production During Experimental Hydrothermal Alteration of Ultramafic Rocks

Peridotite exposed on slow-spreading ridge systems is often severely serpentinized as a result of hydration due to long-term reaction with seawater (Ildefonse et al. 2007; Morishita et al. 2009; Nakamura et al. 2009). Isotopic investigations have shown that such serpentinized peridotites, similar to altered MORB, are the products of high-temperature and -pressure reactions between peridotite and seawater (Wenner and Taylor 1971, 1973; Sakai and Tsutsumi 1978). Hydrothermal alteration of peridotite was studied experimentally (Seyfried and Dibble 1980; Hajash and Chandler 1981; Janecky and Seyfried 1986) even before the discovery of natural peridotite-hosted hydrothermal vents at Logachev field (Krasnov et al. 1995). Reactions between peridotite and seawater under high temperature and pressure have also been examined in batch-type experiments (see Chap. 7). Under high temperature and pressure, the oxidation of ferrous ion [Fe(II)] in primary minerals such as olivine and pyroxene to Fe(III)

in secondary minerals such as magnetite reduces water and releases H_2 gas:

where $(\text{FeO})_{\text{rock}}$ denotes the ferrous constituent of a primary silicate mineral such as olivine and $(\text{Fe}_2\text{O}_3)_{\text{rock}}$ denotes the ferric constituent of an secondary alteration mineral such as magnetite. The serpentinization of olivine (Fo_{90}) generates H_2 as follows:

In this reaction, olivine supplies Fe(II), and hydrogen and magnetite, which contains Fe(II) and Fe(III) in equal amounts, are produced. Note that Eq. (8.2) is a simplified formula provided to illustrate the generation of H_2 by serpentinization; in natural systems, the produced serpentine and brucite commonly contain Fe as well.

Janecky and Seyfried (1986) experimentally investigated the serpentinization of harzburgite at 300 °C and 50 MPa, but they did not measure hydrogen continuously during their experiment. Instead, they collected samples for hydrogen measurement only twice, after 10 months and 2 years. In both samples, they found high concentrations of H_2 (0.1 and 0.33 mmol/kg), but because they did not measure H_2 in other fluid samples collected during the experiment, the H_2 generation reaction cannot be unambiguously interpreted. Using a similar experimental apparatus, McCollom and Seewald (2001) assessed the potential of olivine serpentinization to reduce CO_2 and produce hydrocarbons. In their experiments, they reacted powdered olivine with a 0.5 M NaCl solution at 300 °C and 35 MPa and obtained abundant H_2 (concentration > 70 mmol/kg, Fig. 8.2) after 700 h.

Allen and Seyfried (2003) reacted olivine, orthopyroxene, and clinopyroxene, the major constituent minerals in peridotite, with a NaCl–MgCl₂ solution at 400 °C and 50 MPa to assess the potential of these minerals to produce H_2 . They used olivine (Fo_{89}), orthopyroxene (En_{85}), and clinopyroxene (Di_{89}) both individually and in combination to constrain the chemical processes of ultramafic-hosted hydrothermal systems in mid-ocean ridges. The olivine alteration rate was slow, as indicated by the absence of hydrous alteration products, whereas reactions including pyroxenes were rapid, resulting in significant increases in dissolved H_2 as well as in Ca, SiO₂, and Fe (Fig. 8.3), and in the formation of SiO₂-rich secondary minerals such as talc, tremolite, and magnetite (Fig. 8.3; McCollom and Bach 2009). High pH and low Fe

concentrations are theoretically predicted under the assumption of full equilibrium at 400 °C and 50 MPa in the MgO-CaO-FeO-Fe₂O₃-SiO₂-Na₂O-H₂O-HCl system (Allen and Seyfried 2003). In their laboratory experiments, however, Allen and Seyfried (2003) reported that when the initial olivine:pyroxene ratio was 3:1 (the typical mineral

composition of abyssal peridotite), pH remained low and Fe concentrations remained high. Moreover, the pH also remained relatively low in their experiments that included orthopyroxene and clinopyroxene, a result that may have been caused by talc–fluid and talc–tremolite–fluid equilibria, respectively.

Allen and Seyfried (2003) suggested that the reactions occurring in ultramafic rock-hosted hydrothermal systems such as the Rainbow system on the Mid-Atlantic Ridge may be very similar to those observed in these experiments, because of the similarity of the experimental conditions to the temperature and chemistry conditions of the vent fluid, in which concentrations of SiO₂, Ca, H₂, and Fe are relatively high (Charlou et al. 1998, 2002). In fact, Fe concentrations in the Rainbow system, which are the highest of any vent system yet discovered, imply a relatively low pH in the subseafloor reaction zone. The findings of Allen and Seyfried (2003) are very important because the experimental results together with the observed Rainbow fluid chemistry suggest that pyroxene dissolution is the dominant reaction in the Rainbow hydrothermal system, even though abundant olivine is present in the reaction zone. To better constrain the temporal evolution of the hydrothermal alteration of ultramafic rocks in subseafloor reaction zones, experimental data

Fig. 8.2 Measured concentrations of dissolved H₂ during a hydrothermal reaction in the presence of olivine at 300 °C and 35 MPa as a function of reaction time. Modified from McCollom and Seewald (2001)

Fig. 8.3 Changes in the dissolved concentrations of selected species with time during four experiments, modified from Allen and Seyfried (2003). All experiments were performed at 400 °C, 50 MPa and with mineral:fluid ratios from 2 to 4. Values on the left vertical axis apply to

species represented by *open symbols* and to Mg, and those on the right axis apply to species represented by *closed symbols*. Cpx clinopyroxene, Opx orthopyroxene

Fig. 8.4 Calculated (line) and measured (inverted triangles) time-series changes in dissolved hydrogen in fluid coexisting with lherzolite and alteration products (Seyfried et al. 2007). The difference between measured and predicted dissolved H_2 concentrations may reflect (in part) the lack of thermodynamic data for Fe-rich alteration minerals in the database used for the theoretical model

obtained under various conditions should be compared with thermodynamic simulation results.

The experiments just described were conducted at high temperatures (300 °C or higher), but actual reaction zone temperatures in ultramafic rocks are likely to be variable. For example, in the Lost City hydrothermal field on the Mid-Atlantic Ridge, H_2 is abundant, the pH is high, and the reactions occur at moderately low temperature. When Seyfried et al. (2007) experimentally constrained hydrogen production at the low temperature of 200 °C and 50 MPa, they found that changes in the concentrations of Ca, Mg, and Si in the reacted fluid agreed quantitatively with those predicted theoretically by a reaction pathway model of seawater–lherzolite interaction (fluid:rock mass ratio, 1:1) at 200 °C and 50 MPa. However, the time variations of pH, dissolved chloride, and H_2 monitored during their experiment did not agree with the theoretically predicted results (Fig. 8.4). Dissolved H_2 increased in a series of abrupt steps, and reached a maximum concentration that was only about 20 % of the theoretically predicted concentration. Interestingly, serpentine, the most abundant alteration mineral, contained both ferric and ferrous iron, and magnetite was present only in trace amounts. They therefore inferred that the low rate of H_2 generation, which was lower than that estimated on the basis of predicted serpentinization rates, was due to the production of diverse Fe-bearing alteration minerals.

In a recent experiment, Mayhew et al. (2013) reacted ultramafic and mafic rocks such as peridotite and minerals such as pyroxene, olivine, and magnetite with an anoxic fluid at 55 °C and 100 °C, temperatures that are habitable

for (hyper)thermophilic microbes. Their synchrotron-based X-ray analysis results showed that the amount of H_2 produced was strongly dependent on whether spinel phases such as magnetite were present, and they observed Fe(III)-oxide products on the surface of the spinels. They proposed, therefore, that Fe(II) and water adsorbed on the surface of spinels reacted together under kinetic control to produce H_2 at low temperatures.

8.3 Experimental H_2 Generation During Komatiite Alteration: Simulation of an Archean Hydrothermal System

The experiments described in Sect. 8.2 showed by simulating peridotite–seawater systems that abundant H_2 is generated in a modern slow-spreading ridge by the serpentinization of peridotite. Sufficiently high concentrations of H_2 occur in the vent fluids of these systems to sustain an ecosystem based on H_2 -driven chemolithoautotrophic primary producers such as methanogens. Moreover, an H_2 -driven ecosystem is the most probable candidate habitat for the earliest life on Earth (Russell and Hall 1997; Sleep et al. 2004; Kelley et al. 2005; Takai et al. 2006; Canfield et al. 2006). On this basis, it has been hypothesized that hydrothermal fluids with H_2 , sufficiently abundant to sustain methanogens, existed in the early Earth (Takai et al. 2006).

In contrast to the modern ocean, in the Hadean and early Archean ocean, peridotite was probably scarce beneath the crust of the ocean floor, because at that time the oceanic crust may have been two to three times the thickness of the modern oceanic crust (Ohta et al. 1996; Moores 2002; Takai et al. 2006). In this tectonic setting, komatiite, a distinctive ultramafic volcanic rock, was most likely the main component of ultramafic rocks on the floor of Hadean and Archean oceans. In fact, komatiite is common in Archean greenstone belts, whereas it is less common in Proterozoic successions and quite rare in Phanerozoic strata (Condie 2005). We therefore hypothesized that komatiite was the dominant ultramafic rock in the early history of the Earth (Yoshizaki et al. 2009) and that the interaction of komatiite with fluid at high temperature produced abundant H_2 . To test this hypothesis, we examined whether we could reproduce a hydrothermal environment habitable for early life by experimentally reacting komatiite with water.

Modern komatiite in exposed Archean rocks has already been severely altered, so komatiite in its present form could not be used for our experiments. Therefore, we synthesized komatiite glass by dehydrating and remelting serpentinized komatiite collected from the Komati Formation, Barberton Greenstone Belt, South Africa, and then reacted the synthesized komatiitic glass with an NaCl solution at 300 °C and 50 MPa (Yoshizaki et al. 2009).

Fig. 8.5 Changes in the dissolved H₂ concentration with time during a komatiitic glass–water reaction experiment (Yoshizaki et al. 2009). The experiment was performed at 300 °C and 50 MPa with an initial water: rock ratio of 4:1

The concentration of produced H₂ reached 2.4 mmol/kg after 1500 h (Fig. 8.5) (Yoshizaki et al. 2009). This concentration is comparable to concentrations obtained by hydration of peridotitic rocks (see Sect. 8.2). These results suggest that hydrothermal alteration of komatiite in the Hadean and early Archean may have provided sufficient H₂ to fuel microbial ecosystems in the vicinity of hydrothermal vents.

8.4 Mechanoradical H₂ Generation During Simulated Faulting

As we have described, hydrothermal alteration of peridotites and komatiites can produce abundant H₂ in subseafloor settings. Other potential sites of H₂ generation are active fault systems. Earthquake or fault-related H₂ generation was first found by gas monitoring along the active Yamasaki fault, southwestern Japan (Wakita et al. 1980), and more recently H₂ generation has been inferred from drill cores obtained near microearthquake hypocenters along the San Andreas fault in California (Wiersberg and Erzinger 2008). Kita et al. (1982) suggested that H₂ may be generated in fault systems by the following reaction, expressed in terms of mechanoradicals on fresh surfaces of silicate minerals and water molecules:

Experiments in which a ball mill was used to crush rocks had previously reproduced possible mechanoradical reactions during faulting (Kita et al. 1982; Kameda et al. 2004), but no quantitative investigations of the generation of H₂ in an active fault system have been carried out. To

Fig. 8.6 (a) Hydrogen generation (μmol) as a function of frictional work (kJ) during high-velocity friction experiments with dry and wet basalt specimens at a slip velocity of 1.6 m/s and a displacement of 10 m under an air or argon gas atmosphere (Hirose et al. 2011). The amount of hydrogen generated tended to increase linearly with frictional work (correlation coefficient of 0.814, 0.899, and 0.777 for wet-air, dry-Ar gas, and dry-air conditions, respectively). (b) A typical shear stress versus displacement curve obtained during a friction experiment. Frictional work was calculated by integrating shear stress over the displacement (area under the shear stress versus displacement curve) and then multiplying the result by the fault surface area

examine whether a natural fault system can produce enough H₂ to sustain a microbial methanogen-based ecosystem, we performed high-velocity sliding experiments that reproduced slip velocities and displacements typical of natural earthquakes and then quantitatively estimated the concentrations of H₂ produced (Hirose et al. 2011). To collect the generated gas samples, we placed a reaction cell around the rock specimen in a high-velocity frictional experiment system (see Chap. 7 in this volume). The results showed that H₂ generation increases with frictional work (i.e., earthquake magnitude) (Fig. 8.6). Therefore, the earthquake-derived H₂ flux in nature can be estimated by establishing the correlation between H₂ production and earthquake magnitude (Hirose et al. 2011, 2012). Moreover, an H₂ fluid concentration higher than 1.1 mmol/kg can be achieved in a fault zone after even a small-magnitude earthquake (Hirose et al. 2011, 2012). This concentration of H₂ is potentially high enough to sustain a methanogen-based ecosystem, we can expect to find diverse microbial ecosystems in deep fault systems. Thus, such fault systems may also have provided habitable zones for early life on Earth.

8.5 Concluding Remarks and Future Perspectives

In modern oceans, H₂-rich hydrothermal fluids (i.e., H₂ concentrations from a few to a few tens mmol/kg) have been observed in ultramafic rock-hosted systems. Such concentrations are sufficiently high to sustain H₂-based lithoautotrophic microbial ecosystems, including phylogenetically ancient microbes (Takai et al. 2006). In this regard, the experiments simulating H₂ generation described here have improved our understanding of not only geochemical and biological interactions driven by serpentinization in modern oceans but also the potential habitability of Hadean/Archean komatiite hydrothermal systems, in which life might have originated. Additionally, the results of friction experiments suggest the intriguing idea that H₂ generated in a fault system might also sustain lithoautotrophic microbial ecosystems. Such fault system ecosystems may also have existed on the early Earth. To verify the fault zone model, we anticipate the results of microbial investigations of fault systems performed using fresh drill cores. It should be emphasized, however, that there are certain basic difficulties associated with the described hydrothermal experiments. For example, as described in Sect. 8.2, some results from experimental hydrothermal reactions conflict with both natural observations and theoretical calculations. The discrepancies may reflect mainly inappropriate initial experimental conditions or inaccurate thermodynamic data. Thus, future experiments must correct these problems.

Acknowledgments We thank Mamiko Umetsu for drawing some of the figures. Comments by Prof. Fumitaka Yanagisawa considerably improved our manuscript. This study was supported by a Grant-in-Aid for Scientific Research on Innovative Areas (no. 20109006) to “Project TAIGA: Trans-crustal Advection and In-situ bio-geochemical processes of Global sub-seafloor Aquifer” from the Ministry of Education, Culture, Sports, Science and Technology (MEXT) of Japan. We also thank Prof. Urabe and members of Project TAIGA for their collaboration and useful discussions.

Open Access This chapter is distributed under the terms of the Creative Commons Attribution Noncommercial License, which permits any noncommercial use, distribution, and reproduction in any medium, provided the original author(s) and source are credited.

References

- Allen DE, Seyfried WE (2003) Compositional controls on vent fluids from ultramafic-hosted hydrothermal systems at mid-ocean ridges: an experimental study at 400 °C, 500 bars. *Geochim Cosmochim Acta* 67:1531–1542. doi:[10.1016/s0016-7037\(02\)01173-0](https://doi.org/10.1016/s0016-7037(02)01173-0)
- Aumento F, Loubat H (1971) The Mid-Atlantic Ridge near 45°N, serpentinized ultramafic intrusions. *Can J Earth Sci* 8:631–663
- Berndt ME, Allen DE, Seyfried WE (1996) Reduction of CO₂ during serpentinization of olivine at 300 °C and 500 bar. *Geology* 24:351–354
- Canfield DE, Rosing MT, Bjerrum C (2006) Early anaerobic metabolisms. *Philos Trans R Soc B Biol Sci* 361:1819–1834
- Cannat M (1993) Emplacement of mantle rocks in the seafloor at mid-ocean ridges. *J Geophys Res* 98:4163–4172
- Cannat M, Lagabrie Y, Bougault H, Casey J, de Coutures N, Dmitriev L, Fouquet Y (1997) Ultramafic and gabbroic exposures at the Mid-Atlantic Ridge: geological mapping in the 15°N region. *Tectonophysics* 279:193–213
- Charlou JL, Fouquet Y, Bougault H, Donval JP, Etoubleau J, Jean-Baptiste P, Dapigny A, Appriou P, Rona PA (1998) Intense CH₄ plumes generated by serpentinization of ultramafic rocks at the intersection of the 15°20′N fracture zone and the Mid-Atlantic Ridge. *Geochim Cosmochim Acta* 62:2323–2333. doi:[10.1016/S0016-7037\(98\)00138-0](https://doi.org/10.1016/S0016-7037(98)00138-0)
- Charlou JL, Donval JP, Fouquet Y, Jean-Baptiste P, Holm N (2002) Geochemistry of high H₂ and CH₄ vent fluids issuing from ultramafic rocks at the Rainbow hydrothermal field (36°14′N, MAR). *Chem Geol* 191:345–359
- Charlou JL, Donval JP, Konn C, Birot D, Sudarikov S, Jean-Baptiste P, Fouquet Y; Scientific Party of the Serpentine Cruise (2008) High hydrogen and abiotic hydrocarbons from new ultramafic hydrothermal sites between 12°N and 15°N on the Mid Atlantic Ridge—Results of the SERPENTINE cruise, EOS Transactions, AGU, Fall Meeting Supplement, Abstract, 88, T51F-04
- Condie KC (2005) *Earth as an evolving planetary system*. Elsevier, Burlington
- Cressey BA (1979) Electron microscopy of serpentinite textures. *Can Mineral* 17:741–756
- Douville E, Charlou JL, Oelkers EH, Bienvu P, Jove Colon CF, Fouquet DJPY, Prieur D, Appriou P (2002) The rainbow vent fluids (36°14′N, MAR): the influence of ultramafic rocks and phase separation on trace metal content in Mid-Atlantic Ridge hydrothermal fluids. *Chem Geol* 184:37–48
- Früh-Green GL, James AD, Plas A, Kelley DS, Geobety B (2004) Serpentinization of oceanic peridotites: implications for geochemical cycles and biological activity. *Subseafloor Biosphere Mid Ocean Ridges* 144:119–136
- Hajash A, Chandler GW (1981) An experimental investigation of high-temperature interactions between seawater and rhyolite, andesite, basalt and peridotite. *Contrib Mineral Petrol* 78:240–254
- Hirose T, Kawagucci S, Suzuki K (2011) Mechanoradical H₂ generation during simulated faulting: Implications for an earthquake-driven subsurface biosphere. *Geophys Res Lett* 38. doi:[10.1029/2011gl048850](https://doi.org/10.1029/2011gl048850)
- Hirose T, Kawagucci S, Suzuki K (2012) Correction to “Mechanoradical H₂ generation during simulated faulting: implications for an earthquake-driven subsurface biosphere”. *Geophys Res Lett* 39:L23304 doi:[10.1029/2012GL054539](https://doi.org/10.1029/2012GL054539)
- Ildefonse B, Blackman DK, John BE, Ohara Y, Miller DJ, MacLeod CJ (2007) Oceanic core complexes and crustal accretion at slow-spreading ridges. *Geology* 35:623–626. doi:[10.1130/G23531A.1](https://doi.org/10.1130/G23531A.1)
- Janecky DR, Seyfried WE Jr (1986) Hydrothermal serpentinization of peridotite within the oceanic crust: experimental investigations of mineralogy and major element chemistry. *Geochim Cosmochim Acta* 50:1357–1378
- Janecky DR, Seyfried WE (1986) Hydrothermal serpentinization of peridotite within the oceanic-crust: experimental investigations of mineralogy and major element chemistry. *Geochim Cosmochim Acta* 50:1357–1378. doi:[10.1016/0016-7037\(86\)90311-x](https://doi.org/10.1016/0016-7037(86)90311-x)
- Kameda J, Saruwatari K, Tanaka H, Tsunomori F (2004) Mechanisms of hydrogen generation during the mechanochemical treatment of biotite within D₂O media. *Earth Planets Space* 56:1241–1245
- Kelley DS, Karson JA, Blackman DK, Früh-Green GL, Butterfield DA, Lilley MD, Olson EJ, Schrenk MO, Roe KK, Lebon GT, Rivizzigno

- P, Party ATS (2001) An off-axis hydrothermal vent field near the Mid-Atlantic Ridge at 30 degrees N. *Nature* 412:145–149. doi:[10.1038/35084000](https://doi.org/10.1038/35084000)
- Kelley DS, Karson JA, Fruh-Green GL, Yoerger DR, Shank TM, Butterfield DA, Hayes JM, Schrenk MO, Olson EJ, Proskurowski G, Jakuba M, Bradley A, Larson B, Ludwig K, Glickson D, Buckman K, Bradley AS, Brazelton WJ, Roe K, Elend MJ, Delacour A, Bernasconi SM, Lilley MD, Baross JA, Summons RT, Sylva SP (2005) A serpentinite-hosted ecosystem: the lost city hydrothermal field. *Science* 307:1428–1434. doi:[10.1126/science.1102556](https://doi.org/10.1126/science.1102556)
- Kita I, Matsuo S, Wakita H (1982) H₂ Generation by reaction between H₂O and crushed rock: an experimental study on H₂ degassing from the active fault zone. *J Geophys Res* 87:789–795. doi:[10.1029/JB087iB13p10789](https://doi.org/10.1029/JB087iB13p10789)
- Klein F, Bach W, Joens N, McCollom T, Moskowitz B, Berquo T (2009) Iron partitioning and hydrogen generation during serpentinization of abyssal peridotites from 15 degrees N on the Mid-Atlantic Ridge. *Geochim Cosmochim Acta* 73:6868–6893. doi:[10.1016/j.gca.2009.08.021](https://doi.org/10.1016/j.gca.2009.08.021)
- Krasnov SG, Cherkashev GA, Stepanova TV, Batuyev BN, Krotov AG, Malin BV, Maslob MN, Markov VF, Poroshina IM, Samovarov MS, Ashadze AM, Lazareva LI, Ermolayev IK (eds) (1995) Detailed geological studies of hydrothermal fields in the North Atlantic. Geological Society of London, Special Publication, pp 43–64
- Mayhew LE, Ellison ET, McCollom TM, Trainor TP, Templeton AS (2013) Hydrogen generation from low-temperature water-rock reactions. *Nat Geosci* 6:478–484. doi:[10.1038/NGEO1825](https://doi.org/10.1038/NGEO1825)
- McCollom TM, Bach W (2009) Thermodynamic constraints on hydrogen generation during serpentinization of ultramafic rocks. *Geochim Cosmochim Acta* 73:856–875. doi:[10.1016/j.gca.2008.10.032](https://doi.org/10.1016/j.gca.2008.10.032)
- McCollom TM, Seewald JS (2001) A reassessment of the potential for reduction of dissolved CO₂ to hydrocarbons during serpentinization of olivine. *Geochim Cosmochim Acta* 65:3769–3778. doi:[10.1016/s0016-7037\(01\)00655-x](https://doi.org/10.1016/s0016-7037(01)00655-x)
- Melchert B, Devey CW, German CR, Lackschewitz KS, Seifert R, Walter M, Mertens C, Yoerger DR, Baker ET, Paulick H, Nakamura K (2008) First evidence for high-temperature off-axis venting of deep crustal/mantle heat: the Nibelungen hydrothermal field, southern Mid-Atlantic Ridge. *Earth Planet Sci Lett* 275:61–69
- Miyashiro A, Shido F, Ewing M (1969) Composition and origin of serpentinites from the Mid-Atlantic Ridge near 24° and 30° North Latitude. *Contrib Mineral Petrol* 23:117–127
- Moores EM (2002) Pre-1 Ga (pre-Rodinian) ophiolites: their tectonic and environmental implications. *Geol Soc Am Bull* 114:80–95. doi:[10.1130/0016-7606\(2002\)114<0080:pgprot>2.0.co;2](https://doi.org/10.1130/0016-7606(2002)114<0080:pgprot>2.0.co;2)
- Morishita T, Hara K, Nakamura K, Sawaguchi T, Tamura A, Arai S, Okino K, Takai K, Kumagai H (2009) Igneous, alteration and exhumation processes recorded in abyssal peridotites and related fault rocks from an oceanic core complex along the Central Indian Ridge. *J Petrol* 50. doi:[10.1093/petrology/egp025](https://doi.org/10.1093/petrology/egp025)
- Nakamura K, Morishita T, Bach W, Klein F, Hara K, Okino K, Takai K, Kumagai H (2009) Serpentinized troctolites exposed near the Kairei Hydrothermal Field, Central Indian Ridge: Insights into the origin of the Kairei hydrothermal fluid supporting a unique microbial ecosystem. *Earth Planet Sci Lett* 280:128–136. doi:[10.1016/j.epsl.2009.01.024](https://doi.org/10.1016/j.epsl.2009.01.024)
- Nealson KH, Inagaki F, Takai K (2005) Hydrogen-driven subsurface lithoautotrophic microbial ecosystems (SLiMEs): do they exist and why should we care? *Trends Microbiol* 13:405–410. doi:[10.1016/j.tim.2005.07.010](https://doi.org/10.1016/j.tim.2005.07.010)
- Ohta H, Maruyama S, Takahashi E, Watanabe Y, Kato Y (1996) Field occurrence, geochemistry and petrogenesis of the Archean mid-oceanic ridge basalts (AMORBs) of the Cleaverville area, Pilbara craton, western Australia. *Lithos* 37:199–221. doi:[10.1016/0024-4937\(95\)00037-2](https://doi.org/10.1016/0024-4937(95)00037-2)
- Russell MJ, Hall AJ (1997) The emergence of life from iron monosulphide bubbles at a submarine hydrothermal redox and pH front. *J Geol Soc* 154:377–402. doi:[10.1144/gsjgs.154.3.0377](https://doi.org/10.1144/gsjgs.154.3.0377)
- Russell MJ, Hall AJ, Martin W (2010) Serpentinization as a source of energy at the origin of life. *Geobiology* 8:355–371. doi:[10.1111/j.1472-4669.2010.00249.x](https://doi.org/10.1111/j.1472-4669.2010.00249.x)
- Sakai H, Tsutsumi M (1978) D/H fractionation factors between serpentine and water at 100 °C to 500 °C and 2000 bars water pressure, and the D/H ratios of natural serpentines. *Econ Geol* 40:231–242
- Seyfried WE Jr, Dibble WE Jr (1980) Seawater-peridotite interaction at 300 °C and 500 bars: implications for the origin of oceanic serpentinites. *Geochim Cosmochim Acta* 44:309–321
- Seyfried WE Jr, Foustoukos DI, Fu Q (2007) Redox evolution and mass transfer during serpentinization: an experimental and theoretical study at 200 °C, 500 bar with implications for ultramafic-hosted hydrothermal systems at Mid-Ocean Ridges. *Geochim Cosmochim Acta* 71:3872–3886. doi:[10.1016/j.gca.2007.05.015](https://doi.org/10.1016/j.gca.2007.05.015)
- Sleep NH, Meibom A, Fridriksson T, Coleman RG, Bird DK (2004), H₂-rich fluids from serpentinization: geochemical and biotic implications. *Proc Natl Acad Sci U S A* 101:12818–12823
- Takai K, Gamo T, Tsunogai U, Nakayama N, Hirayama H, Nealson KH, Horikoshi K (2004) Geochemical and microbiological evidence for a hydrogen-based, hyperthermophilic subsurface lithoautotrophic microbial ecosystem (HyperSLiME) beneath an active deep-sea hydrothermal field. *Extremophiles* 8:269–282. doi:[10.1007/s00792-004-0386-3](https://doi.org/10.1007/s00792-004-0386-3)
- Takai K, Nakamura K, Suzuki K, Inagaki F, Nealson KH, Kumagai H (2006) Ultramafics-Hydrothermalism-Hydrogenesis-HyperSLiME (UltraH³) linkage: a key insight into early microbial ecosystem in the Archean deep-sea hydrothermal systems. *Paleontological Res* 10:269–282
- Van Dover CL, Humphris SE, Fornari D, Cavanaugh CM, Collier R, Goffredi SK, Hashimoto J, Lilley MD, Reysenbach AL, Shank TM, Von Damm KL, Banta A, Gallant RM, Gotz D, Green D, Hall J, Harmer TL, Hurtado LA, Johnson P, McKiness ZP, Meredith C, Olson E, Pan IL, Turnipseed M, Won Y, Young CR, Vrijenhoek RC (2001) Biogeography and ecological setting of Indian Ocean hydrothermal vents. *Science* 294:818–823. doi:[10.1126/science.1064574](https://doi.org/10.1126/science.1064574)
- Wakita H, Nakamura Y, Kita I, Fujii N, Notsu K (1980) Hydrogen release: new indicator of fault activity. *Science* 210:188–190. doi:[10.1126/science.210.4466.188](https://doi.org/10.1126/science.210.4466.188)
- Wenner DB, Taylor HP (1971) Temperatures of serpentinization of ultramafic rocks, based on ¹⁶O/¹⁸O fractionation between coexisting serpentine and magnetite. *Contrib Mineral Petrol* 32:165–185
- Wenner DB, Taylor HP (1973) Oxygen and hydrogen isotope studies of the serpentinization of ultramafic rocks in oceanic environments and continental ophiolite complexes. *Am J Sci* 273:207–239
- Wetzel LR, Shock EL (2000) Distinguishing ultramafic- from basalt-hosted submarine hydrothermal systems by comparing calculated vent fluid compositions. *J Geophys Res Solid Earth* 105:8319–8340. doi:[10.1029/1999JB900382](https://doi.org/10.1029/1999JB900382)
- Wiersberg T, Erzinger J (2008) On the origin and spatial distribution of gas at seismogenic depths of the San Andreas Fault from drill mud gas analysis. *Appl Geochem* 23:1675–1690. doi:[10.1016/j.apgeochem.2008.01.012](https://doi.org/10.1016/j.apgeochem.2008.01.012)
- Yoshizaki M, Shibuya T, Suzuki K, Shimizu K, Nakamura K, Takai K, Omori S, Maruyama S (2009) H₂ generation by experimental hydrothermal alteration of komatiitic glass at 300 °C and 500 bars: a preliminary result from on-going experiment. *Geochem J* 43: E17–E22