Erratum

The molecular systematics of some agar- and carrageenan-containing marine red algae based on *rbcL* sequence analysis

Suzanne Fredericq^{1,4}, Max. H. Hommersand² & D. Wilson Freshwater³

¹ Department of Botany, Duke University, Durham, NC 27708-0338, USA

² Department of Biology, University of North Carolina, Chapel Hill, NC 27599-3280, USA

³ Center for Marine Science Research, 7205 Wrightsville Avenue, Wilmington, NC 28403, USA

⁴ Present address: Laboratory of Molecular Systematics and Department of Botany, NHB-166, National Museum of Natural History, Smithsonian Institution, Washington, DC 20560, USA; e-mail: Frederic@onyx.si.edu

Hydrobiologia 326/327: 125–135, 1996.

S. C. Lindstrom & D. J. Chapman (eds), Fifteenth International Seaweed Symposium.

On p. 129 of the above article, Figure 3 was unfortunately omitted (a duplicate of Figure 2 appearing in its place). We now present Figure 3 below.

Figure 3. Single maximum parsimony tree of length = 803, CI = 0.540 and RI = 0.583, resulting from a multiple heuristic search analysis of *rbcL* sequence data from nine Halmeniaceae, five Schizymeniaceae and three Ceramiales (as outgroup) species. Decay indices are indicated above and bootstrap values (%) below branches.