- Löhden, J., Bernerth, R., Frentzen, M. (1990) Acyl-CoA:1-acylglycerol-3-phosphate acyltransferase from developing seeds of *Limnanthes douglasii* (I. Br.) and *Brassica napus* (L.). In: Plant lipid biochemistry, structure and utilization, pp. 175–179, Quinn, P.J., Harwood, J.L., eds. Portland Press, London
- Lynch, D.V., Steponkus, P.L. (1987) Plasma membrane lipid alterations associated with cold acclimation of winter rye seedlings (*Secale cereale* L. cv Puma). *Plant Physiol.* **83**, 761–767
- Ohlrogge, J.B., Kuhn, D.N., Stumpf, P.K. (1979) Subcellular localization of acyl-acyl carrier protein in leaf protoplasts of *Spinacia oleracea*. *Proc. Natl. Acad. Sci. USA* **76**, 1194–1198
- Oo, K.-C., Huang, A.H.C. (1989) Lysophosphatidate acyltransferase activities in the microsomes from palm endosperm, maize scutellum, and rapeseed cotyledon of maturing seeds. *Plant Physiol.* **91**, 1288–1295
- Rochester, C.P., Kjellbom, P., Andersson, B., Larsson, C. (1987) Lipid composition of plasma membranes isolated from light-grown barley (*Hordeum vulgare*) leaves: identification of cerebroside as a major component. *Arch. Biochem. Biophys.* **255**, 385–391
- Rock, C.O., Garwin, J.L. (1979) Preparative enzymatic synthesis and hydrophobic chromatography of acyl-acyl carrier protein. *J. Biol. Chem.* **254**, 7123–7128
- Roughan, P.G., Slack, C.R., Holland, R. (1976) High rates of [14 C]acetate incorporation into the lipids of isolated spinach chloroplasts. *Biochem. J.* **158**, 593–601
- Roughan, P.G., Slack, C.R. (1982) Cellular organization of glycerolipid metabolism. *Annu. Rev. Plant Physiol.* **33**, 97–132
- Sauer, A., Robinson, D.G. (1985) Subcellular localization of enzymes involved in lecithin biosynthesis in maize roots. *J. Exp. Bot.* **36**, 1257–1266
- Stymne, S., Stobart, A.K. (1984) Evidence for the reversibility of the acyl-CoA:lysophosphatidylcholine acyl-transferase in microsomal preparations from developing safflower (*Carthamus tinctorius* L.) cotyledons and rat liver. *Biochem. J.* **223**, 305–314
- Stymne, S., Bafor, M., Jonsson, L., Wiberg, E., Stobart, K. (1990) Triacylglycerol assembly. In: Plant lipid biochemistry, structure and utilization, pp. 191–197, Quinn, P.J., Harwood, J.L., eds. Portland Press, London
- Sparace, S.A., Moore, T.S. Jr. (1979) Phospholipid metabolism in plant mitochondria. Submitochondrial sites of synthesis. *Plant Physiol.* **63**, 963–972
- Vick, B., Beevers, H. (1977) Phosphatidic acid synthesis in castor bean endosperm. *Plant Physiol.* **59**, 459–463
- Verhoek, B., Haas, R., Wrage, K., Linscheid, M., Heinz, E. (1983) Lipids and enzymatic activities in vacuolar membranes isolated via protoplasts from oat primary leaves. *Z. Naturforsch.* **38c**, 770–777
- Yoshida, S., Uemura, M. (1986) Lipid composition of plasma membranes and tonoplasts isolated from etiolated seedlings of mung bean (*Vigna radiata* L.). *Plant Physiol.* **82**, 807–812

Erratum

Accumulation of phenolic conjugates and betacyanins, and changes in the activities of enzymes involved in feruloylglucose metabolism in cell-suspension cultures of *Chenopodium rubrum* L.

Maria Bokern¹, Victor Wray², and Dieter Strack¹

¹ Institut für Pharmazeutische Biologie der Technischen Universität Braunschweig, Mendelssohnstrasse 1, W-3300 Braunschweig, and

² GBF – Gesellschaft für Biotechnologische Forschung, Mascheroder Weg 1, W-3300 Braunschweig, Federal Republic of Germany

Planta (1991) 184:261–270

On page 264, right-hand column, sentence in line 7 to 10 in the legend of Fig. 2 should read:

“... Multiwavelength detection of 260 nm for the hydroxybenzoic acid conjugates (1 and 2), 320 nm for the hydroxycinnamic acid conjugates (7, 8, 10, 11, 12), and 540 nm for the betacyanins (3, 4, 6, 9, 13, 14) (0.128 absorption unit full scale). ...”