Erratum

Spinal abnormalities and the atypical form of the Mayer-Rokitansky-Küster-Hauser syndrome

Ernst H. Strübbe¹, J. Albert M. Lemmens², Cornelis J.P. Thijn³, Wim N.P. Willemsen⁴, Bert S.J. van Toor⁵

Skeletal Radiol (1992) 21:459–462

The authors of the above article would like to make the following corrections:

p. 460, Table 1, line 8, 3rd column: Spina bifida occulta

p. 461, left column, lines 28 and 29:

Table 1. Spinal anomalies as radiographic finding in 96 patients with Mayer-Rokitansky-Küster-Hauser (MRKH) syndrome

Anomalies	Number of features			
	Patients with atypical MRKH (N=52)		Patients with typical MRKH (N=44)	
	n	%	n	%
KF syndrome	14	26.9	—	
subtype 1	5	9.6	—	
subtype 2	6	11.5	—	
subtype 3	3	5.8	—	
Scoliosis	10	19.2	—	
Sacral agenesis	2	3.9	—	
Sacralization L5	5	9.6	1	2.3
Spina bifida occulta	25	48.1	4	9.1
Fusion L2–3	1	1.9	—	
No. of patients	37	71.2	5	11.4

The incidence of spina bifida occulta in **the typical group** is only slightly higher than

^a Fifty-seven features were found in 37 patients. Scoliosis combined with Klippel-Feil (KF) syndrome was found in 8 patients, while spina bifida occulta in combination with KF syndrome was seen in 12 patients.