- port and RuBP carboxylase activity in seedlings of *Pinus sylvestris*. *Physiol. Plant.* **48**, 526–531
- Öquist, G., Greer, D.H., Ögren, E. (1987) Light stress at low temperature. In: *Photoinhibition*, pp. 67–87, Kyle, D.J., Osmond, C.B., Arntzen, C.J., eds. Elsevier, Amsterdam
- Öquist, G., Chow, W.S., Anderson, J.M., (1992a) Photoinhibition of photosynthesis represents a mechanism for the long-term regulation of photosynthesis. *Planta* **186**, 450–460
- Öquist, G., Hurry, V.M., Öquist, M.G., Huner, N.P.A. (1992b) Differential resistance of frost-hardened and non-hardened winter rye to photoinhibition of photosynthesis is due to an increased capacity of frost-hardened rye to keep Q_A oxidized under similar light and temperature conditions. *Photosynthetica*, in press
- Osmond, C.B. (1981) Photorespiration and photoinhibition. Some implications for the energetics of photosynthesis. *Biochim. Biophys. Acta* **639**, 77–89
- Porra, R.J., Thompson, W.A., Kriedemann, P.E. (1989) Determination of accurate extinction coefficients and simultaneous equations for assaying chlorophyll a and b with four different solvents: verification of the concentration of chlorophyll by atomic absorption spectroscopy. *Biochim. Biophys. Acta* **975**, 384–394
- Powles, S.B. (1984) Photoinhibition of photosynthesis induced by visible light. *Annu. Rev. Plant Physiol.* **35**, 15–44
- Schöner, S., Krause, G.H. (1990) Protective systems against active oxygen species in spinach: response to cold acclimation in excess light. *Planta* **180**, 383–389
- Somersalo, S., Krause, G.H. (1989) Photoinhibition at chilling temperature. Fluorescence characteristics of unhardened and cold acclimated spinach leaves. *Planta* **177**, 409–416
- Tognetti, J.A., Salerno, G.L., Crespi, M.D., Pontis, H.G. (1990) Sucrose and fructan metabolism of different wheat cultivars at chilling temperatures. *Physiol. Plant.* **78**, 554–559
- van Kooten, O., Snel, J.F.H. (1990) The use of chlorophyll fluorescence nomenclature in plant stress physiology. *Photosyn. Res.* **25**, 147–150

Erratum

Observations on the subcellular distribution of the ammonium ion in maize root tissue using in-vivo ^{14}N -nuclear-magnetic-resonance spectroscopy

R.B. Lee¹ and R.G. Ratcliffe²

¹ AFRC Institute of Arable Crops Research, University of Bristol, Long Ashton Research Station, Bristol BS18 9AF, UK

² Department of Plant Sciences, University of Oxford, Agricultural Science Building, Parks Road, Oxford OX1 3PF, UK

Published in *Planta* (1991) 183:359–367

On page 360, right hand column, the second sentence in the legend for Fig. 1 should read:

“Spectra were recorded with (○) and without (●, □, ■, △) ^1H -decoupling from solutions containing 5 mM (○, △), 10 mM (○, ■), 20 mM (○, □) and 40 mM (○, ●) ammonium”.