Erratum

Expression of mouse *Tla* region class I genes in tissues enriched for $\gamma\delta$ cells

Pirootz Eghtesady, Kurt A. Brorson, Hilde Cheroutre, Robert E. Tigelaar, Leroy Hood, and Mitchell Kronenberg

Immunogenetics 36: 377–388, 1992

The following references were missing from the above-mentioned article:

- Stanton, T. and Boyse, E.: A new serologically defined locus, Qa-1, in the *Tla* region of the mouse. *Immunogenetics* 3: 525–532, 1976
- Steinmetz, M., Winoto, A., Minard K., and Hood L.: Clusters of genes encoding mouse transplantation antigens. *Cell* 28: 489–492, 1982
- Strominger, J.: The $\gamma\delta$ T-cell receptor and class Ib major histocompatibility molecules of immune recognition. *Cell* 57: 895–898, 1989
- Stroynowski, I., Soloski, M., Low M., and Hood, L.: A single gene encodes soluble and membrane-bound forms of the major histocompatibility Qa-2 antigen: anchoring of the product by a phospholipid tail. *Cell* 50: 759–768, 1987
- Stroynowski, I.: Molecules related to class-I major histocompatibility complex antigens. *Annu Rev Immunol* 8: 501–530, 1990
- Transy, C., Nash, S. R., David-Watine, B., Cochet, M., Hunt, S. W. III, Hood, L. E., and Kourilsky, P.: A low polymorphic mouse *H-2* class I gene from the *Tla* complex is expressed in a broad variety of cell types. *J Exp Med* 166: 341–361, 1987
- Vidovic, D., Roglic, M., McKune, K., Guerder, S., MacKay, C., and Dembic, Z.: *Qa-1* restricted recognition of foreign antigen by a $\gamma\delta$ T-cell hybridoma. *Nature* 340: 646–650, 1989
- Weiss, E., Golden, L., Fahrner, K., Mellor, A., Devlin, J., Bullman, H., Tiddens, H., Bud, H., and Flavell, R.: Organization and evolution of the class I gene family in the major histocompatibility complex of the C57BL/10 mouse. *Nature* 310: 650–655, 1984
- Widmark, E., Ronne, H., Hammerling, U., Serenius, B., Larhammar, D., Gustafsson, K., Bohme, J., Peterson, P., and Rask, L.: Family relationships of murine major histocompatibility complex class I genes. Sequence of the *T2A^a* pseudogene, a member of gene family 3. *J Biol Chem* 263: 7055–7059, 1988
- Wolf, P. R. and Cook, R. G.: The *TL* region gene 37 encodes a Qa-1 antigen. *J Exp Med* 172: 1795–1804, 1990
- Wu M., Van Kaer, L., Itohara, S., and Tonegawa, S.: Highly restricted expression of the Thymus Leukemia antigens on intestinal epithelial cells. *J Exp Med* 174: 213–218, 1991
- Wysocki, W. L. and Sato, V. L.: Panning for lymphocytes: A method for cell selection. *Proc Natl Acad Sci USA* 75: 2844–2848, 1978
- Yasumura, Y., Tashjian, A., and Sato, G.: Establishment of four functional, clonal strains of animal cells in culture. *Science* 154: 1186–1189, 1966