Letters to the editor

Type 1 (insulin-dependent) diabetes and genetic heterogeneity

Dear Sir,

We are glad that Wolf et al. [1] have confirmed several of our previous findings regarding the genetics of Type 1 (insulin-dependent) diabetes [2]. We add some important qualifications and additional data pertinent to their paper.

The main issue revolves around the question: is the genetic susceptibility to Type 1 diabetes due to a homogeneous state (one gene) or a heterogeneous state (several genes or polygenic) and, therefore, several aetiological pathways? An additional question is: is the genetic heterogeneity, if it exists, in the HLA region of chromosome 6 or outside it or both? We have always been a proponent of heterogeneity of both types (within and outside the HLA region), but we think it is important to qualify the strength of the available evidence. As Neil and Spielman [3] have stated, only evidence clearly incompatible with one gene models would represent definite evidence for heterogeneity.

Like Wolf et al., we have also found a large excess of the heterozygotes, DR3/DR4 versus other heterozygotes or homozygotes, DR3/DR3 or DR4/DR4 [2]. However, we also found that there was a small excess of homozygotes (DR3/DR3 or DR4/DR4) and the difference between these 'high risk' homo- and heterozygotes was not statistically significant. Since a large number of normal families has never been typed for DR, adequate control frequencies for homozygotes are lacking. Thus, although the apparent excess of heterozygotes DR3/DR4 over homozygotes is suggestive of the existence of two susceptibility genes in the HLA area, it is not proof. A similar conclusion has been reached by others [4].

Additional important evidence for genetic heterogeneity in Type 1 diabetes comes from linkage studies. A much tighter linkage between the Type 1 diabetic trait and HLA in DR3/DR4 probands than in other probands, recently reported by us [5, 6] and others [7], constitutes evidence for genetic heterogeneity that may extend beyond the HLA region, incorporating additional non-HLA-linked Type 1 diabetic determinants. This putative non-HLA-linked genetic susceptibility may be related to other associations found between the genetic markers, Gm [8] and Kidd [9], and Type 1 diabetes.

The inability of Wolf et al. to find a relationship between HLA and age at diagnosis confirms our findings [2]. Similarly, many groups, including our own [10], have shown that the frequency of HLA recombinants in Type 1 diabetes is normal. Further, it is true that HLA typing *does* identify siblings of probands at considerably higher risk of diabetes. However, this risk is still relatively small in absolute value (approximately 10%). For informative studies of the aetiology of Type 1 diabetes, which may lead to rational, preventative strategies, a large number (several thousand) of non-diabetic siblings of diabetic subjects with appropriate ages need to be studied prospectively for

immune, viral and metabolic parameters. HLA studies are likely to have only a secondary role in these prospective trials.

Yours sincerely,
J. Barbosa and S. Rich

References

1. Wolf E, Spencer KM, Cudworth AG (1983) The genetic susceptibility to Type 1 (insulin-dependent) diabetes: analysis of the HLA-DR association. *Diabetologia* 24: 224–230
2. Barbosa J, Bach FH, Rich SS (1982) Genetic heterogeneity of diabetes and HLA. *Clin Genet* 21: 25–32
3. Neil JV, Spielman R (1982) Evidence for genetic heterogeneity in insulin-dependent diabetes. A discussion of contrasting viewpoints. In: Kobberling J, Tattersall R (eds) *The genetics of diabetes mellitus*. Academic Press, London, pp 138–140
4. Svejgaard A, Ryder LP (1981) HLA genotype distribution and genetic models of insulin-dependent diabetes mellitus. *Ann Hum Genet* 45: 293–298
5. Dunsworth TS, Rich SS, Morton NE, Barbosa J (1982) Heterogeneity of insulin-dependent diabetes – new evidence. *Clin Genet* 21: 233–236
6. Morton NE, Green A, Dunsworth T, Svejgaard A, Barbosa J, Rich SS, Iselius L, Platz P, Ryder LP (1983) Heterozygous expression of insulin-dependent diabetes mellitus determinants in the HLA system. *Am J Hum Genet* 35: 201–213
7. Green A, Morton NE, Iselius L, Svejgaard A, Platz P, Ryder LP (1982) Genetic studies of insulin-dependent diabetes mellitus: segregation and linkage analyses. *Tissue Antigens* 19: 213–221
8. Schanfield MS, Brown R, Sarjeantson S, Kirk RL (1981) Susceptibility to juvenile onset diabetes in families: DR, Gm, Km, and gender. *Am J Hum Genet* 33: 133A
9. Barbosa J, Rich S, Dunsworth T, Swanson J (1982) Linkage disequilibrium between insulin-dependent diabetes and the Kidd blood group Jk^b allele. *J Clin Endocrinol Metab* 55: 193–195
10. Barbosa J, Chern MM, Anderson VE, Noreen H, Johnson S, Reinsmoen N, McCarty R, King R, Greenberg L (1980) Linkage analysis between the major histocompatibility system and insulin-dependent diabetes in families with patients in two consecutive generations. *J Clin Invest* 65: 592–601

Dr. Jose Barbosa
Department of Medicine
Box 716 Mayo Memorial Building
University of Minnesota
Minneapolis, MN 55455
USA

Erratum

Diabetologia, Volume 25, Number 6 (December), 1983, pages 457–459: Editorial by G. E. Sonnenberg and M. Berger

Title on front cover and in list of volume contents should read:

“Human Insulin: Much Ado About *One* Amino Acid?”

Last line of article (p. 458) should read:

“Human insulin *has* not made this difficult task any easier.”