Erratum

Neuropeptide Y inhibits nicotinic cholinergic currents but not voltage-dependent calcium currents in bovine chromaffin cells

Wolfgang Nörenberg¹, Peter Illes¹, and Kenneth Takeda²

¹University of Freiburg, Department of Pharmacology, Hermann-Herder-Strasse 5, W-7800 Freiburg i. Br., Federal Republic of Germany

²Université Louis Pasteur de Strasbourg, Laboratoire de Pharmacologie Cellulaire et Moléculaire-CNRS URA600, B.P. 24, F-67401 Illkirch, France

Pflügers Arch (1991) 418:346–352

Unfortunately, an error occurred in the order of the figures. Please note that Fig. 1 and Fig. 2 were interchanged though the respective legends are in the right places. The correct figures with their legends are printed below.

Fig. 1. **A** Effect of neuropeptide Y (NPY; 1–36) on I_{ACh} . I_{ACh} produced by consecutive pressure ejections of 30 μ mol/l nicotine before (T₁), during simultaneous application of 2 μ mol/l NPY (1–36) (T₂), and after washout (T₃). Drugs were applied for 20 s (horizontal bars). Trials were separated by 3 min of bath superfusion. Holding potential was –90 mV; GTP 200 μ mol/l was included in the patch pipette. ---, Zero current level. **B** Average values of peak amplitudes of nicotinic currents for three successive applications (T₁–T₃) of nicotine (30 μ mol/l) alone ($n = 6$ cells, 6 T₁–T₃ series; left panel) and when 2 μ mol/l NPY (1–36) was co-applied (T₂) with nicotine (30 μ mol/l; $n = 5$ cells, 8 T₁–T₃ series; right panel). * $P < 0.001$ compared to control (T₁)

Fig. 2A,B. Effect of NPY fragment 16–36 on I_{ACh} . **A** I_{ACh} produced by pressure applications of nicotine (30 μ mol/l) before (T₁), during co-administration of 2 μ mol/l NPY (16–36) (T₂), and after washout (T₃). Trials were separated by 3 min of bath superfusion. Holding potential –90 mV, 200 μ mol/l GTP included in the patch pipette. ---, Zero current level. **B** Average values of peak I_{ACh} induced by nicotine 10 μ mol/l (left panel; $n = 5$ cells, 5 T₁–T₃ series) and 30 μ mol/l (centre and right panels; $n = 5$ cells, 7 T₁–T₃ series; $n = 9$ cells, 11 T₂–T₃ series, respectively) alone or together with NPY (16–36) 2 μ mol/l (left and right panels) or 0.2 μ mol/l (centre panel). * $P < 0.001$ compared to controls (T₁)