Diabetic eye disease in Central Africa

M. Rolfe

Nkana Division, Zambia Consolidated Copper Mines Ltd, Kitwe, Zambia

Summary. Six hundred African diabetic patients were examined using a standardised technique based on the World Health Organisation Multinational Study, in which no country from Africa was represented. Thirty-four percent of patients had retinopathy and 13% were affected by cataract. They were older and the duration of diabetes was longer than patients without retinopathy or cataract. There was no association with glycaemic control. Systolic blood pressure was higher in patients with retinopathy but there was no as-

sociation with diastolic pressure. Only one patient had proliferative retinopathy and blindness compared to 25 (4.2%) patients blinded by cataract. In Central Africa cataract is a more important cause of blindness and visual disability than retinopathy.

Key words: Diabetes, Central Africa, retinopathy, cataract, blindness.

The reported prevalence of retinopathy and cataract [1–21] in African diabetic patients has varied widely (Table 1) due to the diversity of the methods and criteria used and failure to take the age of the patient and duration of diabetes into account. In 1979 the World Health Organisation (WHO) reported on a Multinational Study of diabetes [22] the object of which was to compare the prevalence of microvascular and macrovascular disease in different populations using a standardised questionnaire and examination technique. Although 14 centres throughout the world participated, no country could be found to represent the African continent. This paper reports on retinopathy and cataract in diabetic patients in Central Africa using a standardised technique based on the WHO protocol.

Subjects and methods

Patients recruited were ethnic Africans, who were currently being treated for diabetes in general medical clinics at seven hospitals belonging to Zambia Consolidated Copper Mines Ltd (ZCCM) and at two Government Hospitals, Kitwe Central and Ndola Central Hospitals, all situated in the Copperbelt region of Zambia. Registers of diabetic patients were not kept but patients attended hospital rather than private practitioners in order to receive free medicine. All diabetic patients currently attending for treatment were personally seen and examined by the author. Most patients spoke and understood English but for those not able to understand a Zambian nurse translated and interpreted the local dialect.

Table 1. Reported prevalence of retinopathy and cataract in African diabetic patients

Country	Retinopathy		Cataract	
	Reference	Prevalence (%)	Reference	Prevalence (%)
Ethiopia	1	7	1	9.5
	2	38	—	—
Kenya	3	52	—	—
Malawi	—	—	4	3
Nigeria	5	8	5	22
	6	7.5	6	12.5
	7	4.6	8	8.7
	9	25	13	10.6
South Africa	10	11.45	14	9.8
	11	25	—	—
	12	14	—	—
	13	12	—	—
Tanzania	15	13.6	15	41
	16	24.6	16	12.3
Uganda	17	4	17	10.5
	18	8.6	18	15
Zimbabwe	19	4	19	10.7
	20	6	—	—
	21	8	—	—

A questionnaire was completed for each patient, on which was recorded the name, sex, date, tribal origin, present age and age at diagnosis of diabetes. The exact age was not always known, especially by older women, and occasionally had to be estimated from the patient's physical appearance, after consultation with the attending Zambian nurse. The duration of the disease was calculated from the

Table 2. Numbers, mean age, duration of diabetes and body mass index (BMI) by sex and type of diabetes

	Type		Non-ketotic insulin-dependent	Overall
	I	II		
<i>Men</i>				
No.	106	132	134	372
Mean age (years)	34.3 SD 14.8	49.8 SD 8.3	49.7 SD 10.3	45.4 SD 13.2
Mean duration of diabetes (years)	5.6 SD 5.6	5.5 SD 4.8	7.2 SD 5.9	6.1 SD 5.5
BMI	21.1 SD 3.7	25.1 SD 4.0	23.0 SD 4.0	23.2 SD 4.3
<i>Women</i>				
No.	63	96	69	228
Mean age (years)	29.2 SD 12.6	49.7 SD 10.5	47.8 SD 9.6	43.5 SD 14.0
Mean duration of diabetes (years)	5.1 SD 5.0	5.1 SD 4.2	5.9 SD 4.7	5.3 SD 4.6
BMI	23.2 SD 5.0	26.5 SD 5.6	26.8 SD 5.2	25.7 SD 5.5

age at diagnosis, with information given by the patient and from hospital records. Details of current treatment were recorded. The type of diabetes was categorised into three groups. Type I patients were all those receiving insulin who had documented ketosis at some time. Type II patients were those on oral hypoglycaemic drugs or treatment with diet alone. Patients who were receiving insulin but who had had no documented ketosis were regarded as having non-ketotic insulin-dependent diabetes. Visual acuity was recorded for each eye using a Snellen chart, without glasses (if worn). This was done because glasses tend to be passed on from person to person in Africa and many patients did not attend with their spectacles. Mydriatic drops (1% tropicamide or 1% cyclopentolate) were then instilled into each eye. Fundus examination was done when the pupil was at least 3 mm dilated by direct ophthalmoscopy in a darkened room. The patient's gaze was fixed except for examination of the macula when the patient was instructed to look directly at the light. Each fundus was examined for 2 min using a timing device. Extra time was allowed, if necessary, to count the number of lesions. The presence or absence of the lens was recorded. If the retina could not be clearly seen the reason was stated. The protocol for assessing retinopathy was exactly as used in the WHO Study [22]. As it is often difficult, or impossible, to differentiate between a haemorrhage and a micro-aneurysm, the general descriptive term, red lesion, was used. A small red lesion was defined as having a diameter less than the retinal artery at the optic disc. A medium red lesion was larger, but with its diameter less than that of the optic disc; while a large red lesion was defined as one with its largest diameter greater than that of the optic disc. They were counted for each eye and categorised as follows: 1 = none, 2 = one, 3 = two-five, 4 = six or more. Hard and soft exudates were also counted and categorised in the same way. Large or extensive exudates with a total area greater than the optic disc were included in the "six or more" category. The presence of new vessels was recorded as absent, doubtful or definite. Microvascular disease was classified according to the type, number and size of abnormalities seen. Minimal small vessel disease of the eye was defined as one small red lesion in any eye; or one hard or soft exudate in any eye; or doubtful new vessels in any eye. Medium small vessel disease was two or more small red lesions in any eye; or one or more medium red lesions in any eye; or two or more hard or soft exudates in any eye. Severe small vessel disease included retinopathic blindness in any eye; and/or new vessels in any eye; and/or vitreous opacity or haemorrhage in any eye; and/or one or more large red lesions in any eye.

Blood pressure was measured using a mercury sphygmomanometer with a cuff measuring 23 × 14 cm. The pressure was measured in the right arm with the patient sitting. The systolic pressure was recorded as the first perception of successive sounds. The diastolic pressure was taken at the fifth-phase at which sounds disappeared. If this was not audible the fourth-phase muffle was taken. All readings were recorded in mmHg to the nearest 2 mm, reading the calibration below the meniscus.

The height of each patient was recorded in centimetres without shoes and the weight, without coat or shoes, in kilograms. The body mass index (BMI) was calculated according to the formula BMI = weight in kilograms divided by the height in metres squared.

Venous blood was taken from each patient. Glycosylated haemoglobin (HbA_{1c}) was measured using a micro-column chromatographic method (Boehringer, Cat. No. 575453, Mannheim, FRG) after elimination of the labile fraction (normal range = 5–8%). Internal quality control was done with each batch. The final HbA_{1c} values were corrected for temperature using the table provided. Haemoglobin concentration, reticulocyte count and haemoglobin electrophoretic pattern were determined. HbA_{1c} estimations were not done in those with elevated serum creatinine, evidence of haemolytic anaemia and patients heterozygous for sickle cell anaemia as these conditions produce artefactual results [23, 24].

Statistical Analysis

Statistical analysis was done using the Student's t-test or the chi-square test [25].

Results

Six hundred African diabetic patients (372 men, 228 women) were examined. There was no difference in age or duration of diabetes between men and women (Table 2) but women were more obese ($p < 0.001$).

Retinopathy

Two hundred and three (34%) patients were found to have small vessel disease of the eye (SVDE), comprising 132 (35%) men and 71 (31%) women. Forty-four (26%) patients had Type I diabetes, 82 (36%) Type II and 77 (38%) non-ketotic insulin-dependent diabetes. The prevalence of retinopathy rose dramatically with duration of diabetes; 74% of those with disease longer than 14 years had retinopathy (Table 3). Patients with SVDE had had diabetes longer ($p < 0.001$) and were older ($p < 0.05$) than patients without SVDE (Table 4). There was no difference ($p > 0.05$) in the degree of glycaemic control between those with and without SVDE (Table 5). However, there was an association with

Table 3. Prevalence of small vessel disease of the eye (SVDE) by duration of diabetes

	Duration (years)	No.	Percent
Minimal SVDE	0-6	25	6.2
	7-13	18	12.9
	14+	5	8.8
Medium SVDE	0-6	59	14.6
	7-13	44	31.4
	14+	28	49.1
Severe SVDE	0-6	8	2.0
	7-13	7	5.0
	14+	9	15.8
All patients			No. + % with SVDE
	0-6	403	92 (22.8)
	7-13	140	69 (49.3)
	14+	57	42 (73.7)
Total		600	203 (34%)

Table 4. Numbers, mean age and duration of diabetes in patients with and without small vessel disease of the eye (SVDE) and cataract

	Men	Women	Overall	Without
SVDE present	132 (35%)	71 (31%)	203 (34%)	397
Mean age (years)	49.5	48.8	49.2 SD 12.0	42.4 SD 13.7
Mean duration of disease (years)	8.9	8.1	8.6 SD 5.9	4.4 SD 4.1
Cataract present	34	30	64 (10.7%)	523
Cataract removed	7	6	13 (2.2%)	-
Mean age (years)	54.9	53.3	54.2 SD 13.0	43.3 SD 13.1
Mean duration of disease (years)	8.2	7.6	7.9 SD 6.2	5.5 SD 4.9

blood pressure; the systolic pressure was higher in both men and women with SVDE ($p < 0.01$) compared to those without (Table 6) but there was no difference ($p > 0.05$) in the diastolic pressure.

There was no difference ($p > 0.05$) in body mass index between those with SVDE and those without (Table 7). There were only two patients with retinopathic blindness. One patient had severe proliferative retinopathy leading to vitreous haemorrhage while the other had detachment of the retina not associated with her diabetes (Table 8). Forty-four (21.7%) patients with SVDE had visual disability (6/18-6/60) compared to 46 (11.6%) of those without ($p < 0.01$); 37% of patients with retinopathy also had proteinuria.

Cataract

One hundred and thirty-four (11.2%) of all diabetic eyes were affected by cataract while 38 (3.2%) were blind.

Table 5. Glycaemic control (mean HbA_{1c}%) in patients with and without small vessel disease of the eye (SVDE) and cataract

	Men (No.)	Women (No.)
With SVDE	9.7 SD 3.1 (113)	9.8 SD 2.3 (58)
Without SVDE	9.6 SD 3.0 (187)	9.4 SD 2.7 (126)
With cataract	9.7 SD 3.2 (38)	10.3 SD 2.7 (31)
Without cataract	9.6 SD 3.0 (262)	9.3 SD 2.6 (153)

Table 6. Relationship of blood pressure (mmHg) to small vessel disease of the eye (SVDE)

	Men	Women
Systolic with SVDE	143.6 SD 30.2	146.1 SD 28.2
Systolic without SVDE	134.6 SD 26.4	134.0 SD 28.6
Diastolic with SVDE	80.3 SD 15.8	82.8 SD 13.1
Diastolic without SVDE	80.3 SD 14.5	80.3 SD 14.7

Table 7. Body mass index in patients with and without small vessel disease of the eye

	Without	Minimal	Medium	Severe
Men	23.2 SD 4.4	23.4 SD 4.4	23.5 SD 4.1	21.1 SD 3.8
No.	240	27	89	16
Women	25.7 SD 5.7	26.7 SD 6.5	25.5 SD 4.7	24.7 SD 3.9
No.	157	21	42	8

Table 8. Visual acuity in patients with small vessel disease of the eye (SVDE) or cataract

	1	2	3	4	5	6
Minimal SVDE	40	5	2	0	1	0
Medium SVDE	96	22	7	0	5	1
Severe SVDE	15	6	2	1	0	0
All SVDE	151	33	11	1	6	1
Cataract present or removed	9	24	16	2	25	1

1 = 6/6 - 6/12; 2 = 6/18 - 6/36; 3 = 6/60; 4 = retinopathic blindness; 5 = blindness from cataract; 6 = blindness other reasons

Cataract was present in 34 men and 30 women, whilst a further 7 men and 6 women had had cataracts removed. Thus 12.8% of the diabetic study population was affected (Table 4). They were older ($p < 0.001$) and had had diabetes longer ($p < 0.01$) than patients without cataract. There was no difference ($p > 0.05$) in mean HbA_{1c} levels between patients with cataract and those without (Table 5). Blindness, defined as the inability to count fingers at one arm's length, affected 25 (4.2%) patients (bilaterally in 13, unilaterally in 12). Their age and duration of diabetes were no different ($p > 0.05$) from other patients with cataract (Table 9).

Table 9. Blindness and cataract: numbers, mean age and duration of diabetes

	Women	Men	Overall
Bilateral blindness			
Numbers	7	6	13 (2.2%)
Mean age (years)	48.3	48.7	48.5 SD 14.3
Mean duration (years)	4.3	8.3	6.2 SD 6.5
Unilateral blindness			
Numbers	5	7	12 (2.0%)
Mean age (years)	63.8	52.1	57.0 SD 19.2
Mean duration (years)	10.0	9.9	9.9 SD 7.2
All blind patients			
Numbers	12	13	25 (4.2%)
Mean age (years)	54.8	50.5	52.6 SD 16.8
Mean duration (years)	6.7	9.2	8.0 SD 6.9

Only 9 patients with cataract had normal vision (Table 8).

There were 20 patients with bilateral cataracts in whom neither retina could be adequately visualised whilst 10 (52.6%) of 19 patients with unilateral cataract- and a non-assessable fundus had evidence of retinopathy in the other eye. Altogether 27 (35.1%) patients with cataract also had retinopathy.

Discussion

Many reports on retinopathy in African diabetic patients fail to give adequate details on the methods employed and how the patients were selected with reference to age and duration of diabetes. In this study a standardised examination procedure was used, identical to the method employed in the WHO Multinational Study [22]. As the author personally examined all patients inter-observer variation was avoided. The prevalence rate of 34% showed that SVDE was as common in Zambian diabetic patients as in other countries in the Multinational Study [22], despite the relatively short duration of disease. However, retinopathy in patients with cataract may have been under-diagnosed as only direct ophthalmoscopy was employed. The low numbers with severe SVDE reflected the small number of diabetic patients recruited with long duration disease, which is the most important factor in the aetiology of microvascular disease [26]. The association of retinopathy with systolic blood pressure was also found in the WHO Study although there was variation between centres. There was a high prevalence of hypertension in Zambian diabetic patients (39%) which increased with age. The higher systolic pressure in patients with retinopathy may be partly explained by the older age of these patients, in addition to the effect of renal disease in raising blood pressure.

The prevalence of cataract (13%) in this study was similar to that reported from other African countries.

Five of the 25 patients blind from cataract were under 30 years of age and visual acuity was seriously impaired (6/18–6/60) in a further 40 (6.7%) patients. Altogether 11% of the diabetic population under review had impaired vision from cataract. In developed countries diabetic retinopathy is a major cause of blindness; and in England and Wales is the most common cause of blindness between the ages of 30 and 64 years [27]. In Central Africa, however, cataract is a far more important cause of blindness and visual handicap as few diabetic patients live long enough to develop severe SVDE and the scarcity of skilled medical resources means that the majority of patients affected never receive corrective treatment.

Acknowledgements. The data in this paper formed part of an M.D. thesis submitted to the University of London. I am grateful to Prof. R.J. Jarrett of Guy's Hospital Medical School, London and Prof. K.G.M.M. Alberti of University of Newcastle Medical School for much helpful advice; to Mr. T. Mulele, Chief Laboratory technician, Kalulushi Mine Hospital for performing HbA₁ levels; to Boehringer-Mannheim Corporation who donated the kits for measuring HbA₁ levels; and to my wife for typing the manuscript.

References

1. Lester FT (1984) The clinical pattern of diabetes mellitus in Ethiopians. *Diabetes Care* 1: 6–11
2. Lester FT (1983) Longstanding diabetes mellitus in Ethiopia: a survey of 105 patients. *Diabetologia* 25: 222–225
3. Steel JM, Awan AM, Mngola EN (1977) Diabetic retinopathy in Kenya. *Trop Doct* 7: 12–14
4. Goodall JWD, Pilbeam STHH (1964) Diabetes in Nyasaland. *Trans R Soc Trop Med Hyg* 6: 575–578
5. Kinnear TWG (1963) The pattern of diabetes mellitus in a Nigerian Teaching Hospital. *East Afr Med J* 5: 288–294
6. Greenwood BM, Taylor JR (1968) The complications of diabetes in Nigerians. *Trop Geogr Med* 20: 1–12
7. Osuntokun BO (1969) Diabetic retinopathy in Nigerians. *Br J Ophthalmol* 53: 652–663
8. Osuntokun BO, Akinkugbe FM, Francis TI, Reddy S, Osuntokun O, Taylor GOL (1971) Diabetes mellitus in Nigerians. A study of 832 patients. *West Afr Med J* 20: 295–312
9. Jackson WPU, Goldin C, Marine N (1966) Diabetics: inter-racial comparisons II. *S Afr Med J* 40: 206–208
10. Seftel HC, Walker ARP (1966) Vascular disease in South African Bantu diabetics. *Diabetologia* 2: 286–290
11. Morley JE, Lowenthal MN, Asvat MS, Kopelowitz W, Klein C, Kokoris N (1977) Problems experienced in a diabetic clinic for Blacks. *S Afr Med J* 8: 215–218
12. Omar MAK, Asmal AC (1984) Complications of early-onset insulin-dependent diabetes mellitus in Blacks and Indians. *S Afr Med J* 65: 75–78
13. Gill GV, Huddle KR, Krige LP (1984) Intensive health screening of young Black diabetics. *S Afr Med J* 20: 815–816
14. Gill GV, Huddle KR (1984) Patterns and problems of diabetes in Soweto. *Cent Afr J Med* 10: 189–195
15. Haddock DRW (1964) Diabetes and its complications in Dar-es-Salaam. *East Afr Med J* 4: 145–155
16. Mhando PA, Yudkin JS (1980) The pattern of diabetic complications in African patients in Dar-es-Salaam. *Trop Geogr Med* 32: 317–323
17. Shaper AG (1958) The pattern of diabetes in Africans in Uganda. *Proceedings International Diabetes Federation, 3rd Congress. Excerpta Medica, Düsseldorf*, pp 664–669

18. Otim MA (1975) Preliminary observations on diabetic retinopathy in Ugandan Africans attending Mulago diabetic clinic. *East Afr Med J* 2: 63-69
19. Gelfand M, Forbes JI (1963) Diabetes mellitus in the Rhodesian African. *S Afr Med J* 37: 1208-1213
20. Levin L, Gelfand M (1973) Diabetic retinopathy in African patients. *S Afr Med J* 47: 993-994
21. Wicks ACB, Jones JJ (1974) Diabetes mellitus in Rhodesia: a comparative study. *Postgrad Med J* 50: 659-663
22. World Health Organisation (1979) Vascular Disease in Diabetics NCD/OND/79.4
23. Peacock I (1984) Glycosylated haemoglobin: measurement and clinical use. *J Clin Pathol* 37: 841-851
24. Aleyassine H (1979) Low proportions of glycosylated haemoglobin associated with haemoglobin S and haemoglobin C. *Clin Chem* 8: 1484-1486
25. Swinscow TDV (1978) *Statistics at Square One*. British Medical Association, London
26. West KM (1978) *Epidemiology of Diabetes and its Vascular Lesions*. Elsevier, New York
27. Kohner EM, McLeod D, Marshall J (1982) Diabetic eye disease. In: Keen H, Jarrett RJ (eds) *Complications of Diabetes*. E Arnold, London, pp 19-108

Received: 30 June 1987
and in revised form: 1 December 1987

Dr. M. Rolfe
Wick Manor
Curry Rivel
Langport
Somerset TA10 ONW
UK