- Saxena, S. K., Ghose, S.: Mg^{2+} - Fe^{2+} order disorder and the thermodynamics of the orthopyroxene crystalline solution. *Am. Mineralogist* **56**, 532-550 (1971)
- Sen, S. K., Chakraborty, K. R.: Magnesium-iron exchange equilibrium in garnet-biotite and metamorphic grade. *Neues Jahrb. Mineral. Abhandl.* **108**, 181-207 (1968)
- Shannon, R. D., Prewitt, C. T.: Effective ionic radii in oxides and fluorides. *Acta Cryst. B* **25**, 925-946 (1969)
- Skinner, B. J.: Physical properties of the end-members of the garnet group. *Am. Mineralogist* **41**, 428-436 (1956)
- Warner, R. D., Luth, W. C.: Two-phase data for the join monticellite ($CaMgSiO_4$) - Forsterite (Mg_2SiO_4): experimental results and numerical analysis. *Am. Mineralogist* **58**, 998-1008 (1973)
- White, W. B., Moore, R. K.: Interpretation of spin-allowed bands of Fe^{2+} in silicate garnets. *Am. Mineralogist* **57**, 1692-1710 (1972)
- Zeeman, J.: Zur Kristallchemie der Granate. *Beitr. Mineral. Petrol.* **8**, 180-188 (1962)

Dr. J. Ganguly
Institute of Geophysics
University of California
Los Angeles, Calif. 90024
USA

Contrib. Mineral. and Petrol. **48**, 148 (1974)
© by Springer-Verlag 1974

Alkali Amphiboles Intermediate in Composition between Actinolite and Riebeckite

Christos Katagas

Department of Geology, The University of Manchester, Manchester M13 9PL, England

Addendum to Contrib. Mineral. Petrol. **46**, 257-264 (1974). The following additional analyses of the coexisting phases were obtained: chlorite: $(Mg_{2.72}Al_{1.24}Fe_{2.01})(Al_{1.13}Si_{2.87})O_{10}(OH)_8$, epidote $Ca_{1.93}(Al_{2.04}Fe_{0.95})(Si_{2.97})O_{12}(OH)$ and albite $Na_{0.90}Ca_{0.04}Al_{0.98}Si_{3.04}O_8$.

C. Katagas
Department of Geology
The University of Manchester
Manchester M13 9PL, England