tion of clusters, as well as the presence of single cells, the absence of bullous pemphigoid antigen, and vimentin would further argue for them being Merkel cells [10, 12–14].

Since Merkel cells are thought to be highly specialized keratinocytes [10, 14, 15] it is remarkable to find them in a tissue of altered epidermal differentiation, e.g., as in psoriasis. In order to gain a better understanding of the pathogenetic relevance of these cells, further studies should investigate their immunoelectromicroscopic morphology and perhaps determine what proportion of them form into psoriatic plaques.

Acknowledgements. We are grateful to Drs. Dráberová and Dráber, Prague, for VI-01, and for the helpful advice of Professor Saurat, Geneva, and Dr Kasper, Görlitz, and the technical assistance of Mrs Sabine Feldrappe. This study was supported by GDR research grant M51.

References

1. Bártek J, Bárteková J, Taylor-Papadimitriou J, Rejthar A, Kovarik J, Lukás Z, Vojtesek B (1986) Differential expression of keratin 19 in normal human epithelial tissues revealed by monospecific monoclonal antibodies. *Histochem J* 18:565–575
2. Breathnach AS (1971) Embryology of human skin. A review of ultrastructural studies. *J Invest Dermatol* 57:133–143
3. Dráberová E, Dráber P, Viklický V (1987) Characterization of monoclonal antibodies against vimentin. *Wiss Z Friedrich-Schiller-Univ Jena, Naturwiss R* 36:237–240
4. Hashimoto K (1972) Fine structure of Merkel cells in human oral mucosa. *J Invest Dermatol* 58:381–387
5. Karsten U, Papsdorf G, Roloff G, Stolley P, Abel H, Walther I, Weiss H (1985) Monoclonal anticytokeratin antibody from a hybridoma generated by electrofusion. *Eur J Cancer Clin Oncol* 23:137–147
6. Karsten U, Papsdorf G, Roloff G (1987) Herstellung und Charakterisierung von drei monoklonalen Antikörpern gegen Zytokeratine. *Wiss Z Friedrich-Schiller-Univ Jena, Naturwiss R* 36:230–232
7. Kasper M, Stosiek P, Typlt H, Karsten U (1987) Histological evaluation of three monoclonal anti-cytokeratin antibodies 1. Normal tissues. *Eur J Cancer Clin Oncol* 23:137–147
8. Kasper M, Stosiek P, Karsten U (1987) Gewebecharakteristik von drei neuen monoklonalen Cytoskelett-Antikörpern. *Wiss Z Friedrich-Schiller-Univ Jena, Naturwiss R* 36:233–236
9. Kidd RL, Krawczyk WS, Wilgram GF (1971) The Merkel cell in human epidermis: Its differentiation from other dendritic cells. *Arch Dermatol Forsch* 241:374–388
10. Mérot Y, Carraux P, Didierjean L, Saurat J-H (1986) Merkel cells do not express bullous pemphigoid antigen. *Dermatologica* 172:69–76
11. Moll R, Franke WW, Schiller DL, Geiger B, Krepler R (1982) The catalog of human cytokeratins: patterns of expression in normal epithelia, tumors and cultured cells. *Cell* 31:11–24
12. Saurat J-H, Didierjean L (1984) The epidermal Merkel cell is an epithelial cell. *Dermatologica* 169:117–120
13. Saurat J-H, Chavaz P, Carraux P, Didierjean L (1983) A human monoclonal antibody reacting with Merkel cells: immunofluorescence, immunoperoxidase, and immunoelectron microscopy. *J Invest Dermatol* 81:249–253
14. Saurat J-H, Didierjean L, Skalli O, Siegenthaler G, Gabbiani G (1984) The intermediate filament proteins of rabbit normal epidermal Merkel cells are cytokeratins. *J Invest Dermatol* 83:431–435
15. Tweedle CD (1978) Ultrastructure of Merkel cell development in aneurogenic and control amphibian larvae (*Ambystoma*). *Neuroscience* 3:481–486

Received December 14, 1987

Erratum

Arch Dermatol Res (1988) 280:67–70

Effect of exogenous lipids on in vitro transepidermal water loss and percutaneous absorption

S. J. Moloney

There were unfortunately two serious printing errors in the above paper. On page 69, right column, line 15 from the top, “dioleoyl phosphatidylglycerol” should be deleted. On page 69, the fourth item in Table 2 should have been “Dioleoyl phosphatidylcholine” instead of “Dibehenoyl phosphatidylcholine”. Please correct the text in your copy of the issue.

Announcements

7th Annual Postgraduate Course in Medical Mycology (Dermatomyology)

Date: October 14–16, 1988.

Place: University of California, San Francisco.

Credit: AMA/CMA/ACCME Category I Credit

For more information, please contact:

Extended Programs in Medical Education
University of California San Francisco
Room U-569
San Francisco, CA 94143-0742, USA
Telephone: (415) 476-4251

International Dermatology Symposium on Interferons and Related Lymphokines – Basic Research Basic Research and Applied Therapy

October 11–14, 1989 in Berlin.

For further information:

Priv. Doz. Dr. R. Stadler
Universitätsklinikum Steglitz
Hautklinik
Hindenburgdamm 30
D-1000 Berlin 45