Erratum

Randomized Trial of Two Beta-Mimetic Drugs (Ritodrine and Fenoterol) in Acute Intra-Partum Tocolysis

J. Gerris¹, M. Thiery¹, M. Bogaert², and A. De Schaepdryver²

¹Department of Obstetrics and ²Heymans Institute of Pharmacology, University of Gent Medical School, Gent, Belgium

Eur. J. Clin. Pharmacol. 18, 443–448 (1980)

On page 445 the legends for Figs. 1 and 2 were exchanged. They should have been printed as follows:

Fig. 1. Uterine activity, expressed as percentage of pre-infusion value, during and after a 30 min infusion of fenoterol 1, 2, or 4 µg/min. Each dose was given to 4 women; mean $\pm$ SEM

Fig. 2. Uterine activity, expressed as percentage of pre-infusion value, during and after a 30 min infusion of ritodrine 100, 200, or 400 µg/min. Each dose was given to 4 women; mean $\pm$ SEM

Information

The **Paul Martini Prize** of DM 20,000 is awarded in remembrance of Professor Paul Martini in appreciation of his contributions to the promotion of clinical therapeutical research.

The papers submitted must be self-contained, and published papers must not be older than two years. The papers should be submitted either in German or English to the Medizinisch Pharmazeutische Studiengesellschaft e.V., Bilhildisstrasse 2, D-6500 Mainz, where further information about the presentation will be available. The deadline for submission is April 28, 1981.