be denoted by $R(L^+)$. If $(x, y) \in R(D)$, then we denote by $T(L^+)$ the interval $0 < z < a_0$, where a_0 is the low extreme point of $Q(B)$ nearest the plane $z = 0$, lying on L^+ . We denote by $N(D)$ the union of all $R(L^+)$ and $T(L^+)$ when (x, y) traverses the entire plane $z = 0$. We denote by $M(D)$ the connected component of $N(D)$ containing $R(D)$, and by $M_0(D)$ the union of all $T(L^+)$ when (x, y) traverses $R(D)$. Clearly $M_0(D) \subset M(D)$, and the union of set $M_0(D)$ with the set $M(D)$, symmetric with respect to the plane $z = 0$, coincides with $H(D)$. In the examples considered above the sets $M_0(D)$ and $M(D)$ coincide.

The lower extreme points of $Q(B)$ can be found by means of equations (23) only in the case where domain D is bounded. If domain D is not bounded, then relations (23) are not obtained from (21). In this case it is necessary to do otherwise. As an example we shall consider the case of a tubular domain. Let $x = x_1 + ix_2$, $y = y_1 + iy_2$. Domain D is defined as follows: point (x_1, y_1) traverses a flat plane $R(D)$, and $-\infty < x_2, y_2 < \infty$. In order to find $H(D)$ we proceed as follows. From the first equation of (21) it follows that there exists a real number λ such that $\beta = \lambda(\gamma - y)$, $\delta = -\lambda(\alpha - x)$. After computing these equations the second equation of (21) takes the form

$$z^2 = \frac{(\lambda^2 + 1)^2}{\lambda^2 - 1} [(\alpha - x)^2 + (\gamma - y)^2].$$

So long as (x, y) is an internal point of domain $R(D)$ and (α, γ) is a boundary point of this domain, the value of $\min [(\alpha - x)^2 + (\gamma - y)^2]$ is the square of the distance from point (x, y) to the boundary of domain $R(D)$. The value of $(\mu + 1)^2 \cdot (\mu - 1)^{-1}$ reaches the positive minimum 8 for $\mu = 3$. Therefore, for the coordinate z of the boundary point of domain $H(D)$ lying over (x, y) we have

$$z^2 = 8 \min_{\alpha, \gamma} [(\alpha - x)^2 + (\gamma - y)^2]. \quad (25)$$

In particular, if $R(D)$ is a circle $x^2 + y^2 < R^2$, then from (25) we obtain

$$z^2 = 8(R - \sqrt{x^2 + y^2})^2,$$

i.e., the domain $H(D)$ satisfies the inequality

$$z^2 < 8(R - \sqrt{x^2 + y^2})^2.$$

The integral representation (18) of the solution to the Cauchy problem (20) for Eq. (19) can be used exactly the same way as the integral representation of (3) for studying various questions dealing with the analytic extension of harmonic functions and the convergence of series of harmonic functions. These integral representations make possible the broad application of the apparatus of the theory of functions in several complex variables for the study of partial differential equations. The integral representation (18) and its corollaries can be generalized for harmonic functions of more than three independent variables.

LITERATURE CITED

1. A. Janushauskas, "The Cauchy problem for the Laplace equation and the expansion operation for harmonic functions," Dokl. Akad. Nauk SSSR, 159, No. 2, 286-289 (1964).
2. A. Krattser and B. Frants, Transcendental Functions, Izd. -vo Inostr. Lit., Moscow (1963).
3. B. A. Fuks, Special Chapters in the Theory of Analytic Functions of Several Complex Variables, Fizmatgiz, Moscow (1963).
4. H. Behnke and P. Thullen, Theorie der Funktionen Mehrerer Komplexer Veränderlichen, Springer, Berlin - Heidelberg - New York (1970).
5. F. Lösch, "Lückensätze und Überkonvergenz bei zweifachen Potenzreihen," Math. Z., 36, 202-262 (1933).

ERRATA

In the article entitled "A generalized Carleman problem with inverse translation for a simply connected domain," by V. I. Slizkii from Sibirsk Matem. Zh., 16, No. 1, 150-152 (1975), on p. 152, line 7 from the top the equation $\kappa = 1/2$ and $[\Delta(t)]$ should read as follows: $\kappa = \text{ind } \Delta(t)$.