coalescence to the neutrino luminosity on the background of the process of single-photon production of neutrinos on a nucleus [2]. In order of magnitude, the power of the neutrino emission can be obtained from (12) by replacing I by I_0 (see also Fig. 1). This gives

$$S_n \sim \alpha^3 (Ze)^2 (G\gamma)^2 (n\lambda_c^3) (kT/m)^{14} (77C_V^2 + 13C_A^2) m^5, \quad (16)$$

where λ_c is the electron Compton wavelength. The relative contribution of the mechanisms of neutrino production to the luminosity is determined in order of magnitude by the expression ($C_V \sim C_A \sim 1$)

$$\frac{S_n[\gamma\gamma(Ze) \rightarrow \nu\bar{\nu}(Ze)]}{S_n[\gamma(Ze) \rightarrow \nu\bar{\nu}(Ze)]} \sim 10^{-3} \alpha \left(\frac{B}{B_0}\right)^2 \left(\frac{kT}{m}\right)^4. \quad (17)$$

Taking for an estimate $kT \sim m$, we find that for $B \sim 10^3 B_0$ (this is entirely possible in accordance with modern ideas about the evolution of neutron stars) both processes should be taken into account in setting up the energy balance of such collapsed objects (see also [2]).

We thank A. Yu. Loskutov for computer analysis of the results.

LITERATURE CITED

1. L. Rozenberg, Phys. Rev. 129, 2786 (1963).
2. Yu. M. Loskutov and V. V. Skobelev, Yad. Fiz., 31, 1279 (1980).
3. Yu. M. Loskutov, Pis'ma Zh. Eksp. Teor. Fiz., 39, 438 (1984); Dokl. Akad. Nauk SSSR, 275, 1396 (1984); Teor. Mat. Fiz., 65, 141 (1985).
4. V. V. Skobelev, Zh. Eksp. Teor. Fiz., 71, 1263 (1976); 72, 1298 (1977); 73, 1301 (1977).
5. Yu. M. Loskutov and V. V. Skobelev, Zh. Eksp. Teor. Fiz., 77, 809 (1979); Teor. Mat. Fiz., 38, 195 (1979); 39, 64 (1979).
6. Yu. M. Loskutov, in: Proc. of the International Seminar on Problems of High Energy Physics and Quantum Field Theory, Vol. 2 [in Russian], Institute of High Energy Physics, Protvino (1983), pp. 213-230.
7. L. L. De Raad, K. A. Milton, and H. D. Hari Dass, Phys. Rev. D, 14, 3326 (1976).

ERRATUM

In Vol. 83, No. 3, 1990, on p. 663, the authors name should read José A. Farrari.