Errata

An Extended IBA Consistent- Q Formalism Applied to Ru and Pd Isotopes

D. Bucurescu, G. Căta, D. Cutoiu, G. Constantinescu, M. Ivaşcu, and N.V. Zamfir
Central Institute of Physics, Bucharest, Romania

Z. Phys. A – Atomic Nuclei 324, 387–392 (1986)

There is a wrong correspondence between the K and K' parameters of the ECQF Hamiltonian (3) and the values given in Fig. 5. In this figure, the values $-\frac{K}{2}$ and $-K'$ should be read instead of K and K' , respectively. Correspondingly, in Fig. 1 the quantity resprsed is $-2\varepsilon/K$ and not ε/K .

D. Bucurescu, G. Căta, D. Cutoiu,
G. Constantinescu, M. Ivaşcu, N.V. Zamfir
Heavy Ion Physics Department
Central Institute of Physics
P.O. Box MG6
Bucharest
Romania

$p\bar{p}$ Annihilation into Two Mesons in the Quark Rearrangement Model and the Quark Annihilation Model

S. Furui*

Division de Physique Théorique, Institut de Physique Nucléaire, Orsay, France

Z. Phys. A – Atomic Nuclei 325, 375–390 (1986)

The value of $C^2 \times 1296$ for the $\rho^+ \rho^-$ production from the $^{33}S_1$ channel in Table 9 is 3549/16 instead of 1521/8. The definition of C^2 in the rearrangement model differs from that of Green, A.M. and Niskanen, J.A.: Nucl. Phys. **A430**, 605 (1984) by a factor $(2J+1) \times 9/2$.

S. Furui
Institut für Theoretische Physik der Universität Tübingen
Auf der Morgenstelle 14
D-7400 Tübingen 1
Federal Republic of Germany