Addendum

Quantum Tunneling at Low Temperatures: Results for Weak Damping

E. Freidkin and P.S. Riseborough

Department of Physics, Polytechnic Institute of New York,
Brooklyn, New York, USA

P. Hänggi

Lehrstuhl für Theoretische Physik, Universität Augsburg,
Federal Republic of Germany

Received March 18, 1986

Z. Phys. B – Condensed Matter 64, 237–246 (1986)

The approximate evaluations of the integrals in (4.8) and (4.12) can be improved. These integrals can be performed exactly, with the results that in Eq. (4.13) the factor 1.16 should be replaced by 0.472 and 4.07 should be replaced by 0.056. In Eq. (4.14) the factor 1.60 should be precisely 0. The contribution of the continuum eigenvalues can be evaluated directly with Eq. (4.12). As a result of this, Eq. (4.21) should be replaced by $15/2 [1 + 3.011 \alpha]$. On combining these contributions, one finds a factor of 2.860 instead of 5.47 in (4.23). Finally, a factor of $\exp [ik\omega_0 \tau/2]$ should be included in (A.5) of Appendix 1.

E. Freidkin, P.S. Riseborough
Department of Physics
Polytechnic Institute of New York
333 Jay Street
Brooklyn, NY 11201
USA

P. Hänggi
Lehrstuhl für Theoretische Physik
Universität Augsburg
Memmingerstrasse 6
D-8900 Augsburg
Federal Republic of Germany