Corrections

Dear Sir

I am indebted to Professor Hanai (Kyoto University) for pointing out, in private correspondence, an editing mistake in my paper

On the electrical properties of heterogeneous mixtures

(Colloid & Polymer Science 263:51–57 (1985)).

In the section on the generalised Looyenga equation (p. 54), the incremental permittivity changes $\varepsilon_1 \rightarrow \varepsilon - \Delta\varepsilon$ and $\varepsilon_2 \rightarrow \varepsilon + \Delta\varepsilon$ should be applied to the generalised Bruggeman equation (Eq. (6)) and Equation (24) should read:

$$\phi' = \frac{1}{2} \left[1 + A_a \frac{\Delta\varepsilon}{\varepsilon} \right].$$

This leads to the mixture permittivity Equation (26) as required

$$\text{i. e. } \varepsilon = [\varepsilon_1^{1-2A_a} + \phi (\varepsilon_2^{1-2A_a} - \varepsilon_1^{1-2A_a})]^{1/1-2A_a}. \quad (1)$$

This equation is also obtained when the Mandel equation (Ref. [18] in my paper), which is a spheroidal version of the Böttcher equation, is used. This approach applied to spheroids parallels that of Looyenga, who used the equations of Bruggeman and Böttcher, both of which are only applicable to spheres.

When the above incremental changes are applied to the Sillars equation, the expression for ϕ' becomes:

$$\phi' = \frac{1}{2} \left[1 + 2A_a \frac{\Delta\varepsilon}{\varepsilon} \right]$$

and, as Professor Hanai has pointed out to me, this leads to a different mixture permittivity equation:

$$\varepsilon = [\varepsilon_1^{1-4A_a} + \phi (\varepsilon_2^{1-4A_a} - \varepsilon_1^{1-4A_a})]^{1/1-4A_a}. \quad (2)$$

Yet another mixture permittivity is obtained when the incremental permittivity changes are applied to the

spheroidal version of the Rayleigh equation, given by van Beek. In this case:

$$\phi' = \frac{1}{2} + \frac{2A_a - 1}{2} \frac{\Delta\varepsilon}{\varepsilon}$$

which leads to

$$\varepsilon = [\varepsilon_1^{3-4A_a} + \phi (\varepsilon_2^{3-4A_a} - \varepsilon_1^{3-4A_a})]^{1/3-4A_a}. \quad (3)$$

The above three expressions generate very different values for ε as illustrated below for spheres ($A_a = \frac{1}{3}$) with $\varepsilon_1 = 78.0$ and $\varepsilon_2 = 2.0$

ϕ	ε Equation 1	Equation 2	Equation 3
0.25	43.59	19.12	65.66
0.50	21.17	7.37	51.57
0.75	8.16	3.58	34.09

In the integral method used to obtain the generalised Bruggeman equation (Eq. (6) in my paper), the same equation is obtained when the Sillars, van Beek or Mandel equation is used.

Dr. M. H. Boyle
Senior Lecturer
Sheffield City Polytechnic
Department of Applied Physics
Pond Street
Sheffield S1 1WB, England