ERRATUM

ADOLESCENT FRIENDSHIP PATTERNS AS PROGNOSTIC INDICATORS FOR SCHIZOPHRENIC ADULTS

BY RAYMOND PITT, M.A., DONALD S. KORNFELD, M.D. AND
LAWRENCE C. KOLB, M.D.

A transposition in column headings in the paper by Raymond Pitt et al. in the July 1963 *PSYCHIATRIC QUARTERLY* presented the conclusions regarding friendship patterns during adolescence, by persons who became schizophrenic adults, in a fashion to reverse the authors' conclusions, which were: "The deterioration in friendship capacity was demonstrated only by ... [a] longitudinal approach. Table 3 shows the changes noted..." As originally printed in the July *QUARTERLY*, Table 3 shows the "Mid-Adolescence" column to the left of the "Early Adolescence" column, with each heading over the other column's figures—thereby reversing the trend noted. The corrected Table 3 follows:

Table 3. Intensity of Interaction in Individual Relationships

Type of Hospitalization	No. of Patients in Early Adolescence Interaction				No. of Patients in Mid-Adolescence Interaction			
	Hi*	Med**	Low†	Total	Hi	Med	Low	Total
Short-term	6	5	0	11	8	2	1	11
Relapsing	3	5	3	11	1	4	6	11
Chronic	5	2	4	11	0	2	9	11
				33				33

*High Interaction=Two or more times a week

**Medium=One time a week

†Low=Less than once a week