Erratum

Squark and gluino decays for large $\tan \beta$

A. Bartl¹, W. Majerotto², W. Porod¹

¹ Institut für Theoretische Physik, Universität Wien, A-1090 Vienna, Austria

² Institut für Hochenergiephysik, Österreichische Akademie der Wissenschaften, A-1050 Vienna, Austria

Z. Phys. C 64 (1994) 499–507

Equation (11) correctly reads

$$\cos \theta_{\tilde{t}} = \frac{-a_{\tilde{t}} m_{\tilde{t}}}{\sqrt{(m_{\tilde{t}_L}^2 - m_{\tilde{t}_1}^2)^2 + a_{\tilde{t}}^2 m_{\tilde{t}}^2}}.$$

In the last expression of Eq. (33), page 507, the left hand side should read f_{2j}^b instead of f_{1j}^b . In Fig. 1b and Fig. 1c we have taken $A_{\tilde{t}} = A_{\tilde{b}} = 330 \text{ GeV}$. In Figs. 2–5 $m_{\tilde{t}} = 150 \text{ GeV}$ was taken.

Zeitschrift für Physik C – Particles and Fields is delivered to some institutions in the former Soviet Union through a program initiated and administered by the European Physical Society and supported by the publisher. This program is funded by INTAS, the international association for the promotion of cooperation with scientists from the independent States of the former Soviet Union. Members of INTAS are the European Union, Austria, Belgium, Denmark, Finland, France, Germany, Greece, Ireland, Italy, Luxembourg, The Netherlands, Portugal, Spain, United Kingdom and Switzerland. (INTAS, rue du Luxembourg 14A, 1040 Bruxelles, Belgique).