Short communications

On partitions of N into summands coprime to N

P. ERDÖS and B. RICHMOND

Let $R(n)$ and $R'(n)$ denote the number of partitions of n into summands and distinct summands respectively that are relatively prime to n . Erdős obtained the first results concerning the asymptotic behaviour of these functions showing that

$$\log R(n) \sim \pi\sqrt{\frac{2}{3}} \phi^{1/2}(n)$$

$$\log R'(n) \sim \pi\sqrt{\frac{1}{3}} \phi^{1/2}(n)$$

where $\phi(n)$ denotes Euler's function. Richmond showed that the error terms in the above result are $O\{\exp((1+\epsilon)\log 2 \log n / \log \log n)\}$ by showing that the asymptotic results of Roth and Szekeres hold for this problem. In this note previous results are improved and a certain set of integers is exhibited for which asymptotic expansions in terms of standard number-theoretic functions may be obtained.

Received July 5, 1976 and, in revised form, November 7, 1976.

About orthogonal permutations of the group $G = G(2)XG(k)$

J.-C. PANAYIOTOPOULOS

Let the group $G = G(2)XG(k)$ of order $n = 2k$, k even, such that:

$$G = \{(0, 0), (0, 1), \dots, (0, k-1), (1, 0), (1, 1), \dots, (1, k-1)\} [1],$$

with

$$a \pm b = (a_1, a_2) \pm (b_1, b_2) = ((a_1 \pm b_1)(\bmod 2), (a_2 \pm b_2)(\bmod k)), a, b \in G$$