Errata

Ambulatory blood pressure in patients with Parkinson's disease without and with orthostatic hypotension

Jean-Michel Senard, Bernard Chamontin, André Rascol and Jean-Louis Montastruc *Clinical Autonomic Research* 1992; 2: 99-104.

In Table 2, the value indicated for heart rate during night for patients with orthostatic hypotension is not 4 beats/min as stated but 69 ± 4 beats/min.

Table 4 should read as follows:

Table 4. Systolic (SBP) and diastolic (DBP) blood pressure variability (%) during the diurnal (from 0900 to 1900) and the nocturnal period (from 2300 to 0600) in Parkinson's disease (PD) patients with or without orthostatic hypotension. The blood pressure variability was assessed by the variation coefficient of the mean blood pressure values obtained during each period of time

	Day		Night	
	SBP	DBP	SBP	DBP
PD without OH	10.3 ± 0.7	11.8 ± 0.7	8.9 ± 0.6	11.6 ± 0.9
PD with OH	$14.6 \pm 1.3^*$	$16.5 \pm 1.0^*$	9.1 ± 0.8	10.8 ± 1.1

* $p < 0.05$ when compared with values obtained in PD without OH. Mean values $\pm$ SEM.

The Publisher apologises for any inconvenience this has caused.