EDITORIAL

*From the Department of Pathology,
McGill University, Montreal (Canada)*

Cardiomyopathies

Kardiomyopathien

G. R o n a

(Received March 13, 1974)

Summary

Cardiomyopathy is defined as heart muscle dysfunction of unknown cause and association. On the basis of functional anatomy and clinical presentation, it is possible to identify three types: congestive, hypertrophic and obliterative cardiomyopathy.

The term "cardiomyopathy" was introduced by *Bridgen* (1) in 1957 for delineating heart muscle disorders of unknown or unusual etiology from the well defined common cardiac conditions associated with rheumatic fever, hypertension, coronary and thyroid disease or congenital development anomaly. Another term considered to be synonymous with cardiomyopathy, primary myocardial disease, was suggested by *Mattingly* (2) for conditions that affect the heart muscle producing faults of myocardial function and spare other anatomic structures within the cardiovascular system. Primary myocardial disease includes diseases of both known and unknown origin, pathogenesis and/or association and could be subdivided into two major categories: inflammatory and non-inflammatory conditions. The former generally is produced by infectious agents and is called myocarditis.

The terminology and classification of cardiomyopathies have undergone major revisions (3). However, it has not been clarified as yet how broadly the term should be used. The proper definition of terms "primary" and "secondary" cardiomyopathies (4, 5) also awaits for general acceptance. Primary cardiomyopathy may be regarded as a disease that affects the myocardium alone (2) or of unknown etiology (6). Consequently, in secondary cardiomyopathy, the heart is affected with other organs in a systemic disease (2) or the cause of myocardial involvement is known albeit not fully understood (6). Clinical analysis, when correlated with pathological findings may indicate the broad etiological group and, in some cases, the myocardial reaction pattern suggests the causative factor.

In addition to the problems of terminology and classification, the difficulty of diagnosis is the main reason why the incidence of cardiomyopathies is still poorly documented. Clinically apparent myocardial disease may not be accompanied by pathognomonic histopathological changes. Clinicians are prone to label too frequently a cardiac condition as hyper-

tensive or to be related to coronary heart disease. This decreases the percentage of myocardial diseases where the etiology and pathogenesis is not clearly established. The data of WHO indicate that there are well established geographical differences in distribution, and that cardiomyopathies, particularly in underdeveloped countries, represent an important health problem. While infectious myocardial disease is becoming less common with control of infection, cardiomyopathies of unknown etiology are being recognized more frequently also in the Western nations (7).

A great effort has been directed for search of the etiological agents, genetic background, immunological, iatrogenic, or nutritional factors responsible as well as for hemodynamic, biochemical and subcellular alterations associated with cardiomyopathies. According to *Goodwin* (8), cardiomyopathy is a subacute or chronic disorder to the heart of unknown or obscure etiology often with associated endocardial and sometimes with pericardial involvement but not atherosclerotic in origin. He formerly differentiated four main forms of primary cardiomyopathies: the congestive, hypertrophic, constrictive (restrictive) and obliterative forms. Congestive cardiomyopathies represent probably a multicausal group in which low cardiac output is associated with marked dilatation of the heart; hypertrophy may develop as a compensatory adjustment, but is overshadowed by the dilatation. Hypertrophic cardiomyopathy has a well documented genetic background. The hypertrophied left ventricle and reduced left ventricular cavity explains the reduced ventricular compliance. Obstruction of left ventricular outflow tract may follow [idiopathic hypertrophic subaortic stenosis (9) or obstructive cardiomyopathy (10)]. In the constrictive form, myocardial rigidity results in the restriction of diastolic ventricular filling. In the obliterative cardiomyopathies, fibrosis leads to obliteration of left ventricular cavity and atrioventricular incompetence. Although the evolution of disease is different, the end results are the same in all forms: congestive heart failure, pulmonary edema, atrial fibrillation and embolism (11). More recently *Goodwin* (12) for the interest of simplification, redefined cardiomyopathies as heart muscle diseases of unknown cause that allow to concentrate efforts on the main aspects of anatomy, function and clinical patterns. Only two forms, the congestive and the hypertrophic cardiomyopathies were retained following this definition.

In 1971, a Committee of the International Study Group for Research in Cardiac Metabolism (13) defined cardiomyopathy as heart muscle dysfunction of unknown cause and association. Three well characterized types were indentified: 1. systolic/pump failure (congestive form, COCM), 2. Diastolic/compliance failure (hypertrophic cardiomyopathy, HOCM) and obliterative cardiomyopathy. This classification was supported recently by the Council on Cardiomyopathies of the International Society of Cardiology (14).

Zusammenfassung

„Kardiomyopathie“ ist definiert als eine Dysfunktion des Herzmuskels unbekannter Ursache. Auf der Grundlage funktioneller Anatomie und des klinischen Erscheinungsbildes ist es möglich, drei Typen zu unterscheiden: kongestive, hypertrophische und obliterierende Kardiomyopathie.

References

1. *Brigden, W.*, Uncommon Myocardial Diseases. The Non-Coronary Cardiomyopathies. *Lancet* **1957/II**, 1179-1184, 1243-1249. – 2. *Mattingly, T. W.*, Clinical Features and Diagnosis of Primary Myocardial Diseases. *Mod. Conc. Cardiovas. Dis.* **30**, 677-682, 683-686 (1961). – 3. *Korb, G.*, Heart Diseases of Unknown Etiology: Problems of Terminology and Classification. In: *Bajusz, E.* and *Rona, G.*, with *A. Brink* and *A. Lochner* (Ed.), *Advances in Cardiac Structures and Metabolism, Cardiomyopathies*, Vol. 2, pp. 9-16 (Baltimore, 1973). – 4. *Fejfar, Z.*, Definition and Classification of the Cardiomyopathies. *Path. Microbiol.* **35**, 17-25 (1970). – 5. *Mattingly, T. W.*, Diseases of the Myocardium (Cardiomyopathies): The Viewpoint of a Clinical Cardiologist. Editorial. *Amer. J. Cardiol.* **25**, 79-80 (1970). – 6. *Hudson, R. E. B.*, The Cardiomyopathies: Order from Chaos. *Amer. J. Cardiol.* **25**, 70-77 (1970). – 7. *Burch, G. E.*, and *Giles, T. D.*, Incidence of Cardiomyopathy. In: *Bajusz, E.* and *Rona, G.* with *A. J. Brink* and *A. Lochner* (Ed.), *Advances in Cardiac Structure and Metabolism, Cardiomyopathies*, Vol. 2, pp. 59-75 (Baltimore, 1973). – 8. *Goodwin, J. F.*, Congestive and Hypertrophic Cardiomyopathies. A decade of Study. *Lancet* **1970/I**, 731-739. – 9. *Loogen, F.*, *Gleichmann, U.* and *Krelhaus, W.*, Die obstruktive Myokardiopathie. *Z. Kreislaufforschg.* **60**, 1044-1066 (1971). – 10. *Ferrans, V. J.*, *Morrow, A. G.* and *Roberts, W. C.*, Myocardial Ultrastructure in Idiopathic Hypertrophic Subaortic Stenosis. A Study of Operatively Excised Left Ventricular Outflow Tract in 14 Patients. *Circulation* **45**, 769-792 (1972). – 11. *Oakley, C.*, Ventricular Hypertrophy in Cardiomyopathy. *Brit. Heart J.* **33**, Suppl. 179-186 (1971). – 12. *Goodwin, J. F.*, Cardiomyopathies in England. In: *Bajusz, E.* and *Rona, G.* with *A. J. Brink* and *A. Lochner* (Ed.), *Advances in Cardiac Structures and Metabolism, Cardiomyopathies*, Vol. 2, pp. 79-93 (Baltimore, 1973). – 13. Committee of the International Study Group for Research in Cardiac Metabolism on Cardiomyopathy. Nairobi, Kenya, October 1971 (to be published in *Myocardiology in Africa*, Edited by *F. Kamunvi*, *H. P. Ojiambo*, *E. Bajusz* and *G. Rona* (Nairobi, Kenya, 1974). – 14. Council on Cardiomyopathies, Internat. Soc. Cardiol. London, Oct. 1973. Proceedings Edited by *Oakley, M. C.* and *Olsen, E. G. J.* To be published in *Ciba Colloquia* 1974.

Author's address:

Prof. Dr. G. Rona, Department of Pathology, McGill University
Montreal 101 (Canada)