References

- [1] O. ZIETSCHMANN, Anat. Anz. 33, 456 (1908).
 [2] M. J. BRODY, R. HÅKANSON, CH. OWMAN and F. SUNDLER, J. Histochem. Cytochem. 20, 945 (1972).
 [3] M. J. BRODY, R. HÅKANSON, I. LUNDQUIST, CH. OWMAN and F. SUNDLER, J. Histochem. Cytochem. 21, 13 (1973).
 [4] H. KETTERER, H.-Y. RUOFF and K.-FR. SEWING, Experientia 29, 1096 (1973).
 [5] L. GRIMELIUS, Acta Soc. Med. upsal. 73, 243 (1968).
 [6] L. HELLERSTRÖM and B. HELLMAN, Acta endocrinol. (Kbh.) 35, 518 (1960).

Errata

Agents und Actions
 vol. 4/2, p. 71 (1974)

The Histaminase Activity of Guinea-Pig's Plasma during Experimental Inflammation

II. Some Properties of the Enzyme

by T. BIEGAŃSKI and CZ. MAŚLIŃSKI

Figure 3: The abscissa should read: $1/S \times 10^{-3} M^{-1}$.

Agents and Action
 vol. 4/2, p. 74-94 (1974)

Chemical Properties of a High Molecular Weight Spleen Extract with Permeability Activity

by H. FF. S. DAVIES, ROSE E. HARTLEY and H.O. SCHILD

The Isolation of High Molecular Weight Proteins from Spleen Tissue, which Alter Vascular Permeability: Investigation of their Pharmacological Properties

by H. FF. S. DAVIES, ROSE E. HARTLEY and H.O. SCHILD

The articles should have been published in the reversed order.