Table 1: Susceptibility of 34 *Eikenella corrodens* strains to 14 antimicrobial agents.

Antimicrobial agent	Cumulative percentage of strains susceptible to increasing concentrations (mg/l)												
	≤ 0.03	0.06	0.13	0.25	0.5	1	2	4	8	16	32	64	128
Mezlocillin	18	26	44	76	94	100							
Azlocillin	3	18	21	47	76	97	100						
Piperacillin	12	24	41	53	97	100							
Ticarcillin			3	6	56	91	100						
Ceftizoxime	68	94	100										
Cefotaxime	50	88	97		100								
Cefmenoxime	38	76	97	100									
Lamoxactam	21	71	91	97	100								
Cefoperazone	9	15		18	41	74	91	100					
Cefotiam				3	26	85	94	97	100				
Cefazedone						6	42	88	97	100			
Cefuroxime					3	12	26	91	94	100			
Cefsulodin											32	88	100
Fosfomycin									3	15	82	97	100

References

1. Goldstein, E. J. C., Cherubin, C. E., Shulman, M.: Comparison of microtiter broth dilution and agar dilution methods for susceptibility testing of *Eikenella corrodens*. *Antimicrobial Agents and Chemotherapy* 1983, 23: 42–45.
2. Ericsson, H. M., Sherris, J. C.: Antibiotic sensitivity testing. Report of an international collaborative study. *Acta Pathologica Microbiologica Scandinavica (B)* 1971, Supplement 217: 1–90.
3. Buttiaux, R., Beerens, H., Tacquet, A.: *Manuel de techniques bactériologiques*. Flammarion, Paris, 1969, p. 514.

Errata

B. K. Pugashetti et al.: Flagellation is a Strain-Dependent Characteristic of Cystic Fibrosis-Associated Mucoid Strains of *Pseudomonas aeruginosa* (Vol. 2, No. 4, p. 360–361).

Page 360, second to last line should read "irreversible loss of mucoidy would yield the ...".

E. Martin: Once-Daily Administration of Ceftriaxone in the Treatment of Meningitis and Other Serious Infections in Children (Vol. 2, No. 5, p. 509–515).

Page 509, 8th line of the abstract should read "Three patients died ...".