Minimum Dissection of a Rectilinear Polygon with Arbitrary Holes into Rectangles

Valeriu Soltan and Alexei Gorpinevich

Institute of Mathematics, Moldavian Academy of Sciences, Grosul str 5,
Kishinev 277028, Republica Moldova

Abstract. In this paper, the problem of dissecting a plane rectilinear polygon with arbitrary (possibly, degenerate) holes into a minimum number of rectangles is shown to be solvable in $O(n^{3/2} \log n)$ time. This fact disproves a famous assertion about the NP-hardness of the minimum rectangular dissection problem for rectilinear polygons with point holes.

1. Introduction

The following decomposition problem is well studied in computational geometry (see, for instance, [1], [3], [4], and [6]–[9]): Given a rectilinear polygon with rectilinear holes, dissect its interior into a minimum number of nonoverlapping rectangles. In the case of nondegenerate holes this problem can be solved in $O(n^{5/2})$ time, where n is the number of vertices of the polygon (see [1], [4], and [6]). Lipski [7], [8] and Imai and Asano [3] gave $O(n^{3/2} \log n)$ algorithms for this minimum dissection problem. Both algorithms make use of the bipartite intersection graph of a set of vertical and horizontal line segments.

If the holes of a rectilinear polygon can degenerate into points, the minimum dissection problem is asserted to be NP-hard [5]. Nevertheless, we prove below the solvability of this problem in $O(n^{3/2} \log n)$ time in the case of arbitrary (possibly, degenerate) holes. In this way, we adapt the respective techniques developed in [3], [4], and [6]–[8], which breaks down in the case of degenerate holes.

2. Description of Rectilinear Polygons

Let P be a (possibly, multiply connected) bounded polygonal region in the coordinate plane E . The topological boundary $\text{bd } P$ of P is assumed to be the

union of a finite number of simple rectilinear contours. A polygonal contour is called *rectilinear* provided all of its line segments are either horizontal or vertical. We assume that any two of these contours will be situated either one inside the other or mutually noninclusive, that they will have common vertices but no common line segments, and that they are not interlaced.

As usual, a point $x \in E$ is called interior for P provided it is contained inside an odd number of simple contours determining $\text{bd } P$. All interior points of P form the topological interior $\text{int } P$.

Inside P a finite family of *closed* line segments $s_1, \dots, s_u$ (either horizontal or vertical), and a finite family of isolated points $v_1, \dots, v_t$, may be situated such that the following conditions are fulfilled:

- (1) $v_1, \dots, v_t$ belong to $\text{int } P \setminus (s_1 \cup \dots \cup s_u)$;
- (2) the interior of each segment s_i is contained in $\text{int } P$;
- (3) if segments s_i, s_j have a common point, then it is a vertex for both s_i, s_j .

The point-set union of these points and segments is called the *ornament* of P , and denoted by $\text{Or } P$.

In order to consider the degenerate holes, we introduce a nonstandard topology in the plane E . Since the description of the respective constructions is sometimes rather formal, we use some illustrative examples. The first of them is shown in Fig. 1.

Example. For rectilinear polygon B , represented in Fig. 1, $\text{bd } B$ is the union of four rectilinear contours $[1, 3, 10, 11, 24, 23, 32, 30, 1]$, $[4, 5, 13, 12, 4]$, $[9, 10, 23, 22, 9]$, and $[16, 17, 26, 25, 16]$. We have

$$\begin{aligned} \text{Or } B = & \{21\} \cup \{29\} \cup [2, 7] \cup [6, 7] \cup [7, 8] \cup [7, 14] \\ & \cup [15, 20] \cup [18, 19] \cup [18, 27] \cup [19, 28] \cup [25, 31] \cup [27, 28]. \end{aligned}$$

Since the segments of contour $[18, 19, 28, 27, 18]$ are defined to be in $\text{Or } P$, the interior of rectangle $[18, 19, 28, 27]$ belongs to $\text{int } B$.

Fig. 1. A rectilinear polygon in the plane.

Definition 1. The sets

$$\text{Bd } P := \text{Or } P \cup \text{bd } P, \quad \text{Int } P := P \setminus \text{Bd } P$$

are called, respectively, the *formal boundary* and the *formal interior* of P . Any bounded connected component of $E \setminus \text{Int } P$ is named a *formal hole* of P , and the unbounded connected component of $E \setminus \text{Int } P$ is called the *formal exterior* of P . We write $\text{Ext } P$ for the formal exterior of P .

From the definitions above, it follows that $\text{Bd } P$, $\text{Ext } P$, and any formal hole of P are *closed* sets, while $\text{Int } P$ is *open*. As usual, a *topological hole* of P is defined to be a bounded connected component of $E \setminus P$. Observe that topological holes are independent of the choice of $\text{Or } P$.

Example. Polygon B in Fig. 1 has five formal holes: points 21 and 29, segment $[15, 20]$, rectilinear contour $[18, 19, 28, 27, 18]$, and rectangle $[4, 5, 13, 12]$. Rectangle $[16, 17, 26, 25]$ is not a formal hole of B , because it is connected with the exterior of B by segment $[25, 31]$. B has three topological holes, namely, the interiors of rectangles $[4, 5, 13, 12]$, $[9, 10, 23, 22]$, and $[16, 17, 26, 25]$.

Now we are able to define the dissection of P into rectangles.

Definition 2. A rectilinear polygon P is said to be *dissected into rectangles* $R_1, \dots, R_q$ if and only if

$$\bigcup_{i=1}^q \text{int } R_i \subset \text{Int } P \subset \bigcup_{i=1}^q R_i, \quad \text{int } R_i \cap \text{int } R_j = \emptyset \quad \text{for } i \neq j.$$

In other words, nonoverlapping rectangles $R_1, \dots, R_q$ dissect P if their union coincides with P and the formal boundary $\text{Bd } P$ belongs to the union of $R_1, \dots, R_q$ boundaries.

Example. A dissection into rectangles of rectilinear polygon B from Fig. 1 is shown in Fig. 2.

We continue the description of rectilinear polygons.

Definition 3. A point w is called a *vertex* of P in any one of the following cases:

- (1) w is a vertex of a contour forming $\text{bd } P$;
- (2) w is a vertex of a segment s_i forming $\text{Or } P$;
- (3) w is one of the isolated points v_i in $\text{Or } P$.

The set of all vertices of P is denoted by $V(P)$.

Fig. 2. A dissection of a rectilinear polygon into rectangles.

Definition 4. A closed line segment $[x, z]$, $x \neq z$, is called an *elementary segment* of $\text{Bd } P$ if $[x, z] \subset \text{Bd } P$ and $[x, z] \cap V(P) = \{x, z\}$. The set of all elementary segments of $\text{Bd } P$ is denoted by $S(P)$.

Example. Polygon B in Fig. 1 has 32 vertices and 32 elementary segments. Note that $[1, 3]$ and $[30, 32]$ are not elementary segments of $\text{Bd } B$, because they contain, respectively, vertices 2 and 31 in their interiors.

From the definitions, it follows that a common point of elementary segments $t_i, t_j \in S(P)$ is a vertex for both segments.

Observation 1. A rectilinear polygon P is considered to be determined if the sets $\text{int } P$, $\text{Bd } P$, and $V(P)$ are known. It is easily seen that $\text{Bd } P$ and $V(P)$ uniquely determine $S(P)$.

3. Measure of Local Nonconvexity

Any vertex v (if it is not an isolated point in $\text{Bd } P$) is the apex of at least one nonempty *inner angle* of P . Every inner angle of P with the apex v is bounded by two elementary segments of the form $[u, v]$, $[v, w]$, and contains no other inner angle with the apex v . We allow the sides of an inner angle to coincide (in this case, the size of the angle equals 2π). Obviously, any inner angle of a rectilinear polygon is of size $\pi k/2$, $k = 1, 2, 3, 4$. Inner angles of size $3\pi/2$ or 2π are called *concave*.

Definition 5. A vertex v of P is called a *point of local nonconvexity* of P provided it is either an isolated point in $\text{Bd } P$ or the apex of a concave inner angle of P ; otherwise, v is called a point of local convexity of P .

Example. In Fig. 1, vertex 7 is the apex of four inner angles; each of the vertices 2, 10, 18, 19, 23, 27, 28, and 31 is the apex of two inner angles; each of the vertices

6, 8, 14, 15, and 20 is the apex of an inner angle of size 2π . Vertices 4–6, 8, 9, 12–22, and 26–29 are points of local nonconvexity for B .

Definition 6. The *measure* $m(v)$ (of local nonconvexity) of a rectilinear polygon P at a vertex is defined as follows:

- $m(v) = 0$ if v is a point of local convexity of P ;
- $m(v) = 1$ if v is a point of local nonconvexity but not an isolated point in $\text{Bd } P$;
- $m(v) = 2$ if v is an isolated point in $\text{Bd } P$.

Let

$$m(P) := \sum m(v),$$

where the sum is taken over the set $V(P)$ of all vertices of P . The number $m(P)$ is called the *measure* (of local nonconvexity) of P .

Example. In Fig. 1, the measure of B at each of the vertices 21, and 29 equals two, at each of the vertices 4–6, 8, 9, 12–20, 22, and 26–28 is equal to one, and at any other vertex it equals zero. Thus $m(B) = 22$.

We now state two simple lemmas that will be of use later.

Lemma 1. Any formal hole of a rectilinear polygon P contains at least one vertex of local nonconvexity of P .

Proof. Let H be any formal hole of P . It is easily seen that any vertex v of H having the maximum sum $X + Y$ of its Cartesian coordinates (X, Y) is a vertex of the local nonconvexity of P . $\square$

Lemma 2. A rectilinear polygon P has no point of local nonconvexity if and only if each component of $\text{Int } P$ is an open rectangle; i.e., if $\text{Bd } P$ divides P into rectangles.

Proof. A rectilinear polygon P has no point of local nonconvexity if and only if $\text{Bd } P$ contains no isolated point and each inner angle of P is of size π or $\pi/2$. Now, the assertion of Lemma 2 readily follows. $\square$

4. Effective Chords

Further we shall see that the following line segments are of significant importance in any dissection of a rectilinear polygon into a minimum number of rectangles.

Definition 7. A line segment $[v, w]$ is called an *effective chord* of P if the following conditions are fulfilled:

- (1) $[v, w]$ is either horizontal or vertical;
- (2) the open interval $]v, w[$ with the exception of a finite (possibly, empty) set of points is contained in $\text{Int } P$;
- (3) v, w are vertices of local nonconvexity of P and each of them is either an isolated point in $\text{Bd } P$ or a vertex of an elementary segment of $\text{Bd } P$ collinear to $[v, w]$;
- (4) if a point $x \in \text{Bd } P$ belongs to $]v, w[$, then x is a vertex of a unique elementary segment of $\text{Bd } P$ orthogonal to $[v, w]$.

Example. Polygon C (see Fig. 3) with the ornament

$$\text{Or } C = [1, 2] \cup [3, 4] \cup [7, 8] \cup [12, 13] \cup \{5\} \cup \{10\} \cup \{11\}$$

has the following effective chords: $[2, 12]$, $[4, 5]$, $[5, 10]$, $[6, 7]$, $[9, 10]$, and $[10, 11]$.

From Definition 7, it follows:

Observation 2. Any two different effective chords of P have at most one common point. Any proper part of an effective chord cannot be an effective chord.

Definition 8. A family of effective chords of a rectilinear polygon P is called *admissible* if any two mutually orthogonal chords of this family have no common point. The maximum number of effective chords forming an admissible family for P is named the *effective number* of P and is denoted by $e(P)$.

Example. Polygon C in Fig. 3 has three maximal admissible families of effective chords: $\{[2, 12], [5, 10]\}$, $\{[2, 12], [4, 5], [9, 10], [10, 11]\}$, and $\{[4, 5], [6, 7]\}$.

Fig. 3. Effective chords of a rectilinear polygon.

$[9, 10], [10, 11]\}$. Thus $e(C) = 4$, and a maximal admissible family need not be a maximum admissible family.

Observation 3. If two effective chords of an admissible family for P have a common vertex, then these two chords are collinear and their common vertex is an isolated point in $\text{Bd } P$.

5. Addition of Segments to Formal Boundary

Any dissection of a rectilinear polygon into rectangles can be considered as a repeated addition of some *closed* line segments to the formal boundary of the polygon. In this section, we study some auxiliary results regarding this addition.

Let $[x, z] \subset P$ be a *closed* line segment (either horizontal or vertical) such that the intersection of $[x, z]$ with $\text{Bd } P$ is a finite (possibly, empty) set. We can say that the polygon P' is obtained from P by the addition of $[x, z]$ to $\text{Bd } P$ provided:

- (1) $\text{Bd } P' := \text{Bd } P \cup [x, z]$;
- (2) $\text{Int } P' := \text{Int } P \setminus [x, z]$;
- (3) $V(P') := V(P) \cup \{x, z\} \cup (\text{Or } P \cap [x, z])$.

Observation 4. Segment $[x, z]$, added to $\text{Bd } P$, is not, in general, an elementary segment of $\text{Bd } P'$; it is, however, a union of elementary segments of $\text{Bd } P'$.

We say that the addition of a segment $[x, z] \subset P$ to $\text{Bd } P$ *decreases by one* the measure $m(z)$ of P at a vertex $z \in V(P)$ if $m'(z) = m(z) - 1$, where $m'(z)$ means the measure of P' at z .

Example. In Fig. 3, the addition of segment $[2, 12]$ to $\text{Bd } C$ decreases by one the measure of C at each of the vertices 2, 7, and 12, while the addition of segment $[11, 12]$ to $\text{Bd } C$ decreases the measure by one at only vertex 11.

Lemma 3. Let P' be a polygon obtained from P by the addition to $\text{Bd } P$ of a segment $[x, w] \subset P$ such that $w \in V(P)$. Then

$$m(w) - 1 \leq m'(w) \leq m(w),$$

where $m'(w)$ denotes the measure of P' at w .

Proof. If w is a point of local convexity of P , then trivially $m(w) = m'(w) = 0$. Assume that w is a point of local nonconvexity of P but not an isolated point in $\text{Bd } P$. Then w is the apex of a concave inner angle of P . If $[X, w]$ does not intersect the interior of this angle, then $m(w) = m'(w) = 1$. Let $[x, w]$ divide this angle in two inner angles of P' . If one of these new angles is concave, then $m(w) = m'(w) = 1$. If both new angles are convex, then $m(w) - 1 = m'(w) = 0$.

Assume now that w is an isolated point in $\text{Bd } P$. Then w is a vertex of a unique

elementary segment $[u, w] \in S(P')$ (obviously, $[u, w]$ lies in $[x, w]$). In this situation $m(w) - 1 = m'(w) = 1$. $\square$

Lemma 4. *Let $[v, w]$ be an effective chord of a rectilinear polygon P , and let $x_1, \dots, x_r$ be all vertices of P contained in $]v, w[$. The addition of $[v, w]$ to $\text{Bd } P$ decreases by one the measure of P at each of the points $v, w, x_1, \dots, x_r$.*

Proof. Each x_i is a vertex of a unique elementary segment, say $[x_i, z_i]$, orthogonal to $[v, w]$ (see item 4 of Definition 7). In this case, $m(x_i) = 1$ for all $i = 1, \dots, r$.

Denote by P' the polygon obtained from P by the addition of $[v, w]$ to $\text{Bd } P$. Relative to P' , each vertex x_i is the apex of three convex inner angles (two of them are of size $\pi/2$ and one is of size π). Hence x_i is a point of local convexity of P' and $m'(x_i) = 0$, where $m'(x_i)$ denotes the measure of P' at x_i . Hence $m'(x_i) = m(x_i) - 1$.

If v is an isolated point in $\text{Bd } P$, then v is a vertex of a unique elementary segment of the form $[v, x_i] \in S(P')$. In this situation, $m'(v) = m(v) - 1 = 1$. Assume that v is not isolated in $\text{Bd } P$. Then v is a vertex of an elementary segment, say $[u, v] \in S(P)$, which is collinear to $[v, w]$ (see item 2 of Definition 7). In this situation, v is a point of local convexity for P' and $m'(v) = m(v) - 1 = 0$. Similarly, $m'(w) = m(w) - 1$. $\square$

We need the following:

Definition 9. A line segment $[x, z]$ is called a *simple chord* of P if the following conditions are fulfilled:

- (1) $[x, z]$ is either horizontal or vertical;
- (2) $x, z \in \text{Bd } P$;
- (3) the open interval $]x, z[$ is contained in $\text{Int } P$.

Observation 5. From Definitions 4 and 9 it follows that if a simple chord $[x, z]$ of P is added to $\text{Bd } P$, then $[x, z]$ becomes an elementary segment for the polygon P' such that

$$\text{Bd } P' = \text{Bd } P \cup [x, z], \quad \text{Int } P' = \text{Int } P \setminus]x, z[, \quad V(P') = V(P) \cup \{x, z\}.$$

Lemma 5. *For any vertex v of local nonconvexity of a rectilinear polygon P there is a simple chord $[v, z]$ of P whose addition to $\text{Bd } P$ decreases by one the measure $m(v)$.*

Proof. The vertex v is either an isolated point in $\text{Bd } P$ or the apex of some inner concave angle of P . If v is an isolated point in $\text{Bd } P$, then we choose a horizontal segment $[v, z]$ such that $z \in \text{Bd } P$ and $]v, z[\subset \text{Int } P$. The chord $[v, z]$ is simple for P and $m'(v) = 1$, where $m'(v)$ denotes the measure at v for the polygon P' obtained from P by the addition of $[v, z]$ to $\text{Bd } P$. In this case, $m'(v) = m(v) - 1 = 1$.

Assume now that v is the apex of an inner concave angle of P formed by

elementary segments, say $[u, v]$, $[v, w]$. Choose the chord $[z, v]$ such that:

- (1) $[z, v]$ is collinear to $[v, w]$, and $v \in]z, x[$;
- (2) $z \in \text{Bd } P$ and $]z, v[\subset \text{Int } P$.

The chord $[z, v]$ is simple for P and $m'(v) = m(v) - 1 = 0$. □

6. Decomposition of Supplementary Ornament

Assume that a rectilinear polygon P is dissected into rectangles $R_1, \dots, R_q$ (see Definition 2). Denote by $L_1, \dots, L_q$ the boundary contours of $R_1, \dots, R_q$, respectively, and put $L := L_1 \cup \dots \cup L_q$.

Define W to be set of vertices of $R_1, \dots, R_q$. Then $V(P) \cup W$ divides the point-set union L of all $R_1, \dots, R_q$ sides into a family $\mathcal{S}$ of *closed* line segments. Therefore, we may consider L as the formal boundary of the rectilinear polygon P_0 such that

$$\text{Bd } P_0 = L, \quad \text{Int } P_0 = \text{Int } P \setminus L, \quad V(P_0) = V(P) \cup W, \quad S(P_0) = \mathcal{S}.$$

Note that P and P_0 have the same topological boundary and topological interior, and any connected component of $\text{Int } P_0$ is the interior of some rectangle R_i .

As mentioned in Section 5, polygon P_0 can be obtained from P by the repeated addition of some *closed* line segments to $\text{Bd } P$. In this section we investigate the inverse approach: to obtain P from P_0 by the repeated deletion of some *open* line intervals. In this way, the “supplementary ornament” $Y := L \setminus \text{Bd } P$ will be studied.

Lemma 6. *The set $Y = L \setminus \text{Bd } P$ can be decomposed into pairwise disjoint open line intervals.*

Proof. Denote by $\mathcal{L}$ the family of all maximal, with respect to inclusion, *open* intervals contained in Y . Since Y is itself an open set, we have $Y = \bigcup \{I : I \in \mathcal{L}\}$. Generally speaking, $\mathcal{L}$ does not satisfy the condition of Lemma 6 because it may contain some elements with nonempty intersection. In order to obtain the desired decomposition, we execute repeatedly the following:

Segment splitting procedure. If open intervals $]x, y[$, $]z, v[$ in $\mathcal{L}$ have a common point w (in this case, $]x, y[$ and $]z, v[$ are orthogonal to each other), then put $\mathcal{L} := (\mathcal{L} \setminus]z, v[\cup \{]z, w[,]w, v[\})$. Since $\mathcal{L}$ is finite, the required decomposition exists. □

Example. Polygon D (see Fig. 4) with the ornament $\text{Or } D = [1, 4] \cup [2, 3] \cup [5, 8] \cup \{6\}$ is dissected into four rectangles. The set $Y := L \setminus \text{Bd } D$ is decomposed into open intervals as follows: $Y =]13, 4[\cup]4, 5[\cup]5, 6[\cup]6, 7[$.

Let $\mathcal{L} = \{I_1, \dots, I_k\}$ be a decomposition of Y into pairwise disjoint open line intervals, and let $I_i =]x_i, z_i[$, $i = 1, \dots, k$. Denote by P_i the rectilinear polygon obtained from P_0 by deleting $I_1 \cup \dots \cup I_i$ from L , $i = 1, \dots, k$.

Generally speaking, the polygon P_k can be different from P because it may contain some new vertices (as vertex 7 of D_0 for polygon D in Fig. 4). At the same

Fig. 4. "Supplementary ornament" of a rectangular dissection.

time $\text{Bd } P_k = \text{Bd } P$, and any vertex $w \in V(P_k) \setminus V(P)$ is a point of local convexity for P . Therefore $m(P_k) = m(P)$.

Since the set $L_i := L \setminus (l_1 \cup \dots \cup l_i)$ may be nonclosed for $1 \leq i < k$, the respective ornaments of $P_1, \dots, P_{k-1}$ are not well defined (by definition, the ornament of a rectilinear polygon is a union of isolated points and closed segments). Therefore we cannot directly apply to P_i the notion of measure of local nonconvexity. Nevertheless, we can avoid this obstacle for vertices of P (but not for vertices of P_i) in the following way.

A set L_i can be nonclosed if we delete, together with some interval l_p , $p \leq i$, an endpoint of an interval l_q , $q > i$. Since $l_q \cap \text{Bd } P = \emptyset$, this deleted vertex cannot be a vertex of $\text{Bd } P$. Hence for any vertex w in $V(P)$, we can choose a small square Q (having vertical and horizontal sides) with center w such that the intersection of Q with L_i is closed. Now we define the measure $m_i(w)$ of P_i at w relative to that part of $\text{Bd } P_i$ which belongs to Q . Since the notion of measure is local, our approach is correct.

Definition 10. For any decomposition of the set $Y = L \setminus \text{Bd } P$ into pairwise disjoint open intervals $l_i =]x_i, z_i[$, $i = 1, \dots, k$, define the numbers $\mu_i(l_i)$, $i = 1, \dots, k$, as follows:

- (1) $\mu_i(l_i) = 0$ if $x_i \notin V(P)$ and $z_i \notin V(P)$;
- (2) $\mu_i(l_i) = m_i(x_i) - m_{i-1}(x_i)$ if $x_i \in V(P)$ and $z_i \notin V(P)$;
- (3) $\mu_i(l_i) = m_i(z_i) - m_{i-1}(z_i)$ if $x_i \notin V(P)$ and $z_i \in V(P)$;
- (4) $\mu_i(l_i) = [m_i(x_i) - m_{i-1}(x_i)] + [m_i(z_i) - m_{i-1}(z_i)]$ if $x_i \in V(P)$ and $z_i \in V(P)$;

where $m_j(w)$ denotes the measure of P_j at a vertex $w \in V(P)$.

From Lemma 3, it follows that

$$0 \leq m_i(x_i) - m_{i-1}(x_i) \leq 1, \quad 0 \leq m_i(z_i) - m_{i-1}(z_i) \leq 1.$$

Hence, from Definition 10, we obtain

Observation 6.

- (1) $0 \leq \mu_i(l_i) \leq 2$ for all $i = 1, \dots, k$;
- (2) $\mu_i(l_i) \leq 1$ if at most one of x_i, z_i is in $V(P)$;
- (3) $\mu_i(l_i) = 0$ if both vertices x_i, z_i are not in $V(P)$.

Observation 7. The values $\mu_i(l_i)$, $i = 1, \dots, k$, depend on the order in which the intervals l_i are considered (see example below).

The remainder of this section is devoted to the proof of:

Theorem 1. Let $\mathcal{L} = \{l_1, \dots, l_k\}$ be a decomposition of the set $Y = L \setminus \text{Bd } P$ into pairwise disjoint open line intervals. The intervals $l_1, \dots, l_k$ can be renumbered so that $0 \leq \mu_i(l_i) \leq 1$ for at least $k - e$ of them, where e denotes the effective number of P .

Before the proof of Theorem 1, we illustrate its statement with the help of the following example.

Example. Polygon D in Fig. 4 has only one effective chord, namely, $[3, 6]$. Hence $e(D) = 1$. The set $Y = L \setminus \text{Bd } D$ is decomposed into open line intervals $]3, 4[$, $]4, 5[$, $]5, 6[$, and $]6, 7[$. If we put

$$l_1 =]3, 4[, \quad l_2 =]5, 6[, \quad l_3 =]4, 5[, \quad l_4 =]6, 7[,$$

then $\mu_1(l_1) = \mu_2(l_2) = 2$, $\mu_3(l_3) = 0$, and $\mu_4(l_4) = 1$. If we renumber these intervals as

$$l_1 =]3, 4[, \quad l_2 =]4, 5[, \quad l_3 =]5, 6[, \quad l_4 =]6, 7[,$$

then $\mu_1(l_1) = 2$ and $\mu_2(l_2) = \mu_3(l_3) = \mu_4(l_4) = 1$, as required in the statement of Theorem 1.

Proof of Theorem 1. An effective chord $[v, w]$ of P is said to be *compatible* with $\mathcal{L}$ provided $[v, w] \setminus V(P)$ is a union of some intervals from $\mathcal{L}$.

Denote by $\mathcal{F} = \{t_1, \dots, t_f\}$ an admissible family of effective chords of P compatible with $\mathcal{L}$ and having maximum cardinality, and put $T := t_1 \cup \dots \cup t_f$. Obviously, $f \leq e$.

Let P_T be the region obtained from P by the addition of T to $\text{Bd } P$; i.e., $\text{Bd } P_T = \text{Bd } P \cup T$, $\text{Int } P_T = \text{Int } P \setminus T$, and $V(P_T) = V(P)$.

Recall that P_i denotes the rectilinear polygon obtained from P_0 by deleting $l_1 \cup \dots \cup l_i$ from L , $i = 1, \dots, k$.

First renumbering. Without loss of generality, we may suppose that the family $\{l_1, \dots, l_k\}$ is renumbered such that, exactly the intervals $l_1, \dots, l_p$, $p \leq k$ have no common vertex with $V(P)$, and all the other intervals $l_{p+1}, \dots, l_k$ are as follows:

- (1) each of the intervals $l_{p+1}, \dots, l_q$, $q \leq k$, has exactly one vertex in $V(P)$;
- (2) each of the intervals $l_{q+1}, \dots, l_s$, $s \leq k$, is not contained in T and has both vertices in $V(P)$;
- (3) each of the intervals $l_{s+1}, \dots, l_k$ is contained in T ; i.e.,

$$l_{s+1} \cup \dots \cup l_k = T \setminus V(P).$$

Obviously, $\text{Bd } P \subset \text{Bd } P_T \subset \text{Bd } P_i$ for all $i = 0, 1, \dots, s$.

Due to Observation 6,

$$\begin{aligned} \mu_i(l_i) &= 0 & \text{for all } i = 1, \dots, p, \\ 0 \leq \mu_i(l_i) &\leq 1 & \text{for all } i = p+1, \dots, q. \end{aligned}$$

Second renumbering is organized for intervals $l_{q+1}, \dots, l_s$ according to the following procedure:

1. Put $i = q+1$ and $k = 0$.
2. If $\mu_i(l_i) \leq 1$, then put $k = 0$ and go to 3, else go to 4.
3. If $i < s$, then put $i = i+1$ and go to 2, else exit.
4. If $k < s-i-1$, then put $k = k+1$, renumber $l_{i+1}, l_{i+2}, \dots, l_s, l_i$ by the indices $i, i+1, \dots, s-1, s$, respectively, and go to 2, else exit.

As a result we obtain a renumbered sequence $l_{q+1}, \dots, l_s$ such that:

- (1) $0 \leq \mu_i(l_i) \leq 1$, $i = q+1, \dots, r$, for some natural r between $q+1$ and s ;
- (2) $\mu_{r+1}(l_{r+1}) = 2$ for any renumbering of $l_{r+1}, \dots, l_s$ by the indices $r+1, \dots, s$.

We are going to prove that the family $l_{r+1}, \dots, l_s$ is empty; i.e., the following assertion is valid.

Lemma 7. $r = s$.

Proof. Suppose, on the contrary, that $r < s$. Choose in the family $l_{r+1}, \dots, l_s$ a maximal subfamily $\mathcal{M}$ such that all the intervals of $\mathcal{M}$ lie on a common line without segment gaps and no common vertex of two intervals from $\mathcal{M}$ is an isolated vertex in $\text{Bd } P_T$. Without loss of generality, consider this line to be horizontal. After an appropriate renumbering of $l_{r+1}, \dots, l_s$, we can represent $\mathcal{M}$ as the union of intervals

$$l_{r+1} =]x_{r+1}, z_{r+1}[, \dots, l_j =]x_j, z_j[, \quad r+1 \leq j \leq s,$$

such that $z_i = x_{i+1}$, $i = r+1, \dots, j-1$ (in the case $r+1 < j$).

We investigate some properties of the segment $[x_{r+1}, z_j]$, which are listed in Claims 1–5 below.

Claim 1. No vertical elementary segment of $\text{Bd } P_r$ is incident with a point of $l_{r+1} \cup \dots \cup l_s$.

Indeed, assume that some vertical elementary segment $m = [v, w]$ of $\text{Bd } P_0$ has its vertex v in $l_{r+1} \cup \dots \cup l_s$. Since $Y \cap \text{Bd } P = \emptyset$, v is not a vertex of P . Hence m is not in $\text{Bd } P$; i.e., m is contained in some interval $l_i \in \mathcal{L}$. Due to First Renumbering, we have $i \leq q$. Hence $m \cap \text{Bd } P_r = \emptyset$.

Claim 2. Each of the vertices $x_{r+1}, z_{r+1}, \dots, z_j$ is incident with at most one vertical elementary segment of $\text{Bd } P_r$.

Suppose, on the contrary, that x_{r+1} is incident with two vertical elementary segments of $\text{Bd } P_r$ (points $z_{r+1}, \dots, z_j$ are handled analogously). In this case, any interior angle of P_{r+1} with the apex x_{r+1} is of size π or $\pi/2$; i.e., x_{r+1} is a point of local convexity for P_{r+1} . Hence $m_{r+1}(x_{r+1}) = 0$. Similarly, $m_r(x_{r+1}) = 0$. By Definition 10, $\mu_{r+1}(l_{r+1}) \leq 1$, which is in contradiction to the condition $\mu_{r+1}(l_{r+1}) = 2$, by the choice of l_{r+1} .

Claim 3 Each of the vertices $z_{r+1}, \dots, z_{j-1}$ (in the case $r + 1 < j$) is incident with a unique vertical elementary segment of $\text{Bd } P_T$.

By the choice of $\mathcal{M}$, each of $z_{r+1}, \dots, z_{j-1}$ is not isolated in $\text{Bd } P_T$. Hence, for each $z_k, k = r + 1, \dots, j - 1$, there exists at least one vertical elementary segment of $\text{Bd } P_T$ whose end is z_k . Due to Claim 2, there exists at most one elementary segment of $\text{Bd } P_T$ whose end is z_k . Since $\text{Bd } P_T \subset \text{Bd } P_r$, there exists exactly one elementary segment of $\text{Bd } P_r$ whose end is z_k and this segment belongs to $\text{Bd } P_T$.

Claim 4. Each of x_{r+1}, z_j is either an isolated point in $\text{Bd } P_T$ or a vertex of a horizontal elementary segment of $\text{Bd } P_T$.

In order to obtain a contradiction, suppose that x_{r+1} is neither an isolated vertex in $\text{Bd } P_T$ nor a vertex of a horizontal segment of $\text{Bd } P_T$ (for z_j , the consideration is similar). Then x_{r+1} is a vertex of a vertical elementary segment of $\text{Bd } P_T$. Due to Claim 2, there is exactly one vertical elementary segment of $\text{Bd } P_T$ incident with x_{r+1} .

If $\text{Bd } P_{r+1}$ were to contain no horizontal elementary segment incident with x_{r+1} , then the deletion of l_{r+1} from $\text{Bd } P_r$ would keep the measure of P_r at x_{r+1} : $m_r(x_{r+1}) = m_{r+1}(x_{r+1}) = 1$. In this case, $\mu_{r+1}(l_{r+1}) \leq 1$, which is impossible by the choice of l_{r+1} . Hence $\text{Bd } P_{r+1}$ contains a horizontal elementary segment, say $[v, x_{r+1}]$.

By this assumption, $[v, x_{r+1}]$ is not in $\text{Bd } P_T$. Hence $[v, x_{r+1}]$ is contained in a horizontal segment $l_i, i \geq r + 2$. Due to the choice of $\mathcal{M}$, we have $i \geq s + 1$. Therefore, by First Renumbering, l_i belongs to a horizontal effective chord $t_h \in \mathcal{F}$. Since l_{r+1} is not contained in t_h , x_{r+1} is a vertex of t_h . In this case, x_{r+1} is either an isolated point of $\text{Bd } P$ or a vertex of a horizontal elementary segment of $\text{Bd } P$ (see Definition 7). Since there is no horizontal elementary segment of $\text{Bd } P$ incident with x_{r+1} , it is an isolated point in $\text{Bd } P$. At the same time, we have shown above that x_{r+1} is a vertex of a unique elementary segment of $\text{Bd } P_T$. The last is possible only if this vertical segment belongs to T ; i.e., x_{r+1} is a vertex of some vertical effective chord $t_v \in \mathcal{F}$.

Thus we have found in $\mathcal{F}$ two orthogonal effective chords t_h, t_v with a common vertex, which is impossible (see Definition 8). The obtained contradiction shows that each of x_{r+1}, z_j is either an isolated point in $\text{Bd } P_T$ or a vertex of a horizontal elementary segment of $\text{Bd } P_T$.

Claim 5. Neither x_{r+1} nor z_j is a vertex of a vertical effective chord from $\mathcal{F}$.

Consider the vertex x_{r+1} . By Claim 4, x_{r+1} is either an isolated point in $\text{Bd } P_T$ or a vertex of a horizontal elementary segment of $\text{Bd } P_T$. If x_{r+1} is isolated in $\text{Bd } P_T$, then it cannot be a vertex of a vertical effective chord from $\mathcal{F}$. Let x_{r+1} be a vertex of a horizontal elementary segment of $\text{Bd } P_T$, say $[v, x_{r+1}]$. Assume, in order to obtain a contradiction, that x_{r+1} is an end of a vertical chord $t_g \in \mathcal{F}$. Since no two orthogonal chords in F have a common point (see Definition 8), the segment $[v, x_{r+1}]$ is not in T ; i.e., $[v, x_{r+1}]$ is an elementary segment of $\text{Bd } P$. Due to Definition 7, there is a vertical elementary segment, say $[w, x_{r+1}]$, in $\text{Bd } P$. Thus we obtained the existence of two vertical elementary segments of $\text{Bd } P_T$ incident with x_{r+1} (one of them is $[w, x_{r+1}]$ and another is contained in t_g), which is in contradiction to Claim 2.

Now we continue the proof of Lemma 7. Claims 1–4 mean that $[x_{r+1}, z_j]$ is an effective chord of P compatible with the family $\mathcal{L}$. Due to the choice of $\mathcal{F}$, the chord $[x_{r+1}, z_j]$ cannot be included in $\mathcal{F}$. Hence there is at least one vertical effective chord in $\mathcal{F}$ having a vertex in $[x_{r+1}, z_j]$. By Claim 5, no vertical effective chord in $\mathcal{F}$ is incident with x_{r+1} or with z_j . Denote by $t'_i = [z'_i, w_i]$, $i = 1, \dots, m$, all vertical chords of $\mathcal{F}$ having their vertices $z'_1, \dots, z'_m$ in $]x_{r+1}, z_j[$ (see Fig. 5 for an example of $[x_{r+1}, z_j]$ in the case $r + 1 = 3$ and $j = 7$; here unbroken lines denote elementary segments of $\text{Bd } P$, wavy lines denote vertical effective chords of $\mathcal{F}$, and dotted lines denote elementary segments of $\text{Bd } P_{r+1}$ which do not belong to $\text{Bd } P_T$; z_4 and z_5 are isolated points of $\text{Bd } P$).

Since each of $z'_1, \dots, z'_m$ is a vertex of exactly one vertical elementary segment of $\text{Bd } P_T$ (see Claim 3) and each chord t'_i contains no elementary segment of $\text{Bd } P$, the vertices $z'_1, \dots, z'_m$ are isolated points in $\text{Bd } P$. In this situation, the chords

$$[x_{r+1}, z'_1], [z'_1, z'_2], \dots, [z'_m, z_j]$$

are effective for P . Put

$$\mathcal{F}' = (\mathcal{F} \cup \{[x_{r+1}, z'_1], [z'_1, z'_2], \dots, [z'_m, z_j]\}) \setminus \{t'_1, \dots, t'_m\}.$$

It is easily seen that $\mathcal{F}'$ is an admissible family of effective chords of P compatible with $\mathcal{L}$, and $\text{card } \mathcal{F}' = \text{card } \mathcal{F} + 1$. The last is in contradiction to the choice of $\mathcal{F}$. The obtained contradiction shows that the assumption $r < s$ is false. $\square$

Fig. 5.

In summary, we obtain that the Second Renumbering gives

$$0 \leq \mu_i(l_i) \leq 1 \quad \text{for all } i = q + 1, \dots, s.$$

Third Renumbering. It remains to renumber the intervals $l_{s+1}, \dots, l_k$. Due to First Renumbering, these intervals are contained in the set $T = t_1 \cup \dots \cup t_f$. Without loss of generality, we may assume that the intervals $l_{s+1}, l_{s+2}, \dots, l_w$, $w \leq k$, form t_1 , i.e.,

$$]x_{s+1}, z_{s+1}[\cup \dots \cup]x_w, z_w[= t_1 \setminus V(P),$$

such that $z_i = x_{i+1}$, $i = s + 1, \dots, w - 1$ (in the case $s + 1 < w$).

From Definition 7, it follows that each of $z_{s+1}, \dots, z_{w-1}$ is a vertex of some elementary segment $[z_i, u_i]$ of $\text{Bd } P$ orthogonal to t_1 , and each of x_{s+1}, z_w is either an isolated point in $\text{Bd } P_T$ or a vertex of a horizontal elementary segment of $\text{Bd } P_T$. It is easily seen that $\mu_{s+1}(l_{s+1}) = 2$, and $\mu_i(l_i) = 1$ for all $i = s + 2, \dots, w$.

Similarly, we renumber repeatedly the intervals $l_{w+1}, \dots, l_k$ such that for any chord t_i , $i = 2, \dots, f$, there exists exactly one interval l_i contained in t_i and having the respective number $\mu_i(l_i) = 2$, while $\mu_j(l_j) = 1$ for all other intervals l_j in t_i .

Since $f \leq e$, the proof of Theorem 1 is complete. $\square$

7. Auxiliary Results

In this section we continue to study the “supplementary ornament” $Y = L \setminus \text{Bd } P$. Let $\mathcal{L} = \{l_1, \dots, l_k\}$ be a decomposition of Y into pairwise disjoint open intervals.

Lemma 8. $m(P) = \mu_1(l_1) + \dots + \mu_k(l_k)$.

Proof. If a vertex $w \in V(P)$ is not an end of l_i , then, by Definition 6, $m_i(w) = m_{i-1}(w)$. Therefore, for any $i = 1, \dots, k$,

$$\mu_i(l_i) = \sum' [m_i(w) - m_{i-1}(w)],$$

where the sum $\sum'$ is taken over the set of all vertices of P .

By Lemma 2, P_0 has no points of local nonconvexity. Hence $m_0(w) = 0$ for any vertex w of P . In Section 6, it was mentioned that $\text{Bd } P_k = \text{Bd } P$. Therefore, $m_k(w) = m(w)$ for all $w \in V(P)$. Thus

$$\begin{aligned} \sum_1^k \mu_i(l_i) &= \sum_1^k (\sum' [m_i(w) - m_{i-1}(w)]) \\ &= \sum' m_k(w) = \sum' m(w) = m(P). \end{aligned} \quad \square$$

Lemma 9. *The number k of open intervals $l_1, \dots, l_k$ decomposing the set Y is at least $k \geq m - e$, where m is the measure of local nonconvexity of P , and e is the effective number of P .*

Proof. By Theorem 1, the intervals $l_1, \dots, l_k$ can be renumbered by the indices $1, \dots, k$ such that $\mu_i(l_i) \leq 1$, for at least $k - e$ of them. Put $\mathcal{L}_r = \{l_i \in \mathcal{L} : \mu_i(l_i) = r\}$, $r = 0, 1, 2$. Since $0 \leq \mu_i(l_i) \leq 2$ for all $i = 1, \dots, k$, we have $\mathcal{L} = \mathcal{L}_0 \cup \mathcal{L}_1 \cup \mathcal{L}_2$ and

$$\mu_1(l_1) + \dots + \mu_k(l_k) = |\mathcal{L}_1| + 2|\mathcal{L}_2|.$$

By the above, $|\mathcal{L}_2| \leq e$. Now from Lemma 8, it follows that

$$\begin{aligned} k &= |\mathcal{L}_0| + |\mathcal{L}_1| + |\mathcal{L}_2| \geq (|\mathcal{L}_1| + 2|\mathcal{L}_2|) - |\mathcal{L}_2| \\ &= \mu_1(l_1) + \dots + \mu_k(l_k) - |\mathcal{L}_2| \geq m - e. \end{aligned} \quad \square$$

The next lemma contains another estimate for the number k of intervals $l_1, \dots, l_k$.

Denote by α_0 and α_1 , respectively, the numbers of all vertices and all elementary segments of a polygon P . The respective numbers for the polygon P_0 are denoted by β_0 and β_1 (see Section 6 for the definition of P_0).

Lemma 10. *If the set $Y = L \setminus \text{Bd } P$ is represented as a union of k pairwise-disjoint open intervals $l_1, \dots, l_k$, then*

$$k = (\beta_1 - \alpha_1) - (\beta_0 - \alpha_0).$$

Proof. Let us represent β_0 and β_1 as

$$\beta_0 = \gamma_0 + \delta_0, \quad \beta_1 = \gamma_1 + \delta_1,$$

where γ_0, γ_1 denote, respectively, the number of P_0 vertices situated in Y and the number of $\text{Bd } P_0$ elementary segments contained in the closure $\bar{Y}$, and let δ_0, δ_1 denote, respectively, the number of vertices of P_0 and the number of elementary segments of $\text{Bd } P_0$ contained in $\text{Bd } P$. Each interval l_i can be decomposed as

$$l_i =]v_1, v_2[\cup \{v_2\} \cup \dots \cup \{v_{r-1}\} \cup]v_{r-1}, v_r[, \quad r \geq 2,$$

where $[v_i, v_{i+1}]$, $i = 1, \dots, r - 1$, are elementary segments of $\text{bd } P_0$, and $v_1, \dots, v_r$ are vertices of P_0 ; i.e., l_i contains the interiors of $r - 1$ elementary segments of $\text{Bd } P_0$ and $r - 2$ vertices of P_0 . Hence $\gamma_1 - \gamma_0 = k$.

If an elementary segment $[v, w]$ of $\text{Bd } P$ contains in its interior some vertices $x_1, \dots, x_q$ of P_0 , then $[v, w]$ contains $q + 1$ elementary segments of $\text{Bd } P_0$. Hence $\delta_0 - \alpha_0 = \delta_1 - \alpha_1$ and

$$(\beta_1 - \alpha_1) - (\beta_0 - \alpha_0) = (\gamma_1 - \gamma_0) + [(\delta_1 - \alpha_1) - (\delta_0 - \alpha_0)] = k. \quad \square$$

8. Generalized Euler's Relation

The following lemma contains a generalized Euler's relation.

Lemma 11. *For a rectilinear polygon P ,*

$$\alpha_0 - \alpha_1 + c + h' = c' + h,$$

where

α_0 is the number of vertices of P ,

α_1 is the number of elementary segments of $\text{Bd } P$,

c is the number of connected components of $\text{Int } P$,

c' is the number of connected components of P ,

h is the number of formal holes of P ,

h' is the number of topological holes of P .

Proof. Consider the planar graph G , whose vertices and edges are precisely the vertices and the elementary segments of P , respectively. Hence G has α_0 vertices and α_1 edges. Denote by q the number of connected components of $E \setminus G$ (in other words, G partitions E into q open connected domains). A well-known assertion states that

$$q = \alpha_1 - \alpha_0 + p + 1,$$

where p is the number of connected components of G . In our notations, $q = h' + c + 1$, where 1 stands for $\text{Ext } P$.

We shall prove the equality $p = c' + h$ by induction on h . The case $h = 0$ is trivial: $p = c'$. Suppose that the assertion " $p = c' + h$ " is true for all rectilinear polygons with at most $h \leq l - 1$ formal holes ($l \geq 1$), and let P be a rectilinear polygon with l formal holes. Let H be any formal hole of P . Then H is separated from other formal holes or from $\text{Ext } P$ by a connected component Q of $\text{Int } P$. It is possible to draw inside Q an open simple polygonal rectilinear path $]x_1, \dots, x_m[$ such that x_1 belongs to H and x_m belongs either to another formal hole or to $\text{Ext } P$. In any case, the addition of $[x_1, \dots, x_m]$ to $\text{Bd } P$ reduces by one both numbers p and l . By inductive assumption, $p - 1 = c' + l - 1$ and $p = c' + h$. According to all the data mentioned above, $\alpha_0 - \alpha_1 + c + h' = c' + h$. $\square$

Example. In Fig. 1, $\alpha_0 = 32$, $\alpha_1 = 32$, $c = 3$, $c' = 1$, $h = 5$, and $h' = 3$.

9. Minimum Number of Rectangles

The following result was announced without proof in [10] (see also [11]).

Theorem 2. *The minimum number of rectangles dissecting a rectilinear polygon P*

is equal to

$$m + c - h - e,$$

where

- m is the measure of P ,
- c is the number of connected components of $\text{Int } P$,
- h is the number of formal holes of P ,
- e is the effective number of P .

Proof. Let r denote the minimum number of rectangles dissecting a rectilinear polygon P . First, we shall prove the inequality

$$r \leq m + c - h - e.$$

Suppose that $e > 0$, and let $[x_i, z_i]$, $i = 1, \dots, e$, be special chords of P forming an admissible family of maximum cardinality. Denote by P_1 the polygon obtained from P by the addition of $[x_1, z_1]$ to $\text{Bd } P$. Some vertices of P can be situated in $]x_1, z_1[$. Denote them by $v_1, \dots, v_t$, respectively. According to Definition 7, any vertex v_i is an end of some elementary segment from $\text{Bd } P$ orthogonal to $[x_1, z_1]$. Hence the $[x_1, z_1]$ addition to $\text{Bd } P$ leads to the following changes:

- (1) the measure of P at each of the points $x_1, v_1, \dots, v_t, z_1$ decreases by one (see Lemma 4);
- (2) the number $h = h(P)$ of formal holes of P decreases by a number $p (\geq 0)$ and the number $c = c(P)$ of connected components of $\text{Int } P$ increases by a number $q (\geq 0)$ such that $p + q = t + 1$;
- (3) the effective number $e(P)$ decreases by at most one.

Thus

$$m(P_1) = m - t - 2, \quad c(P_1) = c + q,$$

$$h(P_1) = h - p, \quad e(P_1) \geq e - 1,$$

and

$$m(P_1) + c(P_1) - h(P_1) - e(P_1) \leq m + c - h - e.$$

Obviously, the admissibility condition implies that chords $[x_i, z_i]$, $i = 2, \dots, e$, remain effective for P_1 .

After repeated additions of the chords $[x_i, z_i]$, $i = 2, \dots, e$, to $\text{Bd } P_1$, we obtain a rectilinear polygon P_e such that

$$\text{Bd } P_e = \text{Bd } P \cup \left(\bigcup_{i=1}^e [x_i, z_i] \right), \quad V(P_e) = V(P).$$

As before,

$$m(P_e) + c(P_e) - h(P_e) - e(P_e) \leq m + c - h - e.$$

We claim that $e(P_e) = 0$. Indeed, if there were at least one effective chord, l , of P_e , then, according to Definition 7, their family $\{l, [x_i, z_i], i = 1, \dots, e\}$ would be an admissible family of cardinality $e + 1$ for P , which is impossible. Hence

$$m(P_e) + c(P_e) - h(P_e) \leq m + c - h - e.$$

Assume that $m(P_e) > 0$. Then some point u of local nonconvexity of P_e exists. By Lemma 5, there is a simple chord $[u, w]$ of P_e whose addition to $\text{Bd } P_e$ decreases by one the measure $m(u)$ of P_e . Denote by P' the polygon obtained from P_e by the addition of $[u, w]$ to $\text{Bd } P_e$. The addition of $[u, w]$ to $\text{Bd } P_e$ decreases by one the measure of P_e at u . Also, this addition either decreases by one the number $h(P_e)$ or increases by one the number $c(P_e)$. So

$$m(P') + c(P') - h(P') \leq m + c - h - e.$$

After repeated application of the above procedure of a simple chord addition, we obtain some rectilinear polygon $\tilde{P}$ without points of local nonconvexity; i.e., $m(\tilde{P}) = 0$. By Lemma 1, $h(\tilde{P}) = 0$. Similarly to the above,

$$c(\tilde{P}) = m(\tilde{P}) + c(\tilde{P}) - h(\tilde{P}) \leq m + c - h - e.$$

Due to Lemma 2, each connected component of $\text{Int } \tilde{P}$ is an open rectangle. The closures of these open rectangles dissect $\tilde{P}$ (and P) into $c(\tilde{P})$ rectangles. Therefore

$$r \leq c(\tilde{P}) \leq m + c - h - e.$$

Now we prove the opposite inequality. Let Q be a polygon obtained from P by the addition of some line segments to $\text{Bd } P$ such that $\text{Bd } Q$ divides $\text{Int } Q$ into rectangles. Denote, respectively, by γ_0 and γ_1 the number of vertices and elementary segments of the polygon Q . By Lemma 11,

$$\alpha_0 - \alpha_1 + c + h' = c' + h,$$

$$\gamma_0 - \gamma_1 + c(Q) + h'(Q) = c'(Q) + h(Q),$$

for polygons P and Q , respectively. It is easily seen that

$$h(Q) = 0, \quad c'(Q) = c', \quad h'(Q) = h'.$$

Therefore

$$c(Q) = c - h + [(\gamma_1 - \alpha_1) - (\gamma_0 - \alpha_0)].$$

Lemmas 9 and 10 give

$$(\gamma_1 - \alpha_1) - (\gamma_0 - \alpha_0) = (k) \geq m - e.$$

Thus

$$c(Q) \geq c - h + m - e.$$

As Q has no points of local nonconvexity, and connected component of $\text{Int } Q$ is an open rectangle (see Lemma 2). The closures of these open rectangles dissect Q (and P) into $c(Q)$ rectangles. Hence any dissection of P contains at least $m + c - h - e$ rectangles. In particular, $r \geq m + c - h - e$. $\square$

10. Optimum Dissection Algorithm

Theorem 2 not only gives a formula for a minimum number of dissecting rectangles, but also contains an informal description of a dissection algorithm which yields a minimum number of rectangles. Below, we describe this algorithm more precisely and show that its computational complexity is at most $O(n^{3/2} \log n)$.

Let P be any rectilinear polygon in the plane E . We assume that the set $S(P)$ of all elementary segments of P is given by two linear arrays forming, respectively, the topological boundary $\text{bd } P$ and the ornament $\text{Or } P$. Any vertex v of P is given by its Cartesian coordinates (X, Y) . Denote by n the number of vertices of P .

OPTIMUM DISSECTION ALGORITHM:

1. Find the family $\mathcal{M}$ of all effective chords of P .
2. Select in $\mathcal{L}$ some admissible family $\mathcal{M} = \{s_1, \dots, s_e\}$ of maximum cardinality.
3. Form the polygon P_m , adding $M := \bigcup_{i=1}^e s_i$ to $\text{Bd } P$.
4. Dissect P_m into rectangles by drawing, in any order, inside P_m some simple chords which remove the local nonconvexity of P_m at each of its vertices.

We discuss each of Steps 1–4 separately.

1. The construction of the family $\mathcal{L}$ is organized for horizontal and vertical chords apart.

Procedure for finding effective horizontal chords:

- (a) Form the family $\mathcal{H}$ of all horizontal chords $[x, z]$ such that $x, z \in V(P)$ and $]x, z[\subset \text{Int } P$ (see [2]).
- (b) Delete repeatedly from any chord $[x, z]$ for which two vertical elementary segments of the form $[x, v]$, $[x, w]$ exist in $\text{Bd } P$.
- (c) Delete repeatedly from $\mathcal{H}$ each pair of chords $[x, v]$, $[v, w]$ for which the vertex v is not an isolated point in $\text{Bd } P$ and include $[x, w]$ in $\mathcal{H}$.

- (d) Delete repeatedly from $\mathcal{H}$ any chord $[x, z]$ such that at least one of x, z is not an isolated vertex of $\text{Bd } P$ or a vertex of a horizontal elementary segment of $\text{Bd } P$.

From Definition 7, it follows that the repeated execution of (a)–(d) yields the family $\mathcal{H} = \{l_1, \dots, l_p\}$ of all horizontal effective chords of P .

The family $\mathcal{V} = \{m_1, \dots, m_q\}$ of all vertical effective chords of P is constructed similarly.

In both cases a sweep-line technique is used. Hence the family $\mathcal{L} := \mathcal{H} \cup \mathcal{V}$ can be found in $O(n \log n)$ time.

2. For the selection of some admissible family of maximum cardinality in $\mathcal{L}$, we use the appropriate $O(n^{3/2} \log n)$ algorithm from [3]. In [3], the family $\mathcal{L}$ is supposed to have the following property: no two collinear chords in $\mathcal{L}$ have a common point. However, in our case some two collinear chords in $\mathcal{L}$ may have a common point (see Observation 3). Hence for the implementation of step 2 in the dissection algorithm, we do the following:

- (1) Transform (one-to-one) the family $\mathcal{L}$ of effective chords of P into a family $\mathcal{L}'$ of segments such that:
 - (i) a chord $[x, z] \in \mathcal{L}$ is horizontal (vertical) if and only if the corresponding segment $[x', z'] \in \mathcal{L}'$ is horizontal (vertical);
 - (ii) any two orthogonal chords $[x, z], [u, v] \in \mathcal{L}$ have a common point if and only if the corresponding segments $[x', z'], [u', v'] \in \mathcal{L}'$ have a common point; and
 - (iii) no two collinear segments in $\mathcal{L}'$ have a common point.
- (2) Select in $\mathcal{L}'$ an admissible family $\mathcal{M}' = \{s'_1, \dots, s'_e\}$ of maximum cardinality (using [3]),
- (3) Obtain in $\mathcal{L}$ the corresponding family $\mathcal{M} = \{s_1, \dots, s_e\}$ of special chords of P , using the inverse transformation $\mathcal{L}' \rightarrow \mathcal{L}$.

Now we describe the $\mathcal{L} \rightarrow \mathcal{L}'$ transformation. Let

$$l_i = [x_i, v_i], \quad i = 1, \dots, p, \quad m_j = [u_j, w_j], \quad j = 1, \dots, q,$$

be, respectively, all the horizontal and vertical chords in $\mathcal{L}$.

Procedure of $\mathcal{L} \rightarrow \mathcal{L}'$ transformation:

- (a) Find a real number $\varepsilon > 0$ such that $\|x - y\| > 2\varepsilon$ for any two different points x, y in the set

$$\{x_1, \dots, x_p, v_1, \dots, v_p, u_1, \dots, u_q, w_1, \dots, w_q\},$$

where $\|\cdot\|$ denotes the Manhattan metric in E .

- (b) For any horizontal chord $l_i = [x_i, v_i] \in \mathcal{L}$ with the ends

$$x_i = (X, Y), \quad v_i = (Z, Y), \quad X < Z,$$

put $l'_i = [x'_i, v'_i] \in \mathcal{L}'$ with the endpoints

$$x'_i = (X - \varepsilon, Y + \delta_i), \quad v'_i = (Z + \varepsilon, Y + \delta_i),$$

where $\delta_i := \varepsilon i / (p + q + 1)$, $i = 1, \dots, p$.

(c) For any vertical chord $m_j = [u_j, w_j] \in \mathcal{L}$ with the ends

$$u_j = (X, Y), \quad w_j = (X, Z), \quad Y < Z,$$

put $m'_j = [u'_j, w'_j] \in \mathcal{L}'$ with the endpoints

$$u'_j = (X + \delta_j, Y - \varepsilon), \quad w'_j = (X + \delta_j, Z + \varepsilon),$$

where $\delta_j := \varepsilon j / (p + q + 1)$, $j = 1, \dots, q$.

Example. For family $\mathcal{L}$ represented in Fig. 6(a), the corresponding family $\mathcal{L}'$ is shown in Fig. 6(b).

Because of the $O(n \log n)$ time complexity of $\mathcal{L} \rightarrow \mathcal{L}'$ and $\mathcal{L}' \rightarrow \mathcal{L}$ transformations, we have the $O(n^{3/2} \log n)$ time complexity of step 2 in the dissection algorithm.

3. Step $P \rightarrow P_M$ has $O(n \log n)$ time complexity (see [2] for technique details).

4. The dissection of P_M into rectangles can be organized in the following manner:

- For each point x of the local nonconvexity of P_M , which is either an isolated point in $\text{Bd } P_M$ or a vertex of some horizontal segment of $\text{Bd } P_H$, draw inside P_H a horizontal simple chord $[x, z]$ ($x, z \in \text{Int } P_M$ and $z \in \text{Bd } P_M$).
- Form the polygon P_T , adding to $\text{Bd } P_M$ all the chords obtained in (a).
- For each vertex x of the local nonconvexity of P_T , draw inside P_T a vertical simple chord $[x, v]$ ($x, v \in \text{Int } P_T$ and $v \in \text{Bd } P_T$).

The computational complexity of step 4 is equal to $O(n \log n)$ [2]. According to all the data mentioned above, the dissection algorithm has at most $O(n^{3/2} \log n)$ time complexity.

Fig. 6. An $\mathcal{L} \rightarrow \mathcal{L}'$ transformation.

Acknowledgment

The authors thank the referees for their many helpful comments on an earlier draft of the paper.

References

1. L. Ferrari, P. V. Sancar, and J. Sklansky, Minimal rectilinear partitions of digitized blocks, *Comput. Vision Graphics Image Process.* **28** (1984), 58–71.
2. A. N. Gorpinevich and V. Soltan, Simple algorithms of the partition of rectilinear regions into rectangles in VLSI engineering, *Bull. Acad. Sci. Mold. SSR. Ser. Phys.—Techn. and Math. Sci.* No. 2 (1989), 19–25 (Russian).
3. H. Imai and T. Asano, Efficient algorithm for geometric graph search problems, *SIAM J. Comput.* **15** (1986), 478–494.
4. N. M. Kornienko, G. V. Matveev, N. N. Metelsky and R. I. Tyshkevich, On tiling of polygons, *Izv. Akad. Nauk. BSSR. Ser. Phys.—Math. Sci.* No. 2 (1978), 25–29 (Russian).
5. A. Lingas, The power of non-rectilinear holes. Lecture Notes on Computer Science, Vol. 140. Springer-Verlag, Berlin, 1982, pp. 369–383.
6. W. Lipski, E. Lord, F. Luccio, C. Mugnai, and L. Pagli, On two-dimensional data organization II, *Fund. Inform.* **2** (1979), 245–260.
7. W. Lipski, Finding a Manhattan path and related problems, *Networks*, **13** (1983), 399–409.
8. W. Lipski, An $O(n \log n)$ Manhattan path algorithms, *Inform. Process. Lett.* **19** (1984), 99–102.
9. V. P. Rubtsov, Realization of VLSI topology by rectangles, *Elektronnaya Tehnika. Ser. 3.* No. 6 (1976), 54–61 (Russian).
10. V. Soltan and A. N. Gorpinevich, Minimum partition of a plane region into rectangles, In: *Republican Seminar on Discrete Optimization*, Kiev, 1985, pp. 107–108 (Russian).
11. V. Soltan and A. N. Gorpinevich, Some algorithms of the partition of a plane region into rectangles, In: *Geometrie und Anwendungen. Vorträge der 7. Tagung der Fachsektion Geometrie*. AdW der DDR, Berlin, 1988, pp. 111–112.

Received July 27, 1990, and in revised form November 10, 1991, and March 15, 1992.