ERRATUM

This letter to the editor appeared on the table of contents of Volume 40, Number 11 but was inadvertently left out of the issue.

IS *HELICOBACTER PYLORI* A CAUSE OF NONULCER DYSPEPSIA?

To The Editor:

It remains controversial whether *H. pylori* plays an important role in nonulcer dyspepsia (1–3). We therefore read with interest the recent report by McCarthy et al (4). Patients with nonulcer dyspepsia were treated with three different regimes to eradicate *H. pylori*. Importantly, it was found at one year that patients who were cured of their infection had significantly lower symptom scores than patients with persisting infection. They conclude, as stated in their abstract, that “These results suggest that *Helicobacter pylori* plays an important role in the symptoms of nonulcer dyspepsia.” While the authors may be correct, we believe that the methodological limitations of their study need to be appreciated before drawing any firm conclusions.

One of the major problems in the McCarthy study was the lack of random allocation of treatments. There is now ample empiric evidence that nonrandomized studies yield larger estimates of therapeutic efficacy than properly randomized trials (5), which could in part account for the promising results reported by McCarthy et al. A further point is that patients who fail to respond to curative therapy may be quite different from those who do respond (6). For example, it is conceivable that those with more severe nonulcer dyspepsia symptoms are less likely to respond to anti-*H. pylori* regimens because the drugs induce more gastrointestinal side effects and hence lead to reduced compliance. Merely comparing those who were successfully treated with those who failed treatment is thus potentially misleading. The lack of a placebo control group is of equal concern because we just do not know whether a placebo group would have performed as well as the curative therapy group over the long term (6). Finally, while the authors state that the patients were unaware of their *H. pylori* or treatment status, this cannot be guaranteed without taking proper precautions. Indeed, if the investigators be-

came aware of those who were and were not cured of their infection, this could conceivably have led to them unconsciously influencing the symptom scoring (6).

We thus remain to be convinced that *H. pylori* plays a key role in symptom generation in nonulcer dyspepsia. We believe that only randomized, double-blind, placebo-controlled studies that pay careful attention to trial methodology will resolve the issue.

G. HOLTMANN, MD

University of Essen

Department of Internal Medicine

Division of Gastroenterology

Hufelandstrasse 55

D-45122 Essen, Germany

N. J. TALLEY, MD, PhD

University of Sydney

Clinical Sciences Building

Nepean Hospital

P.O. Box 63

Penrith, NSW, Australia

REFERENCES

1. Talley NJ: A relationship between *Helicobacter pylori* and non-ulcer dyspepsia: Is there enough data to know? *Eur J Gastroenterol Hepatol* 6:567–570, 1994
2. Kemmer TP, Dominguez Munoz JE, Klingel H, Zemmler T, Kuhn K, Mallertheiner P: The association between non-ulcer dyspepsia and *Helicobacter pylori* infection. *Eur J Gastroenterol Hepatol* 6:571–577, 1994
3. Holtmann G, Goebell H, Holtmann M, Talley NJ: Dyspepsia in healthy blood donors: Pattern of symptoms and association with *Helicobacter pylori*. *Dig Dis Sci* 39:1090–1098, 1994
4. McCarthy C, Patchett S, Collins RM, Beattie S, Keane C, O'Morain C: Long-term prospective study of *Helicobacter pylori* in nonulcer dyspepsia. *Dig Dis Sci* 40:114–119, 1995
5. Schulz KF, Chalmers I, Hayes RJ, Altman DG: Empirical evidence of bias. Dimensions of methodological quality associated with estimates of treatment effects in controlled trials. *JAMA* 273:408–412, 1995
6. Talley NJ: A critique of therapeutic trials in *Helicobacter pylori* positive functional dyspepsia. *Gastroenterology* 106:1174–1183, 1994