Erratum

Psychopharmacology (1994) 115:320–324

Caffeine augmentation of electroconvulsive seizures

Andrew Francis, Laura Fochtman

Department of Psychiatry and Behavioral Sciences, SUNY Stony Brook, Stony Brook, NY 11794, USA

Received: 28 September 1993 / Final version: 8 December 1993

On pages 321 and 322 of this paper, Figs. 1 and 3 were unfortunately interchanged. Figures 1 and 3 are reprinted below together with the correct legends.

Fig. 1. Correspondence of behavioral estimates of seizure duration by length of tonic-clonic convulsions (T-C) and EEG determined seizure length. There was a significant increase in seizure length by caffeine ($F = 18.3$, $df = 1$, $***P < 0.002$) but no difference between behavioral and EEG measures of seizure length ($F < 1$, $df = 1$, $P > 0.9$)

Fig. 3. Time course for caffeine augmentation of ECS at various intervals after a fixed dose of 100 mg/kg. The duration of tonic-clonic convulsions (T-C) is expressed as percent increase over vehicle injection at 0 min. A significant increase was detectable at 1 min (Tukey- $t = 6.54$, $**P < 0.01$) and at all intervals afterward except for 1440 min. Seizure duration at 5 min was greater than that at 1 min (Tukey $t = 3.40$, $††P < 0.10$) and at 230 min (Tukey $t = 2.66$, $††P < 0.01$) but not different at intervening periods. The seizure duration at 1440 min was not different from control.