ERRATUM

Valentina Bassareo · Gianluigi Tanda
Paola Petromilli · Corrado Giua · Gaetano Di Chiara

Non-psychostimulant drugs of abuse and anxiogenic drugs activate with differential selectivity dopamine transmission in the nucleus accumbens and in the medial prefrontal cortex of the rat

Psychopharmacology (1996) 124:293–299

After publication of the above article, the authors supplied corrected versions of Figs. 5, 6 and 8, which are now published below.

Fig. 5 A, B Effect of pentylenetetrazole (PTZ) (○ 20 mg/kg IP; □ 10 mg/kg IP) on DA output in dialysates from the prefrontal cortex (A) and from the nucleus accumbens (B). Results are means±SEM of the results obtained in at least four rats. * $P < 0.05$ compared with basal values

Fig. 6 A, B Effect of picrotoxin (PTX) (△ 0.50 mg/kg SC; □ 0.75 mg/kg SC; ○ 1 mg/kg SC) on DA output in dialysates from the prefrontal cortex (A) and from the nucleus accumbens (B). Results are means±SEM of the results obtained in at least four rats. * $P < 0.05$ compared with basal values

Fig. 8 Effect of the 5HT₃ antagonist ICS 205930 (15 and 30 µg/kg SC) on DA output in dialysates from the rat prefrontal cortex administered 1 h before pentylentetrazole (PTZ, *top*) 20 mg/kg IP or picrotoxin (PTX, *bottom*) 1 mg/kg SC. Results are means±SEM of the results obtained in at least four rats. ○ Saline SC+PTZ 20 mg/kg IP or PTX 1 mg/kg SC; △ ICS 205930 15 µg/kg SC+PTZ 20 mg/kg IP; □ ICS 30 µg/kg SC+PTZ 20 mg/kg IP or PTX 1 mg/kg SC. * $P < 0.05$ compared with saline+PTZ or saline+PTX