ERRATUM

Boris V. Safronov · Werner Vogel

Modulation of delayed rectifier K⁺ channel activity by external K⁺ ions in *Xenopus* axon

Pflügers Arch – Eur J Physiol (1995) 430:879–886

On page 881 of the above article parts of the lettering of Figure 2C disappeared in printing. Figure 2 is reprinted below.

Fig. 2A–C Voltage dependence of K⁺ conductance in Ringer and high-K_o solutions. **A** Activity of a single potassium intermediate channel, I-channel, at -40 mV in Ringer solution. **B** Single-channel current/voltage (*i/E*) relationship of unitary K⁺ currents in Ringer solution. Each point represents the average of the mean current amplitudes from 6–9 outside-out patches. Error bars represent the standard error of the mean (SEM). **C** Potential dependence of K⁺ conductance in external Ringer (open symbols) and high-K_o (filled symbols) solutions. Each point was obtained by dividing the peak K⁺ current amplitude shown in Fig. 1B by the driving force under the assumption that the equilibrium potential for K⁺ ions, $E_K = -89$ mV for Ringer solution and $E_K = 0$ mV for high-K_o. The data are fitted by the sum of two Boltzmann functions. The conductance components activated at more negative potentials were fitted with parameters, voltage of half-maximal activation, $E_{50} = -75$ mV and slope factor $k = 2.6$ mV for high-K_o and $E_{50} = -64$ mV and $k = 5.3$ mV for external ringer solution and are shown as dotted lines