Feasibility Testing for Systems of Real Quadratic Equations*

Alexander I. Barvinok

Department of Mathematics, Royal Institute of Technology,
S-100 44 Stockholm, Sweden
barvinok@math.kth.se

Abstract. We consider the problem of deciding whether a given system of quadratic homogeneous equations over the reals has nontrivial solution. We design an algorithm which, for a fixed number of equations, uses a number of arithmetic operations bounded by a polynomial in the number of variables only.

1. Introduction

Let $G_i = \langle x, \Psi_i x \rangle$, $i = 1, \dots, m$, be a family of quadratic forms on $\mathbb{R}^n$, so Ψ_i , $i = 1, \dots, m$, are $n \times n$ real square symmetric matrices and $\langle \cdot, \cdot \rangle$ is the standard scalar product in $\mathbb{R}^n$. Let $S^{n-1} = \{x \in \mathbb{R}^n, \|x\| = 1\}$ be the unit sphere. We denote by $\|\Psi\|$ the usual norm of Ψ : $\|\Psi\| = \max\{\|\Psi(x)\|, x \in S^{n-1}\}$. We consider the following problem:

(1.1) Problem. Find whether there exists an $x \in S^{n-1}$ such that

$$G_1(x) = \dots = G_m(x) = 0.$$

Without loss of generality we assume that $\|\Psi_i\| \leq \frac{1}{2}$ for $i = 1, \dots, m$.

In other words we are interested in whether a given family of projective quadrics has nonempty intersection. We study the computational complexity of this problem. If $m = 1$, then Problem (1.1) has no solution if and only if the form G_1

* A preliminary version of this paper with a weaker result appeared in the *Proceedings of STOC'92*. This research was supported by the Mittag-Leffler Institute and KTH, Stockholm.

is definite. In this case the Sylvester criterion provides a polynomial algorithm. If $m = 2$, then the Toeplitz–Hausdorff theorem (see, for example, [11]) can be used to design a polynomial-time algorithm for the “generic” forms G_i . No such results seem to be known for $m = 3$. In this paper we prove the following main result.

(1.2) Theorem. *Assume that m is fixed. Then, for any $n \in \mathbb{N}$ and any quadratic forms $G_1, \dots, G_m: \mathbb{R}^n \rightarrow \mathbb{R}$, Problem (1.1) can be solved using a number of arithmetic operations which is polynomial in n .*

Problem (1.1) is universal in a class of semialgebraic problems since an arbitrary system of polynomial equations and inequalities over the field $\mathbb{R}$ can be reduced to Problem (1.1). Of course, the number m of quadratic forms will be large in general. Usually, algorithms in real algebraic geometry have a complexity which is exponential in the number of variables (for an exposition of algorithmic problems in real algebraic geometry and the history of the subject see, for example, [14]–[16]). Theorem (1.2) allows the distinction of “simple” systems of polynomial equations and inequalities, namely, those which can be reduced to a few quadratic equations. It also inspires a hope that other algorithmic problems of algebraic geometry involving intersections of a small number of quadrics can be solved polynomially.

As the main tool to solve (1.1) we study the following optimization problem.

(1.3) Optimization Problem. Let $F_i = \langle x, \Phi_i x \rangle$, $i = 1, \dots, k$, be positive definite quadratic forms on $\mathbb{R}^n$. Find

$$l = \max \left\{ \prod_{i=1}^k F_i(x) : x \in S^{n-1} \right\}.$$

Without loss of generality we assume that $F_i(x) \geq \frac{1}{2}\|x\|^2$ for $i = 1, \dots, k$.

Putting $k = 2m$, $F_{2i-1}(x) = \|x\|^2 - G_i(x)$, $F_{2i}(x) = \|x\|^2 + G_i(x)$ for $i = 1, \dots, m$ for Problem (1.1) we conclude that

$$\max \left\{ \prod_{i=1}^k F_i(x) : x \in S^{n-1} \right\} = \begin{cases} 1 & \text{if the forms } G_1, \dots, G_m \text{ have a} \\ & \text{common nontrivial zero,} \\ l < 1 & \text{if the forms } G_1, \dots, G_m \text{ do not} \\ & \text{have a common nontrivial zero.} \end{cases}$$

The main part of this paper deals with Problem (1.3) and only in Section 4 do we consider the source problem (1.1) which initiates and justifies the study of (1.3). In Section 2 we characterize the optimal value l of (1.3) by constructing a univariate polynomial P of degree $O(n^k)$ such that $P(l) = 0$. In Section 3 we design a polynomial algorithm for (1.3) when k is fixed and the forms $F_1, \dots, F_k$ are in a

general position. We also construct a polynomial algorithm for pushing forms into general position.

By arithmetic operations we mean addition, subtraction, multiplication, division, and comparison of real numbers.

2. A Polynomial Equation for the Maximal Value

Here we prove the following main result.

(2.1) Theorem. *Assume that k is fixed. Then, for any given $n \in \mathbb{N}$ and any given quadratic forms $F_1, \dots, F_k: \mathbb{R}^n \rightarrow \mathbb{R}$, a univariate nonzero polynomial $P(z)$ of degree not more than $(k+1) \cdot n^k$ such that $P(l) = 0$, where l is the solution of (1.3), can be computed. To do that a number of arithmetic operations which is polynomial in n (the degree of this polynomial is linear in k^2) can be used.*

Let I denote the identity $n \times n$ matrix. Consider an expansion

$$(2.2) \quad \det^{-1/2} \left(I - \sum_{i=1}^k t_i \Phi_i \right) = 1 + \sum_{0 \leq m_1, \dots, m_k} q(m_1, \dots, m_k) \cdot t_1^{m_1} \cdots t_k^{m_k}$$

in a small neighborhood of the point $t_1 = \dots = t_k = 0$. Our first lemma deals with the geometric meaning of the coefficients $q(m_1, \dots, m_k)$.

Let $\Gamma(z) = \int_0^{+\infty} x^{z-1} \exp\{-x\} dx$ be the usual Gamma function and let ds be the measure on the sphere S^{n-1} .

(2.3) Lemma. *The following identity for the coefficients of (2.2) holds:*

$$q(m_1, \dots, m_k) = \pi^{-n/2} \cdot \frac{\Gamma(m_1 + \dots + m_k + n/2)}{2 \cdot m_1! \cdots m_k!} \int_{S^{n-1}} F_1^{m_1}(s) \cdots F_k^{m_k}(s) ds.$$

Proof. Put $G(x) = F_1^{m_1}(x) \cdots F_k^{m_k}(x)$. For $r > 0$ set $S(r) = \{x: \|x\| = r\}$ and $\psi(r) = \int_{S(r)} G(s) ds$, where ds is the measure on the sphere $S(r)$ induced from $\mathbb{R}^n$. Since $G(x)$ is homogeneous of degree $2m = 2(m_1 + \dots + m_k)$ we have

$$\psi(r) = \psi(1) \cdot r^{n+2m-1}.$$

Therefore we have

$$\begin{aligned} \int_{\mathbb{R}^n} G(x) \cdot \exp\{-\|x\|^2\} dx &= \psi(1) \cdot \int_0^{+\infty} r^{n+2m-1} \exp\{-r^2\} dr \\ &= \frac{1}{2} \cdot \psi(1) \cdot \Gamma\left(m + \frac{n}{2}\right). \end{aligned}$$

The left-hand side integral is equal to

$$\frac{\partial^m}{\partial t_1^{m_1} \cdots \partial t_k^{m_k}} \int_{\mathbb{R}^n} \exp\{-\|x\|^2 + t_1 F_1(x) + \cdots + t_k F_k(x)\} dx \Big|_{t_1 = \cdots = t_k = 0}.$$

To compute the last integral the well-known formula for the integral over $\mathbb{R}^n$ of the exponential function of a quadratic form can be used (see, for example, [10]). So we have

$$\int_{\mathbb{R}^n} \exp\{-\|x\|^2 + t_1 F_1(x) + \cdots + t_k F_k(x)\} dx = \pi^{n/2} \cdot \det^{-1/2} \left(I - \sum_{i=1}^k t_i \Phi_i \right).$$

Finally we obtain

$$q(m_1, \dots, m_k) = \pi^{-n/2} \cdot \frac{\Gamma(m_1 + \cdots + m_k + n/2)}{2 \cdot m_1! \cdots m_k!} \psi(1),$$

and the proof follows. $\square$

We also need the following result which is known in many different forms.

(2.4) Lemma. *Let $H: \mathbb{R}^n \rightarrow \mathbb{R}$ be a continuous function which is positive on S^{n-1} . Assume that $\rho: S^{n-1} \rightarrow \mathbb{R}$ is a continuous density such that $\rho(s) > 0$ for all $s \in S^{n-1}$. Then*

$$\lim_{m \rightarrow +\infty} \left(\frac{\int_{S^{n-1}} H^{m+1}(s) \rho(s) ds}{\int_{S^{n-1}} H^m(s) \rho(s) ds} \right) = \max\{H(x): x \in S^{n-1}\}.$$

Proof. For the one-dimensional interval this result is proved, for example, in Section 2, Chapter 5, §1, Ex. 199 of [13]. We omit the proof for S^{n-1} since it is completely analogous. $\square$

(2.5) Corollary. *Let us fix $a_1, \dots, a_k \in \mathbb{N}$ and denote $Q_i = q(a_1 + i, \dots, a_k + i)$. Then, for any $j \in \mathbb{N}$,*

$$\lim_{i \rightarrow +\infty} \frac{Q_{j+i}}{Q_i} = k^{kj} \cdot l,$$

where l is the maximal value in (1.3).

Proof. Put $\rho(s) = F_1^{a_1}(s) \cdots F_k^{a_k}(s)$ and $H(s) = F_1(s) \cdots F_k(s)$ in Lemma (2.4). Then using Lemma (2.3) we deduce that $\lim_{i \rightarrow +\infty} Q_{i+1}/Q_i = k^k \cdot l$. Since $Q_{j+i}/Q_i = \prod_{r=1}^j (Q_{i+r}/Q_{i+r-1})$ the proof follows. $\square$

In the previous version of this paper [2] an analogous relation was used to design an approximate algorithm for solving (1.1).

To prove Theorem (1.2) we need a computational version of one result of Gessel (see Theorem 2 of [5]) on rational power series in few variables.

Let $Z(t_1, \dots, t_k)$ be a polynomial in complex variables $t_1, \dots, t_k$ with constant term 1 and let α be a complex number. Let us consider the expansion

$$Z^{-\alpha}(t_1, \dots, t_k) = 1 + \sum_{0 \leq m_1, \dots, m_k} \zeta(m_1, \dots, m_k) \cdot t_1^{m_1} \cdots t_k^{m_k}$$

in a small neighborhood of the point $t_1 = \dots = t_k = 0$. Then Theorem 2 from [5] asserts that there exist polynomials $r_0(m_1, \dots, m_k), \dots, r_d(m_1, \dots, m_k)$, not all equal to zero, such that

$$(2.6) \quad \sum_{j=0}^d r_j(m_1, \dots, m_k) \cdot \zeta(m_1 + j, \dots, m_k + j) = 0$$

for all $m_1, \dots, m_k$.

In fact, in [5] a more general result is proved not only for polynomials but also for rational functions. We need explicit estimates of d and of the computational complexity of these polynomials r_j . The desired estimates can be easily extracted from the proof in [5], but since [5] does not deal with computational complexity questions we briefly describe its method. We assume that the polynomial Z is given by its coefficients. To compute a polynomial means to compute its decomposition into a sum of monomials.

(2.7) Lemma. *Let us fix k . For any given α and any given polynomial Z such that $\deg Z \leq v$, polynomials $r_j(m_1, \dots, m_k)$, $j = 0, \dots, d$, can be computed such that (2.6) holds, $d \leq (k+1) \cdot v^k$, and $\deg r_j \leq k \cdot (k+1) \cdot v^k$ for all j . To compute these polynomials r_j it is necessary to perform $v^{O(k^2)}$ arithmetic operations.*

Proof. We follow Theorem 2 of [5] converting the proof into an algorithm and adding explicit estimates.

Let us choose $D \in \mathbb{N}$. For $\beta = (\beta_1, \dots, \beta_k) \in \mathbb{N}_0^k$, $j \in \mathbb{N}_0$: $\beta_1 + \dots + \beta_k + k \cdot j \leq D$, let us consider the following family of functions:

$$\left(\frac{\partial^k}{\partial t_1 \cdots \partial t_k} \right)^j \prod_{i=1}^k \left(t_i \cdot \frac{\partial}{\partial t_i} \right)^{\beta_i} Z^{-\alpha}(t_1, \dots, t_k) = U_{j, \beta}(t_1, \dots, t_k) \cdot Z^{-\alpha-D}(t_1, \dots, t_k).$$

Here $U_{j, \beta}$ are polynomials of degree at most $D \cdot v$. These polynomials $U_{j, \beta}$ can be computed in a straightforward way using $(D \cdot v)^{O(k)}$ arithmetic operations. It turns out by counting arguments that if D is chosen so that

$$\binom{k + D \cdot v}{k} < \frac{1}{k} \binom{k + D}{k + 1},$$

then the polynomials $U_{j,\beta}$ are linearly dependent. If $\{c_{j,\beta}\}$ are the coefficients of this dependence, then we put

$$r_j(m_1, \dots, m_k) = \prod_{i=1}^k \frac{(m_i + j)!}{m_i!} \sum_{\beta} c_{j,\beta} (m_1 + j)^{\beta_1} \cdots (m_k + j)^{\beta_k}.$$

Thus we have $\deg r_j \leq D$ for all j . We can rewrite r_j as a sum of monomials in $m_1, \dots, m_k$ using $D^{O(k)}$ arithmetic operations.

Hence the problem is reduced to finding a linear dependence between certain polynomials $U_{j,\beta}$ in k variables of degree less than $D \cdot v$. We can choose $D = k \cdot (k+1) \cdot v^k$, so $d \leq (k+1) \cdot v^k$. Now we obtain the desired estimates. $\square$

We need the following purely technical result on the expansion of the determinant of a matrix of polynomials.

(2.8) Proposition. *Let us fix $k \in \mathbb{N}$. Then, for any given $n \times n$ square matrices $A_0, \dots, A_k$, the expansion of*

$$Z(t_1, \dots, t_k) = \det \left(A_0 + \sum_{i=1}^k t_i \cdot A_i \right)$$

into a sum of monomials in $t_1, \dots, t_k$ can be computed using $n^{O(k)}$ arithmetic operations.

Proof. First we note that the degree of Z does not exceed n . Note that the determinant of an $n \times n$ square matrix can be computed using $O(n^3)$ arithmetic operations. Therefore computing the values of $Z(t_1, \dots, t_k)$ in points

$$(t_1, \dots, t_k) \in [0:n]^k$$

from the resulting system of linear equations using $n^{O(k)}$ arithmetic operations we obtain an explicit decomposition of $Z(t_1, \dots, t_k)$ into a sum of monomials. $\square$

Now we can prove the main result of this section.

Proof of Theorem (2.1). Let us denote

$$Z(t_1, \dots, t_k) = \det \left(I - \sum_{i=1}^k t_i \Phi_i \right).$$

So $Z(t_1, \dots, t_k)$ is a polynomial in $t_1, \dots, t_k$ of degree not more than n . The right-hand side of (2.2) is the expansion of $Z^{-1/2}(t_1, \dots, t_k)$ into a power series in $t_1, \dots, t_k$. By Proposition (2.8) we obtain an explicit decomposition of $Z(t_1, \dots, t_k)$

into a sum of monomials in $t_1, \dots, t_k$. Then by Lemma (2.7) using $n^{O(k^2)}$ arithmetic operations we compute polynomials $r_0(m_1, \dots, m_k), \dots, r_d(m_1, \dots, m_k)$, not all equal to zero, such that

$$\sum_{j=0}^d r_j(m_1, \dots, m_k) \cdot q(m_1 + j, \dots, m_k + j) = 0$$

for all $m_1, \dots, m_k$. Here $d \leq (k + 1) \cdot n^k$.

Let us choose $a_1, \dots, a_k$ such that $r_u(a_1, \dots, a_k) \neq 0$ for some u . Let us put

$$Q_i = q(a_1 + i, \dots, a_k + i), \quad R_j(i) = r_j(a_1 + i, \dots, a_k + i), \quad j = 0, \dots, d, \quad i \in \mathbb{N}.$$

So we have got a polynomial recursion

$$\sum_{j=0}^d R_j(i) \cdot Q_{i+j} = 0 \quad (*)$$

for all $i \in \mathbb{N}$, where R_j are polynomials not all of which are identically zero. Let $g = \max\{\deg R_j, j = 0, \dots, d\}$, so $R_j(i) = \alpha_j \cdot i^g + \text{lower-order terms}$. Divide each summand of $(*)$ by $Q_i \cdot i^g$ as $i \rightarrow +\infty$. Since by Corollary (2.5)

$$\lim_{i \rightarrow +\infty} Q_{i+j}/Q_i = k^{kj} \cdot l^j,$$

we obtain finally the desired polynomial equation:

$$\sum_{j=0}^d \alpha_j \cdot k^{kj} \cdot l^j = 0.$$

So we put $P(z) = \sum_{j=0}^d \alpha_j \cdot k^{kj} \cdot z^j$. Since by Lemma (2.7) we have that

$$\deg r_j \leq k \cdot (k + 1) \cdot n^k$$

for all j , we get the desired estimate of the complexity of the algorithm. $\square$

Remark. In the proof above we show that for a certain choice of $a_1, \dots, a_k$ the sequence $q(a_1 + i, \dots, a_k + i)$ is polynomially recursive. In fact, this sequence is polynomially recursive for any $a_1, \dots, a_k$ (see [12]). However, known bounds on the degree of the resulting polynomial equation are much worse than for a sequence with a “generic” starting point.

Example. If $k = 1$, then l is the maximal eigenvalue of the matrix Φ_1 . Then we have $\chi(l) = 0$, where χ is the characteristic polynomial of degree not more than n .

3. Maximum in a General Position

Here we consider the case of “general position” in (1.3). We begin with the following standard result.

(3.1) Lemma. *Let $x \in S^{n-1}$ be a point where the maximum l in (1.3) is attained. Then for some positive $t_1, \dots, t_k$ the following equation holds:*

$$\left(I - \sum_{i=1}^k t_i \Phi_i \right) x = 0.$$

Proof. Note that the maximum of $H = \sum_{i=1}^k \ln F_i$ on S^{n-1} is also attained in x . Thus for the differential dH we get

$$\sum_{i=1}^k \frac{1}{F_i(x)} \Phi_i(x) = \lambda \cdot x$$

for some $\lambda \in \mathbb{R}$. Applying $\langle \cdot, x \rangle$ to both sides of the relation we deduce that $\lambda = k$. $\square$

It is known that in the space of real symmetric $n \times n$ matrices the set of matrices of corank r is a real analytic submanifold of codimension $r(r+1)/2$ (see, for example, the corollary on p. 994 of the English translation of [1]). From this it can be derived that, for k symmetric $n \times n$ matrices $\Phi_1, \dots, \Phi_k$ in general position, the following condition holds:

$$\text{rank} \left(I - \sum_{i=1}^k t_i \cdot \Phi_i \right) \geq n - \frac{\sqrt{1+8k}-1}{2}$$

for all $t_1, \dots, t_k \in \mathbb{R}$.

The words “in general position” mean that the last inequality holds for all matrices from an open dense set in the vector space of all k -tuples $(\Phi_1, \dots, \Phi_k)$ of symmetric $n \times n$ matrices. In fact, for us it is essential that the corank of a linear combination cannot be greater than a certain function in k alone. We say that $\Phi_1, \dots, \Phi_k$ are in general position if

$$(3.2) \quad \text{rank} \left(I - \sum_{i=1}^k t_i \cdot \Phi_i \right) \geq n - f(k)$$

for all $t_1, \dots, t_k \in \mathbb{R}$, where $f(k)$ is a certain function such that

$$f(k) \geq \frac{\sqrt{1+8k}-1}{2}.$$

For example, $f(k) = k$ can be chosen.

Instead of Problem (1.3) we now consider the following “yes-or-no” problem.

(3.3) Problem. For given $a, \varepsilon \in \mathbb{R}$, decide whether $|l - a| < \varepsilon$, where l is the maximal value in (1.3).

The idea of the following result was suggested by A. Megretsky.

(3.4) Theorem. Assume that k is fixed. Then, for given quadratic forms $F_1, \dots, F_k: \mathbb{R}^n \rightarrow \mathbb{R}$ such that (3.2) holds and any given $a, \varepsilon \in \mathbb{R}$, Problem (3.3) can be solved using a number of arithmetic operations which is polynomial in n .

Proof. Let us denote $H(x) = \prod_{i=1}^k F_i(x)$. Put

$$\mathcal{A} = \left\{ x \in S^{n-1} : \left(I - \sum_{i=1}^k t_i \cdot \Phi_i \right) x = 0 \text{ for some } t_1, \dots, t_k \right\}.$$

By Lemma (3.1) it follows that $l = \max\{H(x) : x \in \mathcal{A}\}$, whereas by (3.2) we deduce that $\mathcal{A}$ is a semialgebraic set of dimension not more than $k + f(k)$. Here it is essential that $\dim \mathcal{A}$ is bounded by a function in k alone. We construct a decomposition of the set $\mathcal{A}$ into a union of (possibly intersecting) semialgebraic sets $\{\mathcal{B}_m : m \in M\}$ called *pieces* such that, for each piece $\mathcal{B}_m$, the problem

(3.4.1) given $b \in \mathbb{R}$, decide whether $H(x) > b$ for some $x \in \mathcal{B}_m$

reduces to solving a system of algebraic equations and inequalities in at most $k + f(k)$ variables. The number card M of such pieces is bounded by a polynomial in n . The answer to Problem (3.3) is “yes” if for some $m \in M$ and $b = a - \varepsilon$ the answer to Problem (3.4.1) is “yes” and for all $m \in M$ and $b = a + \varepsilon$ the answer to Problem (3.4.1) is “no.”

An index $m \in M$ consists of a number $r \in \mathbb{N}$ such that $n - f(k) \leq r < n$ and of a pair (I, J) , where $I, J \subset \{1, \dots, n\}$: $\text{card } I = \text{card } J = r$. For $t = (t_1, \dots, t_k)$ let us denote the matrix $I - \sum_{i=1}^k t_i \cdot \Phi_i$ by $\Phi(t)$ and its $r \times r$ submatrix with row indices in I and column indices in J by $\Phi(t; I, J)$. Put

$$T_m = \{t = (t_1, \dots, t_k) \text{ such that all } (r+1) \times (r+1) \text{ minors of } \Phi(t) \text{ are equal to 0 and } \det \Phi(t; I, J) \neq 0\}.$$

Then define

$$\mathcal{B}_m = \{x \in S^{n-1} : \Phi(t)x = 0 \text{ for some } t \in T_m\}.$$

Now we can design the desired system of polynomial equations and inequalities for solving (3.4.1). To simplify notation we assume that the nonsingular submatrix $\Phi(t; I, J)$ occupies the upper left-hand side corner of the matrix $\Phi(t)$. Let us denote

by $u_j(t), j = r + 1, \dots, n$, the vector consisting of the first r entries of the j th column of $\Phi(t)$. Finally put

$$x_j(t) = \begin{cases} -\Phi(t; I, J)^{-1} u_j(t) & \text{for the first } r \text{ coordinates,} \\ 1 & \text{for the } j\text{th coordinate,} \\ 0 & \text{elsewhere.} \end{cases}$$

Then

$$\mathcal{B}_m = \left\{ \sum_{j=r+1}^n \lambda_j x_j(t), \lambda_j \in \mathbb{R}, t \in T_m, \left\| \sum_{j=r+1}^n \lambda_j x_j(t) \right\|^2 = 1 \right\}.$$

Using Proposition (2.8) we obtain an explicit representation of the entries of $\Phi(t; I, J)^{-1}$ as rational functions in $t_1, \dots, t_k$. Now it is clear that (3.4.1) is written as a system of polynomial equations and inequalities in at most $k + f(k)$ variables $t_1, \dots, t_k, \lambda_{r+1}, \dots, \lambda_n$. Since the degree of these equations and inequalities is $O(n)$ and their number is polynomial in n when k is fixed then (see, for example, [14] and [15]) it follows that Problem (3.4.1) can be solved using a number of arithmetic operations which is polynomial in n (the degree of this polynomial is linear in the number of variables, i.e., in $k + f(k)$). Since $\text{card } M \leq n \cdot n^{2 \cdot f(k)}$ we have reduced the initial problem (3.3) to a set of problems of type (3.4.1) whose cardinality is bounded by a certain polynomial in n . $\square$

Now we describe a way to disturb effectively given matrices $\Phi_i \mapsto \hat{\Phi}_i$ to ensure (3.2) with $f(k) = k$. Here we basically follow [7] although we present a weaker construction (in [7] a sharp bound for $f(k)$ is achieved).

(3.5) Theorem. *Assume that k is fixed. Then, for any given symmetric $n \times n$ matrices $\Phi_1, \dots, \Phi_k$ and any given $\varepsilon > 0$, $n \times n$ matrices $\hat{\Phi}_1, \dots, \hat{\Phi}_k$ such that condition (3.2) holds with $f(k) = k$ and $\|\Phi_i - \hat{\Phi}_i\| < \varepsilon$ can be constructed using a number of arithmetic operations which is polynomial in n . (The degree of this polynomial is linear in k .)*

First we reduce the problem to the following one, written in symmetric form.

(3.5.1) Problem. Given real symmetric matrices $A_0, \dots, A_k$ and $\varepsilon > 0$ find symmetric matrices $\hat{A}_i, i = 0, \dots, k$, such that $\|A_i - \hat{A}_i\| < \varepsilon$ for all i and

$$\text{rank} \left(\sum_{i=0}^k t_i \cdot \hat{A}_i \right) \geq n - k$$

for all complex $t_0, \dots, t_k$, not all of which are equal to 0.

If (3.5.1) can be solved in polynomial time, then Theorem (3.5) is proved. One has to choose $A_0 = I$, $A_i = \Phi_i$, $i = 1, \dots, k$, and then put $\hat{\Phi}_i = G' \hat{A}_i G$, where G is a nondegenerate matrix such that $G' \hat{A}_0 G = I$ and $\hat{A}_i$ are computed with regard to $\varepsilon/2$ (we assume that $\varepsilon < \frac{1}{2}$).

Let B_i , $i = 0, \dots, k$, be the following diagonal matrices:

$$B_i(j, j) = j^i.$$

Then for the family B_i , $i = 0, \dots, k$, condition (3.2) obviously holds with $f(k) = k$. We construct the desired deformation of A_i using B_i .

(3.6) Lemma. *There exist not more than $N = n^{O(k)}$ different numbers $z \in \mathbb{C}$ such that*

$$\text{rank} \left(\sum_{i=0}^k t_i \cdot (A_i + z \cdot B_i) \right) < n - k$$

for some $t_0, \dots, t_k$, not all of which are equal to 0.

Proof. Let us consider two complex projective spaces $\mathbf{P}^k = \{t = (t_0 : t_1 : \dots : t_k)\}$, $\mathbf{P}^1 = \{z = (z_0 : z_1)\}$ and the algebraic variety

$$V = \left\{ (t, z) \in \mathbf{P}^k \times \mathbf{P}^1 : \text{rank} \left(\sum_{i=0}^k t_i \cdot z_1 \cdot A_i + t_i \cdot z_0 \cdot B_i \right) < n - k \right\}$$

together with the projection $pr: V \rightarrow \mathbf{P}^1$, $(t, z) \mapsto z$. The image $pr(V)$ is a certain subvariety in $\mathbf{P}^1$ such that the point $(1 : 0)$ does not belong to the image. Therefore $pr(V)$ is a finite set in $\mathbf{P}^1$ and the number of points in $pr(V)$ does not exceed the number of irreducible components of V . Note that V can be defined by $O(n^{2k})$ polynomial equations of degree not more than n in $2k + 2$ variables $w_{ij} = t_i \cdot z_j$ of $\mathbf{P}^{2k+1}$. To estimate the number of irreducible components of V the results of [6] and [3] (see also [8]) can be used. $\square$

Proof of Theorem (3.5). We design an algorithm for Problem (3.5.1). For a given ε choose sufficiently small $\delta > 0$ such that $\|\delta \cdot B_i\| < \varepsilon$ for $i = 0, \dots, k$. Then let us put consecutively $z = 0, \delta/N, 2 \cdot \delta/N, \dots, \delta$, $\hat{A}_i = A_i + z \cdot B_i$, where N is an upper bound from Lemma (3.6). Note that, for any given z , condition (3.2) can be tested using a number of arithmetic operations which is polynomial in n , since it reduces to solving systems of polynomial equations in a fixed number of variables. By Lemma (3.6) it follows that for at least one z from these N the matrices $\hat{A}_i$ are desired. Another way to get such a z is to use a quantifier elimination method (see [15] and [16]) which has polynomial complexity since the number of variables is fixed. $\square$

4. Feasibility Testing

Now we turn to Problem (1.1) and prove the main result of this paper.

Proof of Theorem (1.2). Our algorithm is the following. First we construct the forms $F_1, \dots, F_k$ as in Section 1. Then we have to check whether $l = 1$ where l is the solution of (1.3). Let us construct the polynomial P as in Theorem (2.1). If P does not vanish on 1, then $G_1, \dots, G_m$ have no common nontrivial root and we are done; thus we may assume that $P(1) = 0$. Then we find a number $\delta > 0$ such that $|\alpha_i - \alpha_j| > \delta$ for any two different real roots of the polynomial P . To do this we divide P by g.c.d. $(P(z), dP/dz)$, reducing to the case of polynomial without multiple roots, and then estimate δ using the usual discriminant argument (see, for example, [4]). To compute such a δ it is necessary to perform a number of arithmetic operations which are polynomial in $\deg P$, and therefore are polynomial in n . Then using Theorem (3.5) we construct a perturbation $F_i \mapsto \hat{F}_i$, $i = 1, \dots, k$, such that $|l - \hat{l}| < \delta/2$, where

$$\hat{l} = \max\{\hat{F}_1(x) \cdots \hat{F}_k(x) : x \in S^{n-1}\}.$$

Finally, using Theorem (3.4) we check whether $|\hat{l} - 1| < \delta/2$. If the inequality holds, then there exists a common nontrivial root of $G_1, \dots, G_m$. Otherwise these forms have no common nontrivial root. $\square$

We conclude the paper with two remarks.

Our algorithm is designed for arbitrary real data. If the forms $G_1, \dots, G_m$ are given by their *rational* coefficients, then it can be checked that the size of all numbers involved in the algorithm is bounded by a polynomial in the input size and thus our algorithm is strongly polynomial in the number of variables.

Theorem (1.2) gives us $n^{O(m^2)}$ as an upper bound for the complexity of an algorithm. Grigor'ev told the author that using ideas from [9] an estimate $n^{O(m)}$ can be achieved. He also noted that an estimate $O(\log^m n)$ for the parallel complexity can be achieved.

Acknowledgments

I am grateful to A. M. Vershik who has introduced me to real algebraic geometry and to N. Mnëv, A. Megretsky, and D. Grigor'ev for many stimulating discussions and valuable suggestions. I am indebted to M.-F. Coste-Roy for her remarks and interest and to anonymous referees for their recommendations.

References

1. A. A. Agrachev, Topology of quadratic maps and Hessians of smooth maps, *Itogi Nauki i Tekhniki, Seriya Algebra, Topologiya, Geometriya*, **26** (1988), 85–124 (translated in *Journal of Soviet Mathematics*, **49**(3) (1990), 990–1013).

2. A. I. Barvinok, Feasibility testing for systems of real quadratic equations, in: *Proceedings of the 24th Symposium on the Theory of Computing*, ACM Press, New York, 1992, pp. 126–132.
3. A. L. Chistov, Algorithm of polynomial complexity for factoring polynomials and finding the components of varieties in subexponential time, *Zapiski Nauchnykh Seminarov LOMI*, **137** (1984), 124–188 (translated in *Journal of Soviet Mathematics*, **34**(4) (1986), 1838–1882).
4. G. E. Collins and R. Loos, Real zeros of polynomials, in: *Computer Algebra. Symbolic and Algebraic Computation* (B. Buchberger, G. E. Collins, and R. Loos, eds.), Springer-Verlag, New York, 1982, pp. 83–94.
5. I. M. Gessel, Two theorems on rational power series, *Utilitas Mathematica*, **19** (1981), 247–254.
6. D. Yu. Grigor'ev, Factorization of polynomials over a finite field and the solution of systems of algebraic equations, *Zapiski Nauchnykh Seminarov LOMI*, **137** (1984), 20–79 (translated in *Journal of Soviet Mathematics*, **34**(4) (1986), 1762–1803).
7. D. Yu. Grigor'ev, Private communication, 1992.
8. D. Yu. Grigor'ev and A. L. Chistov, Fast decomposition of polynomials into irreducible ones and the solution of systems of algebraic equations, *Doklady Akademii Nauk SSSR*, **275**(6) (1984), 1302–1306 (translated in *Soviet Mathematics. Doklady*, **29**(2) (1984), 380–383).
9. D. Yu. Grigor'ev and N. N. Vorobjov, Solving systems of polynomial inequalities in subexponential time, *Journal of Symbolic Computation*, **5**(1–2) (1988), 37–64.
10. J. Hadamard, *Cours D'Analyse*, vol. 1, Hermann, Paris, 1927.
11. P. R. Halmos, *A Hilbert Space Problem Book*, Van Nostrand, London, 1967.
12. L. Lipshitz, The diagonal of a D -finite series is D -finite, *Journal of Algebra*, **113** (1988), 373–378.
13. G. Pólya and G. Szegő, *Aufgaben und Lehrsätze aus der Analysis*, Vol. 1, Dover, New York, 1945.
14. J. Renegar, On the computational complexity and geometry of the first order theory of the reals. Part 1. Introduction. Preliminaries. The geometry of semi-algebraic sets. The decision problem for the existential theory of the reals, *Journal of Symbolic Computation*, **13**(3) (1992), 255–299.
15. J. Renegar, On the computational complexity and geometry of the first order theory of the reals. Part 2. The general decision problem. Preliminaries for quantifier elimination, *Journal of Symbolic Computation*, **13**(3) (1992), 301–327.
16. J. Renegar, On the computational complexity and geometry of the first order theory of the reals. Part 3. Quantifier elimination, *Journal of Symbolic Computation*, **13**(3) (1992), 329–352.

Received November 9, 1991, and in revised form September 14, 1992.