In reply:—We share Gozum's enthusiasm that computer diagnostic assistants (CDAs) may be an important addition to the practice of internal medicine in the future. In the proper context, the computer might expand knowledge, increase efficiency, improve quality, and lower cost. The article by Bacchus et al.¹ demonstrated that although the improvement in interns' diagnostic scores after using QMR was statistically significant, they felt the clinical improvement did not warrant the time expenditure for training. The limitations of space did not allow us to discuss the potential cost implications of CDAs in our editorial,² but we do not think it is clear that CDAs will produce cost savings. Cost studies have not been completed. Improvements in diagnostic accuracy may ultimately be cost-effective. On the other hand, improperly entered findings leading to inappropriate diagnoses on the differential might increase use of unnecessary testing. Primary care physicians using CDAs might be able to avoid certain consultations, but there is currently no system to compensate the primary physician for the not-inconsequential amount of time required to use the CDA.

The figures cited for CDA accuracy in the Berner paper³ represent the proportion of the cases in which the correct diagnosis was listed in the *entire list produced by the CDA*. When the correct diagnosis was present in the CDA's database, the correct diagnosis still was ranked only between 6th and 13th on the CDAs' differentials. Increasing the number of diagnoses present in the database is logical. However, although it is obvious that the computer cannot rank a disease that is not in its database, it is not obvious that increasing the number of

diseases included in the database will lead to a proportional increase in accuracy.

In the future, CDAs may play a supportive role in quality assurance and cost containment as suggested by Dr. Gozum. However, from our experience, we believe that the diagnostic accuracy and clinical utility of computer-aided diagnosis are highly dependent on the skill of the user. Laypersons and medical students often are unable to properly select features to enter into the CDA, and the resultant differential may be seriously misleading. We therefore are *strongly* opposed to the idea that CDAs (at this stage of development) might be used by third-party payers or quality assurance workers to second-guess practicing physicians. —**ANDREW S. KANTER, MD, MPH**, Chief Medical Resident, Providence Medical Center, and Acting Instructor, Department of Medicine, University of Washington, Seattle, WA; and **DAWN E. DEWITT, MD, MSc**, Acting Assistant Professor, Department of Medicine, University of Washington, Seattle, WA 98195

REFERENCES

1. Bacchus CM, Quinton C, O'Rourke K, Detsky AS. A randomized crossover trial of Quick Medical Reference (QMR) as a teaching tool for medical interns. *J Gen Intern Med*. 1994;9:616–21.
2. DeWitt DE, Kanter AS. Are computer diagnostic assistants useful tools for clinicians and educators? *J Gen Intern Med*. 1994;9:653–4.
3. Berner ES, Webster GD, Shugerman AA, et al. Performance of four computer based diagnostic systems. *N Engl J Med*. 1994;330:1792–6.

REFLECTIONS

Optimistic lies have such immense therapeutic value that a doctor who cannot tell them convincingly has mistaken his profession. —GEORGE BERNARD SHAW (1856–1950), Irish dramatist and critic. *Misalliance*, Preface