Themenschwerpunkt:

„Neue Trends in der Therapie der portalen Hypertonie“

Editorial

Ph. Gertsch

Since the Russian surgeon Nicolas Eck performed the first portacaval shunts in dogs in 1877, treatment of the hemorrhagic complications of portal hypertension has been a subject of constant and renewed interest for surgeons. Research in this field has resulted in new treatments, and several therapeutic options are now offered to clinicians treating patients with cirrhosis and portal hypertension. Until recently, the only effective treatment for variceal hemorrhage has been portacaval decompression. This therapy was based on the concept that resistance to portal flow was responsible for the elevation of portal pressure and variceal bleeding. However, this proved to be partially true, and the initial studies on portacaval diversion also put into focus hepatic encephalopathy as another important complication related to portal hypertension and portosystemic shunting. Significant progress has been made since the first surgical attempts at solving the problem of portal hypertension, and surgeons have become more critical and aware of the limits of their successes. Based on better pathophysiological understanding of the manifestations of portal hypertension, new treatments have been proposed, which often required more than a decade for their complete assessment.

Selective shunting has been performed with the hope of achieving better results than those obtained with total portal diversion, but long-lasting selective portal decompression has proved difficult to achieve (1). Partial shunting preserving hepatic portal perfusion, as a goal, is still considered preferable to total portal diversion. Time will tell whether the recently introduced small diameter interposition portacaval shunts will maintain their initial hemodynamic advantages on a long term basis (2). The technique for placement of the "Transjugular Intrahepatic Portosystemic Stent Shunts" (TIPS) has improved remarkably and this new treatment of por-

tal hypertension already represents a concrete therapeutic option in many centers around the world. Because of their small diameter, TIPS might achieve partial shunting (3). The long-term results and place of this new treatment remains to be defined.

Control of variceal hemorrhage by direct action on the source of bleeding has been performed since over 50 years either surgically, or endoscopically by injection of sclerosant into varices. Since the advent of fibroscopes in the seventies, sclerotherapy has progressively been adopted in most centers as the primary treatment for active variceal hemorrhage as well as for prevention of future bleeding.

Understanding the importance of the hypercirculatory state in the rise of portal pressure has resulted in the introduction of drugs reducing portal pressure. This has opened new perspectives in the prophylaxis of variceal bleeding (4). Liver transplantation provides a simultaneous and definitive cure of end stage liver insufficiency and portal hypertension, and has been proposed as a therapeutic option in patients with cirrhosis and bleeding varices (5). Variceal hemorrhage does not necessarily mean that the patient has end-stage liver disease. Furthermore, the evolution towards liver insufficiency may differ, as each disease has its own natural history. Therefore liver transplantation, should not be regarded as the optimum therapeutic option in all cases, and other choices may be considered in patients with preserved hepatic function. A recent comparative analysis of selective shunting and liver transplantation highlights the respective merits of both treatments (6).

Treatments for variceal hemorrhage related to liver cirrhosis may be regarded as palliative in most cases. Some patients presenting variceal bleeding may be or may become candidates for liver transplantation. This puts into a new light all palliative measures aimed at controlling variceal hemorrhage as they may offer time to the patient waiting for a new liver. No consensus exists as to whether recurrent variceal hemorrhage in transplant candidates

should be treated by conservative measures, sclerotherapy, or surgical shunt. TIPS may be the best choice (3) and certainly deserves complete investigation. This procedure does not have the drawback of operative portacaval decompression which adds to the technical difficulty and operative blood loss of subsequent liver transplantation (5, 7). Furthermore TIPS might be a technical help to the transplant surgeon in reducing intraoperative bleeding caused by elevated portal pressure. Portal hypertension secondary to liver cirrhosis is related to a wide pattern of underlying conditions. Detailed investigations are necessary to select the most appropriate treatment for each particular case.

The authors of this issue of *Acta Chirurgica Austriaca*, chosen for their expertise and broad knowledge in the field of portal hypertension, will give the readers some perspectives on recent advances and research in this field and help them choose the appropriate investigations and therapies for different situations.

I wish to express my gratitude to the authors for their erudite text covering the major problems related to portal hypertension.

References

- (1) Henderson JM, Gong-Liang J, Galloway J, et al: Portacaval collaterals following distal splenorenal shunt: incidence, magnitude and associated portal perfusion changes. *J Hepatol* 1985;1:649-661.
- (2) Sarafteh IJ, Rypins EB, Raiszadeh M, et al: Serial measurement of portal hemodynamics after partial portal decompression. *Surgery* 1986;100:52-57.
- (3) Ring EJ, Lake JR, Roberts JP, et al: Using Transjugular Intrahepatic Portosystemic Shunts to control variceal bleeding before liver transplantation. *Ann Int Med* 1992;116:304-309.
- (4) Elder CA, Restino MS: Beta-adrenergic antagonists for primary prevention of gastrointestinal hemorrhage in patients with cirrhosis and esophageal varices. *Clin Pharm* 1992;11:337-341.
- (5) Iwatsuki S, Starzl TE, Todo S, et al: Liver transplantation in the treatment of bleeding esophageal varices. *Surgery* 1988;104:697-705.
- (6) Henderson JM, Gilmore TG, Hooks MA, et al: Selective shunt in the management of variceal bleeding in the era of liver transplantation. *Ann Surg* 1992;216:248-255.
- (7) Mazzafe V, Todo S, Tzakis A, et al: Liver transplantation in patients with previous portosystemic shunt. *Am J Surg* 1990;160:111-116.

**ACA-Manuscripts
should be submitted to:**

B. Niederle, M.D.
Editor-in-Chief
c/o Blackwell-MZV
Feldgasse 13,