Gradient of Cosmic-Ray Intensity Perpendicular to the Elliptic Plane.

R. P. KANE

Physical Research Laboratory - Ahmedabad

(*Nuovo Cimento*, **57** B, 36 (1968))

In this communication a serious error has occurred, *viz.* the dates for farthest Northern and Southern heliolatitudes have been taken as June 7 and Dec. 7 whereas these should have been March 7 and September 7. After reanalysis, the Table II therein should read as follows:

TABLE II. — *For climax neutron monitor.*

Year	Northern hemisphere (March centring)			Southern hemisphere (September centring)		
	Correlation coefficient	Gradient (%) over		Correlation coefficient	Gradient (%) over	
		1/8 A.U.	1 A.U.		1/8 A.U.	1 A.U.
1960	0.18	0.3	2.5	0.62	1.1	8.9
1961	0.93	1.8	14.8	— 0.89	— 2.4	— 19.4
1962	— 0.81	— 0.7	— 5.2	— 0.83	— 0.7	— 5.7
1963	0.91	1.7	13.7	— 0.29	— 0.2	— 2.0
1964	0.15	0.1	0.6	— 0.23	— 0.2	— 1.9
1965	— 0.33	— 0.3	— 2.2	— 0.91	— 0.9	— 7.5
1966	0.20	0.5	2.4	— 0.37	— 0.3	— 2.7
1967	— 0.75	— 1.1	— 8.5	0.05	0.1	0.5
1968	0.24	0.3	2.6	0.83	2.0	16.0
Average (1960-68)	0.68	0.3	2.3	— 0.71	— 0.2	— 1.5

Conclusions are roughly the same as given in the previous communication. Analysis of Deep River neutron monitor also shows similar results.