**Approximate Relativistic Equations of Motion
for an Extended Charged Particle
in an Inhomogeneous External Electromagnetic Field.**

P. NYBORG

Mathematics Department, Agricultural College of Norway - Vollebekk

(*Nuovo Cimento*, **31**, 1209 (1964))

For eq. (86), p. 1223, read

$$(86) \quad O_{\mu\varrho} O_{\nu\sigma} \dot{\mu}_{\varrho\sigma} = g N_{\mu\nu}.$$

This equation together with (78) determines the motion of $\mu_{\mu\nu}$.

For eq. (88), read

$$(88a) \quad N_{\mu\nu} = \mu_{\mu\sigma} (O_{\nu\varrho} F_{\varrho\sigma}) - \mu_{\nu\sigma} (O_{\mu\varrho} F_{\varrho\sigma}).$$

Eqs. (86a) and (78) are satisfied by

$$(88b) \quad \begin{cases} \dot{\mu}_{\mu\nu} = g N_{\mu\nu} + R_{\mu\nu}, \\ R_{\mu\nu} = \frac{1}{mc^2} (\mu_{\mu\sigma} U_\nu - \mu_{\nu\sigma} U_\mu) F_\sigma, \end{cases}$$

with $N_{\mu\nu}$ and F_μ given by (88a) and (85) respectively.

As $R_{\mu\nu}$ is proper timelike, the equation of motion (99) for t_μ will not be affected by these changes in (86) and (88).