Singularities and Discontinuities of the Triangle Graph, as a Functions of an Internal Mass.

I. J. R. AITCHISON

High-Energy Physics Group, Cavendish Laboratory - Cambridge

C. KACSER

Physics Department, The University of Maryland - College Park, Md.

(*Nuovo Cimento*, **40 A**, 576 (1965))

The right-hand sides of each of the following equations should be multiplied by a single extra factor of π : eqs. (6a), (6b), (7), (9), (10) and the final equation. (Cf.: *The discontinuities of the triangle graph*, C. KACSER: *Journ. Math. Phys.*, **7**, to appear (1966)).

Comparison of $K^\pm p$ Dispersion Relations with Experiment.

M. LUSIGNOLI, M. RESTIGNOLI and G. VIOLINI

Istituto di Fisica dell'Università - Roma

G. A. SNOW

CERN - Geneva

(*Nuovo Cimento*, **45 A**, 792 (1966))

In ref. ⁽⁵⁾	instead of GIACOMELL read GIACOMELLI
Every time	instead of FOGGART read FROGGATT
At page 793, line 7 in the $X(Y)$ numerator	instead of m_K read m_K^2
At page 797, line 1	instead of vicinity read vicinity
line 6	instead of solution II read solution III
In the table's footnotes	instead of c_4 read c_6
In Table I, line 2	instead of oole read pole
In Table II, at $k = 9.00 \text{ GeV}/c$, $c_3 = -76.40 \pm 0.69$ and $D_- = -1.96 \pm 3.74$.	